



Full wwPDB EM Validation Report ⓘ

Apr 28, 2026 – 10:49 pm BST

PDB ID : 9R9A / pdb_00009r9a
EMDB ID : EMD-53855
Title : CryoEM structure of a membrane protein associated with a contractile injection in *Salmonella enterica* subspecies *salamae*
Authors : Ejaz, R.N.; Tillmann, H.P.; Taylor, N.M.I.; Siborova, M.; Sofos, N.H.
Deposited on : 2025-05-19
Resolution : 2.60 Å(reported)
Based on initial model : .

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	864	 53% 8% 39%
1	C	864	 53% 8% 39%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 17112 atoms, of which 8548 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	B	530	Total	C	H	N	O	S	0	0
			8556	2778	4274	707	780	17		
1	C	530	Total	C	H	N	O	S	0	0
			8556	2778	4274	707	780	17		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



A	W717	T437	GLN	ASN	SER
	N718	M438	ILE	ILE	GLY
	L732	G439	ASN	ASN	LYS
	F733	T440	THR	LEU	ALA
	L734	G443	SER	ILE	ASP
	E739	F444	GLN	LYS	THR
	T745	Y445	ASP	SER	SER
	E777	Q449	THR	ILE	ALA
	I789	H450	MET	GLU	GLU
	L793	Q455	THR	ILE	PRO
B	C796	S456	ALA	HIS	SER
	I797	I457	PRO	PRO	PRO
	Y800	R461	GLN	SER	E197
	L802	L462	LEU	LYS	E212
	R803	V468	GLN	SER	Q222
	L814	G469	THR	THR	T223
	V827	P484	ASP	SER	I224
	Q833	R502	ASP	SER	LEU
	I845	Y503	ALA	ALA	ALA
	A853	R504	ASN	THR	THR
C	V857	K551	LYS	ASN	LYS
	R858	Q559	ILE	THR	LEU
	LYS	T575	LYS	THR	GLU
	GLU	D578	THR	PRO	LYS
	GLY	T579	ILE	PRO	LYS
	ASP	E581	GLN	PRO	GLN
	LYS	V626	PHE	ALA	GLN
	ASP	R634	LYS	SER	PHE
	LYS	G635	T357	SER	ASP
	PRO	N636	D364	GLU	GLN
D	GLY	E581	VAL	VAL	ARG
	GLY	L365	GLU	GLU	SER
	ASP	V626	ASN	ASN	GLN
	LYS	E374	THR	THR	TTR
	LYS	R634	SER	SER	ASN
	PRO	G635	GLU	GLU	GLN
	GLY	N636	GLN	GLN	SER
	GLY	R637	SER	SER	GLU
	GLY	F638	ILE	ILE	LEU
	PRO	Y639	S411	GLY	PHE
E	L647	P412	ASP	ASP	ALA
	L653	I413	THR	THR	GLN
	R657	H414	THR	THR	LEU
	L688	W420	SER	SER	ASN
	E704	D425	PRO	PRO	ILE
	T714	D430	VAL	VAL	ALA
	T714	D430	PRO	PRO	ALA
	T714	W432	THR	THR	THR
	T714	W432	THR	THR	GLY
	T714	W432	THR	THR	PRO

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	239630	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	41	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.332	Depositor
Minimum map value	-0.174	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.03033	Depositor
Map size (Å)	232.95999, 232.95999, 232.95999	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.728, 0.728, 0.728	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.20	0/4394	0.25	0/5976
1	C	0.20	0/4394	0.25	0/5976
All	All	0.20	0/8788	0.25	0/11952

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4282	4274	4272	46	0
1	C	4282	4274	4272	46	0
All	All	8564	8548	8544	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:634:ARG:NH1	1:B:634:ARG:O	2.11	0.83
1:C:634:ARG:O	1:C:634:ARG:NH1	2.11	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:363:GLU:N	1:C:363:GLU:OE2	2.19	0.75
1:B:363:GLU:N	1:B:363:GLU:OE2	2.19	0.75
1:C:212:GLU:N	1:C:212:GLU:OE1	2.25	0.70
1:B:468:VAL:O	1:B:468:VAL:HG12	1.91	0.70
1:C:468:VAL:HG12	1:C:468:VAL:O	1.91	0.69
1:B:777:GLU:N	1:B:777:GLU:OE1	2.26	0.69
1:B:212:GLU:OE1	1:B:212:GLU:N	2.25	0.69
1:C:777:GLU:OE1	1:C:777:GLU:N	2.26	0.69
1:C:374:GLU:OE2	1:C:450:HIS:NE2	2.28	0.66
1:B:374:GLU:OE2	1:B:450:HIS:NE2	2.28	0.65
1:C:445:TYR:OH	1:C:449:GLN:O	2.14	0.64
1:B:717:VAL:HG12	1:B:745:THR:HG21	1.80	0.63
1:B:445:TYR:OH	1:B:449:GLN:O	2.14	0.62
1:B:551:LYS:NZ	1:B:559:GLN:OE1	2.27	0.62
1:C:433:LYS:HD3	1:C:438:MET:HE1	1.83	0.61
1:C:717:VAL:HG12	1:C:745:THR:HG21	1.80	0.61
1:C:714:THR:O	1:C:718:ASN:ND2	2.35	0.60
1:B:714:THR:O	1:B:718:ASN:ND2	2.35	0.60
1:B:433:LYS:HD3	1:B:438:MET:HE1	1.83	0.59
1:B:430:ASP:OD1	1:B:504:ARG:NH1	2.38	0.56
1:B:827:VAL:O	1:B:833:GLN:NE2	2.37	0.56
1:C:382:PHE:CZ	1:C:386:ILE:HD11	2.41	0.56
1:C:551:LYS:NZ	1:C:559:GLN:OE1	2.27	0.56
1:B:382:PHE:CZ	1:B:386:ILE:HD11	2.41	0.55
1:B:575:THR:HG22	1:B:636:ASN:HD21	1.72	0.55
1:B:843:PHE:CE2	1:B:847:ILE:HD11	2.42	0.55
1:C:827:VAL:O	1:C:833:GLN:NE2	2.37	0.55
1:C:575:THR:HG22	1:C:636:ASN:HD21	1.72	0.54
1:C:430:ASP:OD1	1:C:504:ARG:NH1	2.38	0.53
1:B:578:ASP:OD1	1:B:579:THR:N	2.42	0.53
1:B:461:ARG:NH2	1:B:739:GLU:OE2	2.39	0.53
1:C:578:ASP:OD1	1:C:579:THR:N	2.42	0.52
1:C:410:LEU:HD23	1:C:411:SER:N	2.25	0.52
1:C:461:ARG:NH2	1:C:739:GLU:OE2	2.39	0.51
1:C:732:LEU:O	1:C:734:LEU:N	2.44	0.51
1:B:732:LEU:O	1:B:734:LEU:N	2.44	0.51
1:B:410:LEU:HD23	1:B:411:SER:N	2.25	0.51
1:C:789:ILE:HD11	1:C:814:LEU:HD22	1.93	0.50
1:C:443:GLY:HA3	1:C:732:LEU:HD13	1.94	0.50
1:B:789:ILE:HD11	1:B:814:LEU:HD22	1.93	0.49
1:C:853:ALA:O	1:C:857:VAL:HG23	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:443:GLY:HA3	1:B:732:LEU:HD13	1.94	0.49
1:B:853:ALA:O	1:B:857:VAL:HG23	2.13	0.49
1:B:365:LEU:C	1:B:365:LEU:HD23	2.38	0.49
1:B:502:ARG:NH2	1:B:777:GLU:OE2	2.46	0.48
1:B:357:ILE:O	1:B:357:ILE:HG22	2.13	0.48
1:C:502:ARG:NH2	1:C:777:GLU:OE2	2.46	0.48
1:B:365:LEU:HD23	1:B:365:LEU:O	2.14	0.48
1:C:357:ILE:O	1:C:357:ILE:HG22	2.13	0.47
1:C:365:LEU:O	1:C:365:LEU:HD23	2.14	0.47
1:C:420:TRP:CD2	1:C:457:ILE:HD12	2.50	0.47
1:C:365:LEU:HD23	1:C:365:LEU:C	2.38	0.47
1:B:420:TRP:CD2	1:B:457:ILE:HD12	2.50	0.47
1:B:636:ASN:OD1	1:B:638:PHE:N	2.49	0.46
1:C:636:ASN:OD1	1:C:638:PHE:N	2.49	0.46
1:C:581:GLU:OE1	1:C:581:GLU:N	2.44	0.45
1:C:634:ARG:NH1	1:C:636:ASN:O	2.49	0.45
1:B:634:ARG:NH1	1:B:636:ASN:O	2.49	0.45
1:B:462:LEU:HD11	1:B:732:LEU:HD11	1.97	0.45
1:C:469:GLY:HA3	1:C:484:PRO:HG3	1.99	0.45
1:C:468:VAL:O	1:C:468:VAL:CG1	2.62	0.44
1:C:462:LEU:HD11	1:C:732:LEU:HD11	1.97	0.44
1:B:581:GLU:OE1	1:B:581:GLU:N	2.44	0.44
1:C:704:GLU:OE1	1:C:704:GLU:N	2.41	0.44
1:C:626:VAL:HG13	1:C:639:TYR:CE1	2.53	0.44
1:B:469:GLY:HA3	1:B:484:PRO:HG3	1.99	0.43
1:B:626:VAL:HG13	1:B:639:TYR:CE1	2.53	0.43
1:B:796:CYS:SG	1:B:797:ILE:N	2.91	0.43
1:C:797:ILE:O	1:C:803:ARG:NH2	2.45	0.43
1:B:793:LEU:HD23	1:B:793:LEU:C	2.44	0.42
1:C:796:CYS:SG	1:C:797:ILE:N	2.91	0.42
1:B:657:ARG:O	1:B:657:ARG:HG3	2.19	0.42
1:B:413:ILE:HD12	1:B:413:ILE:N	2.34	0.42
1:C:413:ILE:HD12	1:C:413:ILE:N	2.34	0.42
1:B:437:THR:O	1:B:437:THR:HG22	2.19	0.42
1:C:437:THR:O	1:C:437:THR:CG2	2.67	0.42
1:C:800:TYR:N	1:C:801:PRO:HD2	2.35	0.42
1:C:437:THR:O	1:C:437:THR:HG22	2.19	0.42
1:B:800:TYR:N	1:B:801:PRO:HD2	2.35	0.42
1:C:793:LEU:C	1:C:793:LEU:HD23	2.44	0.42
1:C:657:ARG:O	1:C:657:ARG:HG3	2.19	0.41
1:B:455:GLN:N	1:B:455:GLN:OE1	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:414:HIS:ND1	1:C:414:HIS:O	2.53	0.41
1:C:455:GLN:N	1:C:455:GLN:OE1	2.53	0.41
1:B:704:GLU:OE1	1:B:704:GLU:N	2.41	0.41
1:B:437:THR:O	1:B:437:THR:CG2	2.67	0.41
1:B:414:HIS:O	1:B:414:HIS:ND1	2.53	0.41
1:B:468:VAL:O	1:B:468:VAL:CG1	2.62	0.40
1:B:802:LEU:HD22	1:B:802:LEU:N	2.36	0.40
1:C:802:LEU:N	1:C:802:LEU:HD22	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	526/864 (61%)	516 (98%)	10 (2%)	0	100	100
1	C	526/864 (61%)	516 (98%)	10 (2%)	0	100	100
All	All	1052/1728 (61%)	1032 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	476/766 (62%)	469 (98%)	7 (2%)	57	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	476/766 (62%)	469 (98%)	7 (2%)	57	80
All	All	952/1532 (62%)	938 (98%)	14 (2%)	55	80

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	222	GLN
1	B	425	ASP
1	B	440	THR
1	B	647	LEU
1	B	653	LEU
1	B	688	LEU
1	B	845	ILE
1	C	222	GLN
1	C	425	ASP
1	C	440	THR
1	C	647	LEU
1	C	653	LEU
1	C	688	LEU
1	C	845	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	429	GLN
1	B	691	GLN
1	C	429	GLN
1	C	691	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-53855. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 160




Y Index: 134




Z Index: 119


6.3.2 Raw map



X Index: 160



Y Index: 134




Z Index: 119


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) ⓘ


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.03033. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 173 nm³; this corresponds to an approximate mass of 156 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.385 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.18	3.67	3.24

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.18 differs from the reported value 2.6 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-53855 and PDB model 9R9A. Per-residue inclusion information can be found in section [3](#) on page [4](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.03033 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03033).

9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.03033) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9720	<div></div> 0.5230
B	<div></div> 0.9720	<div></div> 0.5230
C	<div></div> 0.9720	<div></div> 0.5230

