



Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2026 – 01:29 pm BST

PDB ID : 9R9T / pdb_00009r9t
Title : [FeFe]-hydrogenase from Nitratidesulfovibrio vulgaris str. Hildenborough at pH 5.81
Authors : Bikbaev, K.; Span, I.
Deposited on : 2025-05-20
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>15%</div> <div>• •</div> </div> </div>
2	B	88	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>15%</div> <div>•</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7646 atoms, of which 3767 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic [Fe] hydrogenase large subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	396	6047	1926	3011	505	570	35	255	5	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	398	TRP	-	expression tag	UNP P07598
A	399	SER	-	expression tag	UNP P07598
A	400	HIS	-	expression tag	UNP P07598
A	401	PRO	-	expression tag	UNP P07598
A	402	GLN	-	expression tag	UNP P07598
A	403	PHE	-	expression tag	UNP P07598
A	404	GLU	-	expression tag	UNP P07598
A	405	LYS	-	expression tag	UNP P07598

- Molecule 2 is a protein called Periplasmic [Fe] hydrogenase small subunit.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	88	1438	462	718	124	133	1	54	2	0

- Molecule 3 is dicarbonyl[bis(cyanide-kappaC)]-mu-(iminodimethanethiolato-1kappaS:2kappaS)-mu-(oxomethylidene)diiron(2+) (CCD ID: 402) (formula: C₇H₅Fe₂N₃O₃S₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	Fe	H	N	O	S	1	0
			24	7	2	7	3	3	2		

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			8	4	4		
4	A	1	Total	Fe	S	0	0
			8	4	4		
4	A	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 5 is 2-(2-ETHOXYETHOXY)ETHANOL (CCD ID: AE3) (formula: $C_6H_{14}O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	3	0
			22	6	13	3		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	S	2	0
			33	8	18	2	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	49	Total 49	O 49	0	0
7	B	9	Total 9	O 9	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Periplasmic [Fe] hydrogenase large subunit



- Molecule 2: Periplasmic [Fe] hydrogenase small subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.35Å 87.13Å 88.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.41 – 2.08 44.41 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.41-2.08) 99.4 (44.41-2.08)	Depositor EDS
R_{merge}	0.30	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.188 , 0.242 0.189 , 0.245	Depositor DCC
R_{free} test set	1182 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.507	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 32.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7646	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, 402, AE3, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.97	23/3130 (0.7%)	2.17	42/4234 (1.0%)
2	B	0.88	4/754 (0.5%)	1.55	8/1018 (0.8%)
All	All	1.81	27/3884 (0.7%)	2.06	50/5252 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
2	B	0	1
All	All	0	5

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320[A]	ASP	CB-CG	52.05	2.82	1.52
1	A	320[B]	ASP	CB-CG	52.05	2.82	1.52
1	A	367	LYS	C-O	-33.63	0.81	1.24
1	A	54	MET	CG-SD	32.25	2.61	1.80
1	A	53	GLU	CG-CD	-28.94	0.79	1.52
1	A	169	ASP	CB-CG	-23.77	0.92	1.52
1	A	367	LYS	CA-C	19.46	1.81	1.52
1	A	93	LYS	CG-CD	11.45	1.86	1.52
1	A	343	THR	CB-CG2	-10.79	1.17	1.52
1	A	346	LYS	CE-NZ	10.65	1.81	1.49
2	B	101	GLU	CD-OE1	10.62	1.45	1.25
1	A	166	LYS	CB-CG	10.30	1.83	1.52
1	A	220	LYS	CE-NZ	-9.41	1.21	1.49
1	A	281	LYS	CB-CG	-9.03	1.25	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	367	LYS	N-CA	-8.27	1.35	1.46
1	A	346	LYS	CD-CE	8.26	1.77	1.52
1	A	247	LYS	CE-NZ	-8.19	1.24	1.49
2	B	38	GLN	CD-NE2	-8.13	1.16	1.33
1	A	320[A]	ASP	CA-CB	-7.57	1.40	1.53
1	A	320[B]	ASP	CA-CB	-7.57	1.40	1.53
1	A	19	LYS	CD-CE	-7.04	1.31	1.52
1	A	317	LYS	CA-CB	-6.76	1.38	1.53
1	A	54	MET	SD-CE	-6.28	1.63	1.79
2	B	101	GLU	CD-OE2	6.02	1.36	1.25
2	B	73	LYS	CD-CE	-5.36	1.36	1.52
1	A	58	HIS	CG-CD2	-5.08	1.30	1.35
1	A	318	LYS	CB-CG	-5.01	1.37	1.52

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	135	GLN	OE1-CD-NE2	-74.21	48.39	122.60
1	A	320[A]	ASP	CB-CG-OD2	-42.01	21.77	118.40
1	A	320[B]	ASP	CB-CG-OD2	-42.01	21.77	118.40
1	A	53	GLU	CB-CG-CD	36.51	174.67	112.60
1	A	169	ASP	CA-CB-CG	23.97	136.57	112.60
2	B	73	LYS	CG-CD-CE	22.76	163.64	111.30
1	A	320[A]	ASP	CA-CB-CG	-17.04	95.56	112.60
1	A	320[B]	ASP	CA-CB-CG	-17.04	95.56	112.60
1	A	54	MET	CG-SD-CE	-14.45	69.10	100.90
1	A	343	THR	CA-CB-CG2	13.32	133.15	110.50
1	A	220	LYS	CD-CE-NZ	12.59	152.18	111.90
1	A	165	THR	CA-C-O	11.13	132.58	119.56
1	A	19	LYS	CG-CD-CE	10.99	136.58	111.30
1	A	367	LYS	O-C-N	10.98	135.83	122.21
1	A	320[A]	ASP	N-CA-CB	10.02	125.82	110.28
1	A	320[B]	ASP	N-CA-CB	10.02	125.82	110.28
1	A	165	THR	O-C-N	-9.99	110.35	122.34
2	B	73	LYS	CD-CE-NZ	-9.32	82.08	111.90
1	A	53	GLU	CG-CD-OE1	-8.81	98.14	118.40
2	B	101	GLU	OE1-CD-OE2	-8.44	102.66	122.90
1	A	169	ASP	CB-CG-OD1	-8.01	99.99	118.40
2	B	38	GLN	OE1-CD-NE2	7.84	130.44	122.60
1	A	367	LYS	CA-C-O	7.80	128.27	120.32
1	A	88	VAL	N-CA-CB	7.23	116.09	110.45
1	A	320[A]	ASP	CB-CG-OD1	-7.17	101.90	118.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	320[B]	ASP	CB-CG-OD1	-7.17	101.90	118.40
1	A	166	LYS	CA-CB-CG	-7.11	99.87	114.10
1	A	53	GLU	CG-CD-OE2	7.08	134.67	118.40
1	A	281	LYS	CA-CB-CG	6.94	127.98	114.10
1	A	346	LYS	CG-CD-CE	-6.86	95.53	111.30
1	A	34	LYS	CD-CE-NZ	-6.59	90.82	111.90
2	B	78	LYS	N-CA-C	6.34	113.83	108.13
1	A	222	ASP	CA-CB-CG	6.33	118.93	112.60
1	A	97	ASP	CA-CB-CG	6.33	118.92	112.60
1	A	163	ARG	N-CA-CB	6.32	119.52	110.16
1	A	247	LYS	CD-CE-NZ	6.24	131.88	111.90
1	A	136	LYS	CD-CE-NZ	6.22	131.80	111.90
1	A	19	LYS	CD-CE-NZ	6.21	131.76	111.90
2	B	94	ASP	CA-CB-CG	6.10	118.70	112.60
1	A	279	ASP	CB-CA-C	5.93	119.56	109.53
1	A	10	GLU	N-CA-CB	5.85	118.89	110.06
2	B	115	GLU	CB-CG-CD	5.76	122.39	112.60
2	B	78	LYS	CB-CA-C	-5.72	102.19	110.03
1	A	91	VAL	N-CA-CB	5.68	117.19	110.55
1	A	344	ASP	OD1-CG-OD2	-5.57	109.54	122.90
1	A	31	ASP	CA-CB-CG	5.53	118.13	112.60
1	A	343	THR	OG1-CB-CG2	-5.52	98.26	109.30
1	A	367	LYS	N-CA-CB	-5.30	103.39	111.66
1	A	24	LYS	CD-CE-NZ	5.15	128.38	111.90
1	A	135	GLN	CB-CA-C	5.09	119.50	110.85

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	135	GLN	Sidechain
1	A	169	ASP	Sidechain
1	A	233	PRO	Peptide
1	A	344	ASP	Sidechain
2	B	101	GLU	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3036	3011	2976	47	3
2	B	720	718	701	5	1
3	A	17	7	5	0	0
4	A	24	0	0	0	0
5	A	9	13	14	0	0
6	A	15	18	18	0	0
7	A	49	0	0	1	0
7	B	9	0	0	0	0
All	All	3879	3767	3714	49	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:MET:CE	1:A:54:MET:CG	2.54	0.86
1:A:351:HIS:HD2	1:A:380:GLY:H	1.30	0.79
1:A:351:HIS:CD2	1:A:380:GLY:H	2.07	0.71
1:A:367:LYS:C	1:A:367:LYS:CB	2.70	0.64
1:A:222:ASP:H	1:A:225:GLN:HE21	1.47	0.60
1:A:236:ALA:O	1:A:239:TYR:HB3	2.01	0.60
1:A:39:ASP:OD1	1:A:58:HIS:HE1	1.86	0.58
1:A:346:LYS:CD	1:A:346:LYS:NZ	2.67	0.57
1:A:54:MET:CG	1:A:54:MET:HE3	2.37	0.54
1:A:54:MET:CG	1:A:54:MET:HE2	2.37	0.54
1:A:54:MET:HE3	1:A:54:MET:CB	2.37	0.53
1:A:56:GLU:O	1:A:58:HIS:HD2	1.91	0.53
1:A:143:TRP:CD2	1:A:286:MET:HE3	2.45	0.52
1:A:141:HIS:HE1	1:A:221:TYR:OH	1.93	0.51
1:A:199:THR:OG1	2:B:89:HIS:HD2	1.94	0.50
1:A:71:GLN:O	1:A:75:HIS:HD2	1.94	0.50
1:A:367:LYS:C	1:A:367:LYS:N	2.69	0.50
1:A:14:HIS:HD2	1:A:85:GLN:OE1	1.95	0.49
1:A:36:ILE:HG12	1:A:78:GLU:HG3	1.95	0.48
1:A:222:ASP:O	1:A:225:GLN:HG2	2.13	0.48
1:A:323:ASP:OD1	2:B:60:ARG:NH2	2.41	0.48
1:A:156:GLU:OE1	7:A:601:HOH:O	2.20	0.47
1:A:295:ILE:O	1:A:301:GLY:HA3	2.16	0.46
1:A:219:MET:HE3	1:A:285:LEU:HD22	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:LYS:N	1:A:368:SER:N	2.64	0.45
1:A:231:ILE:HG21	1:A:263:LEU:HD22	1.99	0.45
1:A:88:VAL:N	1:A:89:PRO:CD	2.80	0.45
1:A:54:MET:HE3	1:A:54:MET:CA	2.48	0.44
1:A:92:GLU:OE1	1:A:266:MET:HE3	2.18	0.44
1:A:395:LEU:O	1:A:396:GLU:C	2.61	0.43
1:A:194:LEU:N	1:A:195[A]:PRO:CD	2.81	0.43
1:A:48:ALA:HA	2:B:116:PHE:CE2	2.52	0.43
1:A:194:LEU:N	1:A:195[B]:PRO:CD	2.81	0.43
1:A:163:ARG:NE	1:A:174:GLN:HE21	2.16	0.43
1:A:22:PRO:HB2	1:A:239:TYR:CD1	2.54	0.42
1:A:331:ASP:O	1:A:351:HIS:HE1	2.02	0.42
1:A:56:GLU:O	1:A:58:HIS:CD2	2.72	0.42
1:A:238:LYS:HE3	1:A:259:THR:HG23	2.02	0.42
1:A:143:TRP:CD2	1:A:286:MET:CE	3.03	0.42
1:A:143:TRP:CG	1:A:286:MET:HE3	2.55	0.42
1:A:54:MET:HE3	1:A:54:MET:HA	2.01	0.41
1:A:367:LYS:N	1:A:368:SER:H	2.17	0.41
1:A:247:LYS:HA	1:A:251:MET:O	2.21	0.41
1:A:58:HIS:CD2	1:A:58:HIS:N	2.88	0.41
2:B:59:VAL:HG13	2:B:64:ASP:HB2	2.02	0.41
1:A:88:VAL:O	1:A:92:GLU:HG2	2.22	0.40
2:B:114:SER:OG	2:B:115:GLU:OE2	2.29	0.40
1:A:141:HIS:CE1	1:A:221:TYR:OH	2.74	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:THR:HG1	2:B:36:VAL:H2[3_444]	1.13	0.47
1:A:14:HIS:HE1	1:A:53:GLU:OE1[4_555]	1.27	0.33
1:A:14:HIS:CE1	1:A:53:GLU:OE1[4_555]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/405 (98%)	393 (98%)	6 (2%)	0	100	100
2	B	88/88 (100%)	83 (94%)	5 (6%)	0	100	100
All	All	487/493 (99%)	476 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/330 (99%)	315 (97%)	11 (3%)	32	34
2	B	78/76 (103%)	76 (97%)	2 (3%)	40	45
All	All	404/406 (100%)	391 (97%)	13 (3%)	36	36

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	ILE
1	A	148	THR
1	A	169	ASP
1	A	220	LYS
1	A	281	LYS
1	A	284	SER
1	A	294	THR
1	A	320[A]	ASP
1	A	320[B]	ASP
1	A	362	ASP
1	A	396	GLU
2	B	82	HIS
2	B	83	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	HIS
1	A	58	HIS
1	A	75	HIS
1	A	85	GLN
1	A	135	GLN
1	A	141	HIS
1	A	174	GLN
1	A	225	GLN
1	A	351	HIS
1	A	358	GLN
2	B	89	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SF4	A	502	1	0,12,12	-	-	-		
4	SF4	A	504	1	0,12,12	-	-	-		
6	EPE	A	506	-	15,15,15	0.55	0	18,20,20	1.17	2 (11%)
4	SF4	A	503	1	0,12,12	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	AE3	A	505	-	8,8,8	0.23	0	7,7,7	0.71	0
3	402	A	501	1	13,19,19	6.08	8 (61%)	2,36,36	2.83	2 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	A	502	1	-	-	0/6/5/5
4	SF4	A	504	1	-	-	0/6/5/5
6	EPE	A	506	-	-	5/9/19/19	0/1/1/1
4	SF4	A	503	1	-	-	0/6/5/5
5	AE3	A	505	-	-	5/6/6/6	-
3	402	A	501	1	-	-	0/5/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	402	C2-S2	-10.95	1.65	1.85
3	A	501	402	S2-FE2	-10.90	2.10	2.26
3	A	501	402	C1-S1	-9.76	1.67	1.85
3	A	501	402	S1-FE1	-7.51	2.15	2.26
3	A	501	402	S1-FE2	-6.73	2.16	2.26
3	A	501	402	S2-FE1	-5.32	2.18	2.26
3	A	501	402	O5-C5	2.93	1.22	1.17
3	A	501	402	O7-C7	2.17	1.20	1.15

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	402	S2-C2-N1	-3.08	108.70	117.18
3	A	501	402	S1-C1-N1	-2.56	110.13	117.18
6	A	506	EPE	O1S-S-C10	-2.24	104.21	106.92
6	A	506	EPE	O3S-S-O2S	-2.01	106.37	111.27

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	506	EPE	S-C10-C9-N1

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	505	AE3	O3-C5-C6-O4
6	A	506	EPE	N4-C7-C8-O8
6	A	506	EPE	C8-C7-N4-C3
6	A	506	EPE	C9-C10-S-O2S
6	A	506	EPE	C8-C7-N4-C5
5	A	505	AE3	C1-C2-O2-C3
5	A	505	AE3	O2-C3-C4-O3
5	A	505	AE3	C3-C4-O3-C5
5	A	505	AE3	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	396/405 (97%)	0.01	4 (1%) 79 82	10, 26, 42, 71	29 (7%)
2	B	88/88 (100%)	0.07	2 (2%) 61 64	14, 28, 45, 50	7 (7%)
All	All	484/493 (98%)	0.02	6 (1%) 76 78	10, 26, 42, 71	36 (7%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	397	ALA	3.1
2	B	36	VAL	2.8
1	A	367	LYS	2.6
1	A	340	VAL	2.4
2	B	85	HIS	2.3
1	A	345	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	AE3	A	505	9/9	0.77	0.13	33,43,48,55	3
3	402	A	501	17/17	0.88	0.17	14,29,33,36	24
6	EPE	A	506	15/15	0.89	0.11	38,50,58,58	2
4	SF4	A	504	8/8	0.98	0.04	19,21,22,24	0
4	SF4	A	502	8/8	0.98	0.04	17,19,19,19	0
4	SF4	A	503	8/8	0.98	0.05	18,19,22,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around AE3 A 505:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 402 A 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.