



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 12:58 pm BST

PDB ID : 5RAV
Title : PanDDA analysis group deposition – Crystal Structure of JMJD1B in complex with FM001763a
Authors : Snee, M.; Nowak, R.; Johansson, C.; Burgess-Brown, N.A.; Arrowsmith, C.H.; Bountra, C.; Edwards, A.M.; Oppermann, U.
Deposited on : 2020-03-16
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Mol	Chain	Length	Quality of chain
1	A	372	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">4% 84% 7% 9%</p>
1	B	372	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">3% 84% 7% 8%</p>

ENTRY-COMPOSITION INFOmissingINFO



2 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific demethylase 3B



- Molecule 1: Lysine-specific demethylase 3B



3 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.75Å 93.69Å 93.57Å 90.00° 107.93° 90.00°	Depositor
Resolution (Å)	89.02 – 1.77 89.02 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.2 (89.02-1.77) 99.2 (89.02-1.77)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.179 , 0.209 0.193 , 0.219	Depositor DCC
R_{free} test set	4600 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtrriage
Anisotropy	0.054	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6292	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, S9G, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/2841	0.89	1/3854 (0.0%)
1	B	0.80	0/2872	0.83	0/3895
All	All	0.82	0/5713	0.86	1/7749 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1661	ARG	CG-CD-NE	-6.05	99.10	111.80

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2768	0	2650	25	0
1	B	2798	0	2685	24	0
2	A	18	0	0	0	0
3	A	4	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	379	0	0	9	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	322	0	0	7	0
All	All	6292	0	5335	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1517[B]:ARG:HH11	1:B:1517[B]:ARG:HG3	1.38	0.89
1:A:1582:HIS:CE1	5:A:1938:HOH:O	2.32	0.83
1:B:1518:ASP:HB3	5:B:2134:HOH:O	1.82	0.79
1:B:1517[B]:ARG:HH11	1:B:1517[B]:ARG:CG	1.95	0.78
1:A:1470:CYS:O	5:A:1902:HOH:O	2.03	0.75
1:A:1406[A]:ARG:NH1	5:A:1905:HOH:O	2.18	0.75
1:B:1517[B]:ARG:NH1	1:B:1517[B]:ARG:HG3	1.98	0.74
1:A:1407:GLU:HG2	5:A:2213:HOH:O	1.90	0.70
1:A:1639:GLU:O	5:A:1903:HOH:O	2.10	0.68
1:A:1582:HIS:O	1:A:1582:HIS:CG	2.47	0.67
1:A:1385:LEU:HD12	1:A:1392[B]:CYS:SG	2.35	0.65
1:B:1715:ARG:HG3	5:B:2119:HOH:O	1.97	0.65
1:A:1470:CYS:C	5:A:1902:HOH:O	2.36	0.64
1:A:1631:LYS:HE2	1:A:1666:TYR:OH	1.97	0.64
1:A:1406[A]:ARG:NH2	5:A:1907:HOH:O	2.29	0.64
1:B:1385:LEU:HD12	1:B:1392:CYS:SG	2.38	0.64
1:A:1583:ASP:O	1:A:1583:ASP:OD1	2.21	0.58
1:A:1502:PHE:CE2	1:A:1506:MET:HE2	2.38	0.58
1:A:1505:LEU:HD22	1:A:1700:VAL:HG11	1.88	0.55
1:B:1644:HIS:NE2	1:B:1649:ASP:OD2	2.39	0.55
1:B:1505:LEU:HD22	1:B:1700:VAL:HG11	1.88	0.54
1:A:1469:ILE:HD12	1:A:1579:GLU:HB2	1.90	0.52
1:B:1506:MET:CE	1:B:1538:PRO:HG3	2.40	0.52
1:B:1494:PHE:CZ	1:B:1498:MET:HE3	2.45	0.52
1:A:1407:GLU:CG	5:A:2213:HOH:O	2.54	0.52
1:A:1385:LEU:CD1	1:A:1392[B]:CYS:SG	2.98	0.51
1:A:1469:ILE:CD1	1:A:1579:GLU:HB2	2.41	0.50
1:B:1506:MET:O	1:B:1516:LYS:HE2	2.12	0.49
1:B:1715:ARG:HD3	5:B:2182:HOH:O	2.12	0.48
1:B:1606:HIS:HB2	1:B:1608:GLU:OE2	2.15	0.46
1:B:1644:HIS:CE1	1:B:1649:ASP:OD2	2.67	0.46
1:A:1582:HIS:HE1	5:A:1938:HOH:O	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1506:MET:HE3	1:B:1538:PRO:HD3	1.98	0.46
1:B:1516:LYS:HE3	5:B:1913:HOH:O	2.15	0.46
1:B:1520:ARG:NE	5:B:1917:HOH:O	2.50	0.45
1:B:1488:TRP:HA	1:B:1489:PRO:C	2.36	0.44
1:A:1583:ASP:C	1:A:1583:ASP:OD1	2.56	0.43
1:B:1453:ILE:HD11	5:B:2090:HOH:O	2.18	0.43
1:B:1604:HIS:HA	1:B:1607:LYS:NZ	2.33	0.43
1:A:1407:GLU:O	1:A:1410:LYS:HG2	2.18	0.43
1:A:1505:LEU:HD22	1:A:1700:VAL:CG1	2.48	0.43
1:A:1402:TRP:O	1:A:1406[A]:ARG:HG2	2.19	0.42
1:B:1602:ARG:HH11	1:B:1608:GLU:HG2	1.83	0.42
1:B:1517[B]:ARG:CB	1:B:1517[B]:ARG:HH11	2.31	0.42
1:A:1561:LEU:C	1:A:1561:LEU:HD12	2.40	0.41
1:A:1582:HIS:CD2	1:A:1585:GLU:HG3	2.56	0.41
1:B:1599:THR:O	1:B:1603:ILE:HG23	2.21	0.41
1:A:1547:ILE:HA	1:A:1582:HIS:NE2	2.35	0.41
1:B:1383:SER:HB2	5:B:2143:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2212:HOH:O	5:A:2242:HOH:O[2_648]	1.82	0.38

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/372 (92%)	336 (98%)	6 (2%)	1 (0%)	41	25
1	B	342/372 (92%)	328 (96%)	14 (4%)	0	100	100
All	All	685/744 (92%)	664 (97%)	20 (3%)	1 (0%)	51	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1581	ALA

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/331 (90%)	292 (98%)	7 (2%)	50	34
1	B	302/331 (91%)	294 (97%)	8 (3%)	46	29
All	All	601/662 (91%)	586 (98%)	15 (2%)	49	31

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1406[A]	ARG
1	A	1406[B]	ARG
1	A	1584	GLU
1	A	1603	ILE
1	A	1644	HIS
1	A	1707	PRO
1	A	1717	THR
1	B	1494	PHE
1	B	1577	ILE
1	B	1582	HIS
1	B	1588	LYS
1	B	1603	ILE
1	B	1604	HIS
1	B	1605	ASP
1	B	1607	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1640	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1650	GLN
1	B	1656	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 could not be matched to an existing wwPDB Chemical Component Dictionary definition at this stage and 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/372 (91%)	-0.18	16 (4%) 31 29	16, 27, 59, 108	12 (3%)
1	B	344/372 (92%)	-0.27	11 (3%) 47 46	18, 30, 61, 126	2 (0%)
All	All	684/744 (91%)	-0.23	27 (3%) 39 38	16, 28, 60, 126	14 (2%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1581	ALA	14.7
1	A	1392[A]	CYS	9.2
1	B	1605	ASP	8.8
1	A	1577	ILE	8.1
1	A	1580	GLY	8.0
1	A	1379[A]	MET	7.7
1	B	1581	ALA	6.9
1	A	1578	GLY	6.6
1	B	1577	ILE	5.9
1	A	1564[A]	SER	5.5
1	A	1582	HIS	5.5
1	A	1406[A]	ARG	5.4
1	A	1588[A]	LYS	5.0
1	A	1579	GLU	4.9
1	B	1517[A]	ARG	4.7
1	B	1604	HIS	4.6
1	B	1606	HIS	4.5
1	B	1587	LEU	3.9
1	B	1664[A]	GLU	3.8
1	A	1378	SER	3.5
1	B	1603	ILE	3.4
1	B	1582	HIS	3.4
1	A	1583	ASP	2.9
1	A	1606	HIS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1642	PRO	2.8
1	B	1601	GLU	2.6
1	A	1639	GLU	2.6

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

5.5 Other polymers [i](#)

There are no such residues in this entry.