



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2023 – 07:08 PM EDT

PDB ID : 7RAH  
Title : Adenylate cyclase toxin RTX domain fragment bound to M1H5 Fab and M2B10 Fab  
Authors : Goldsmith, J.A.; McLellan, J.S.  
Deposited on : 2021-07-01  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	87% (green), 13% (yellow)
2	B	233	89% (green), 6% (yellow), 5% (grey)
3	C	214	7% (red), 74% (green), 12% (yellow), 7% (grey)
4	D	235	10% (red), 69% (green), 16% (yellow), 5% (grey)
5	E	458	71% (green), 7% (yellow), 21% (grey)

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9183 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M1H5 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	1671	1035	282	347	7	0	0	0

- Molecule 2 is a protein called M1H5 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	223	1680	1060	278	333	9	0	0	0

- Molecule 3 is a protein called M2B10 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	185	1395	870	232	286	7	0	0	0

- Molecule 4 is a protein called M2B10 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	199	1499	948	252	294	5	0	0	0

- Molecule 5 is a protein called Bifunctional adenylate cyclase toxin/hemolysin CyaA,Bifunctional adenylate cyclase toxin/hemolysin CyaA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	360	2726	1664	484	573	5	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	1011	GLY	-	expression tag	UNP A0A380ZZA1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1012	PRO	-	expression tag	UNP A0A380ZZA1
E	1013	GLY	-	expression tag	UNP A0A380ZZA1
E	1014	SER	-	expression tag	UNP A0A380ZZA1

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O P 5 4 1	0	0

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	E	19	Total Ca 19 19	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	48	Total O 48 48	0	0
8	B	44	Total O 44 44	0	0
8	C	18	Total O 18 18	0	0

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
8	D	10	Total	O	0	0
			10	10		
8	E	68	Total	O	0	0
			68	68		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: M1H5 Fab Light Chain

Chain A: 87% 13%




- Molecule 2: M1H5 Fab Heavy Chain

Chain B: 89% 6%




- Molecule 3: M2B10 Fab Light Chain

Chain C: 7% 74% 12% 14%




- Molecule 4: M2B10 Fab Heavy Chain

Chain D: 10% 69% 16% 15%



- Molecule 5: Bifunctional adenylate cyclase toxin/hemolysin CyaA, Bifunctional adenylate cyclase toxin/hemolysin CyaA

Chain E:  71% 7% 21%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.56Å 117.02Å 254.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.81 – 2.60 68.72 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (55.81-2.60) 99.7 (68.72-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 2.62Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122+SVN	Depositor
R, $R_{free}$	0.195 , 0.218 0.192 , 0.214	Depositor DCC
$R_{free}$ test set	3076 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.8	Xtrriage
Anisotropy	0.437	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9183	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/1705	0.50	0/2314
2	B	0.26	0/1721	0.51	0/2339
3	C	0.25	0/1427	0.49	0/1940
4	D	0.26	0/1534	0.51	0/2087
5	E	0.24	0/2772	0.53	0/3756
All	All	0.25	0/9159	0.51	0/12436

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1671	0	1597	19	0
2	B	1680	0	1644	8	0
3	C	1395	0	1339	11	0
4	D	1499	0	1458	27	0
5	E	2726	0	2493	20	0
6	A	5	0	0	0	0
7	E	19	0	0	0	0
8	A	48	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	44	0	0	3	0
8	C	18	0	0	1	0
8	D	10	0	0	2	0
8	E	68	0	0	6	1
All	All	9183	0	8531	82	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:1288:ARG:NH1	8:E:1601:HOH:O	2.00	0.95
1:A:132:SER:O	8:A:401:HOH:O	1.89	0.88
4:D:121:VAL:HG21	4:D:207:VAL:HG11	1.56	0.84
5:E:1360:HIS:ND1	8:E:1603:HOH:O	2.10	0.82
5:E:1324:GLU:OE1	8:E:1602:HOH:O	2.01	0.78
1:A:156:ARG:NH1	8:A:403:HOH:O	2.18	0.73
3:C:105:GLU:OE2	3:C:140:TYR:OH	2.08	0.70
1:A:38:GLN:HG3	1:A:44:VAL:HG22	1.72	0.69
2:B:13:LYS:HG3	8:B:311:HOH:O	1.95	0.66
1:A:166:ASP:OD1	8:A:402:HOH:O	2.13	0.66
4:D:2:VAL:N	8:D:302:HOH:O	2.29	0.65
1:A:191:ASN:HD21	1:A:213:ASN:H	1.47	0.62
4:D:19:LYS:NZ	4:D:81:GLN:HB2	2.14	0.62
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.82	0.60
3:C:171:SER:OG	8:C:301:HOH:O	2.16	0.60
1:A:24:ARG:HG3	1:A:24:ARG:HH11	1.68	0.58
1:A:191:ASN:ND2	1:A:213:ASN:H	2.01	0.58
4:D:200:HIS:CD2	4:D:202:PRO:HD2	2.37	0.58
5:E:1324:GLU:HG3	5:E:1353:LEU:HB2	1.85	0.57
1:A:24:ARG:HG3	1:A:24:ARG:NH1	2.20	0.57
1:A:29:ILE:HA	1:A:92:ASN:HD22	1.70	0.57
1:A:115:THR:HG23	1:A:138:ASN:HB3	1.87	0.57
3:C:136:LEU:HD21	3:C:196:VAL:HG21	1.88	0.55
4:D:209:LYS:HB2	4:D:209:LYS:NZ	2.21	0.55
4:D:45:LEU:N	8:D:301:HOH:O	2.05	0.54
5:E:1153:GLU:HG2	5:E:1171:ILE:HB	1.89	0.54
4:D:148:GLU:HG2	4:D:149:PRO:HA	1.89	0.54
5:E:1140:ARG:N	8:E:1604:HOH:O	2.15	0.53
5:E:1178:LEU:HD12	5:E:1197:VAL:HG13	1.90	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:ARG:NH1	1:A:182:LEU:HD21	2.24	0.52
4:D:40:ARG:HG2	4:D:41:PRO:HD2	1.92	0.52
3:C:117:ILE:HD11	3:C:207:LYS:HB3	1.93	0.50
2:B:119:PRO:HB3	2:B:145:TYR:HB3	1.94	0.50
3:C:110:VAL:HG12	3:C:141:PRO:HD3	1.93	0.50
4:D:19:LYS:NZ	4:D:81:GLN:OE1	2.45	0.50
1:A:185:ASP:O	1:A:189:ARG:HG3	2.12	0.49
4:D:19:LYS:CE	4:D:81:GLN:HB2	2.42	0.49
3:C:135:LEU:HD22	4:D:181:VAL:HG11	1.95	0.49
4:D:159:LEU:HD11	4:D:182:VAL:HG11	1.96	0.48
5:E:1057:THR:HG23	5:E:1077:GLN:HB2	1.96	0.48
5:E:1211:ASP:OD1	5:E:1213:SER:OG	2.29	0.48
1:A:191:ASN:OD1	1:A:211:ASN:ND2	2.38	0.47
4:D:68:THR:HB	4:D:81:GLN:HB3	1.95	0.47
2:B:155:ASN:O	8:B:301:HOH:O	2.20	0.47
2:B:165:THR:OG1	8:B:302:HOH:O	2.21	0.46
1:A:124:GLU:N	1:A:124:GLU:OE1	2.47	0.46
4:D:38:LYS:HB2	4:D:48:ILE:HD11	1.97	0.46
3:C:134:CYS:HB2	3:C:148:TRP:CH2	2.51	0.46
4:D:201:LYS:HB3	4:D:202:PRO:HD3	1.98	0.46
3:C:135:LEU:CD2	4:D:181:VAL:HG11	2.47	0.45
5:E:1404:VAL:HG11	5:E:1407:TRP:HD1	1.81	0.45
5:E:1374:ARG:HG2	5:E:1419:HIS:HB2	1.98	0.45
4:D:156:SER:H	4:D:197:ASN:HD21	1.65	0.45
1:A:18:ARG:HG3	1:A:76:SER:HA	1.99	0.45
4:D:96:LEU:HD11	4:D:100(B):GLN:HB3	1.99	0.45
4:D:119:PRO:HD3	4:D:200:HIS:ND1	2.32	0.45
4:D:170:LEU:HB2	4:D:176:TYR:HE1	1.82	0.45
5:E:1135:ASN:CG	5:E:1153:GLU:HB2	2.38	0.44
1:A:24:ARG:NH1	1:A:70:ASP:OD1	2.51	0.43
4:D:119:PRO:HA	4:D:144:ASP:O	2.18	0.43
5:E:1392:GLU:OE2	5:E:1394:ARG:NE	2.48	0.43
5:E:1255:ARG:NE	8:E:1605:HOH:O	2.17	0.43
3:C:133:VAL:HG22	3:C:178:THR:HG23	2.00	0.43
5:E:1073:VAL:HA	5:E:1126:SER:HB2	1.99	0.43
3:C:122:ASP:N	3:C:122:ASP:OD1	2.52	0.43
1:A:14:SER:O	1:A:17:ASP:HB2	2.19	0.43
4:D:171:GLN:H	4:D:171:GLN:HG2	1.70	0.42
5:E:1110:ASP:OD1	5:E:1166:ARG:NH1	2.51	0.42
3:C:108:ARG:HG2	3:C:109:THR:N	2.34	0.42
5:E:1212:TYR:O	5:E:1263:THR:HG22	2.20	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:199:ASN:HD22	2:B:206:LYS:HE2	1.85	0.41
4:D:40:ARG:NH2	4:D:84:SER:O	2.53	0.41
2:B:168:ALA:HA	2:B:178:LEU:HB3	2.02	0.41
4:D:67:ALA:HA	4:D:81:GLN:O	2.19	0.41
4:D:119:PRO:HD2	4:D:205:THR:OG1	2.21	0.41
2:B:12:VAL:HG21	2:B:82(C):LEU:HD13	2.02	0.41
4:D:144:ASP:OD1	4:D:171:GLN:NE2	2.54	0.41
5:E:1380:ASP:OD1	5:E:1380:ASP:N	2.49	0.41
1:A:121:PRO:HD3	1:A:133:VAL:HG22	2.03	0.41
2:B:100:GLY:HA2	5:E:1270:ILE:HD11	2.04	0.40
4:D:19:LYS:HZ1	4:D:81:GLN:HB2	1.85	0.40
5:E:1408:TYR:O	8:E:1606:HOH:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:E:1619:HOH:O	8:E:1636:HOH:O[3_555]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/215 (99%)	209 (98%)	4 (2%)	0	100	100
2	B	221/233 (95%)	218 (99%)	3 (1%)	0	100	100
3	C	179/214 (84%)	174 (97%)	5 (3%)	0	100	100
4	D	193/235 (82%)	188 (97%)	5 (3%)	0	100	100
5	E	356/458 (78%)	339 (95%)	17 (5%)	0	100	100
All	All	1162/1355 (86%)	1128 (97%)	34 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	191 (99%)	2 (1%)	76	90
2	B	191/200 (96%)	188 (98%)	3 (2%)	62	82
3	C	161/186 (87%)	152 (94%)	9 (6%)	21	42
4	D	169/200 (84%)	162 (96%)	7 (4%)	30	56
5	E	277/341 (81%)	271 (98%)	6 (2%)	52	76
All	All	991/1120 (88%)	964 (97%)	27 (3%)	44	71

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	170	LYS
2	B	92	CYS
2	B	98	ARG
2	B	197	ASN
3	C	23	CYS
3	C	27(A)	SER
3	C	30	SER
3	C	33	LEU
3	C	45	LYS
3	C	91	TYR
3	C	142	ARG
3	C	146	VAL
3	C	203	SER
4	D	56	ASN
4	D	66	LYS
4	D	84	SER
4	D	114	ARG
4	D	138	LEU
4	D	145	TYR
4	D	188	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
5	E	1131	LYS
5	E	1149	ASP
5	E	1198	SER
5	E	1220	ARG
5	E	1288	ARG
5	E	1412	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	199	ASN
3	C	53	ASN
5	E	1329	ASN
5	E	1381	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 19 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	PO4	A	301	-	4,4,4	0.91	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/215 (100%)	0.25	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	38, 53, 74, 86	0
2	B	223/233 (95%)	0.24	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	37, 49, 64, 93	0
3	C	185/214 (86%)	0.50	14 (7%) <span style="border: 1px solid red; padding: 2px;">13</span> <span style="border: 1px solid red; padding: 2px;">10</span>	46, 65, 142, 164	0
4	D	199/235 (84%)	0.71	24 (12%) <span style="border: 1px solid red; padding: 2px;">4</span> <span style="border: 1px solid red; padding: 2px;">2</span>	49, 100, 142, 238	0
5	E	360/458 (78%)	0.27	1 (0%) <span style="border: 1px solid blue; padding: 2px;">94</span> <span style="border: 1px solid blue; padding: 2px;">93</span>	40, 52, 71, 85	0
All	All	1182/1355 (87%)	0.37	39 (3%) <span style="border: 1px solid red; padding: 2px;">46</span> <span style="border: 1px solid red; padding: 2px;">39</span>	37, 56, 125, 238	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	1338	ARG	4.1
4	D	138	LEU	4.1
4	D	207	VAL	3.9
4	D	122	PHE	3.9
4	D	11	LEU	3.8
3	C	133	VAL	3.8
4	D	182	VAL	3.8
4	D	137	ALA	3.7
4	D	178	LEU	3.7
3	C	132	VAL	3.7
3	C	194	CYS	3.7
4	D	170	LEU	3.6
3	C	134	CYS	3.4
4	D	143	LYS	3.3
4	D	120	SER	3.2
4	D	121	VAL	3.1
4	D	159	LEU	3.1
3	C	122	ASP	3.0
3	C	118	PHE	3.0
3	C	160	GLN	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	C	131	SER	2.8
3	C	146	VAL	2.8
3	C	116	PHE	2.7
3	C	148	TRP	2.7
4	D	177	SER	2.6
4	D	150	VAL	2.6
4	D	183	THR	2.5
4	D	169	VAL	2.5
4	D	202	PRO	2.4
4	D	198	VAL	2.4
3	C	2	ILE	2.4
4	D	82	LEU	2.3
4	D	141	LEU	2.3
3	C	176	SER	2.2
4	D	136	ALA	2.1
4	D	153	SER	2.1
4	D	206	LYS	2.1
4	D	24	ALA	2.1
3	C	178	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	CA	E	1517	1/1	0.73	0.15	121,121,121,121	0
7	CA	E	1519	1/1	0.85	0.12	79,79,79,79	0
7	CA	E	1501	1/1	0.95	0.25	76,76,76,76	0
7	CA	E	1518	1/1	0.96	0.16	48,48,48,48	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	PO4	A	301	5/5	0.96	0.20	74,81,94,106	0
7	CA	E	1503	1/1	0.97	0.20	53,53,53,53	0
7	CA	E	1502	1/1	0.97	0.22	76,76,76,76	0
7	CA	E	1516	1/1	0.98	0.18	44,44,44,44	0
7	CA	E	1510	1/1	0.98	0.24	41,41,41,41	0
7	CA	E	1509	1/1	0.99	0.18	48,48,48,48	0
7	CA	E	1504	1/1	0.99	0.20	39,39,39,39	0
7	CA	E	1511	1/1	0.99	0.24	31,31,31,31	0
7	CA	E	1513	1/1	0.99	0.19	33,33,33,33	0
7	CA	E	1514	1/1	0.99	0.20	37,37,37,37	0
7	CA	E	1515	1/1	0.99	0.16	47,47,47,47	0
7	CA	E	1505	1/1	0.99	0.22	37,37,37,37	0
7	CA	E	1506	1/1	0.99	0.25	36,36,36,36	0
7	CA	E	1507	1/1	0.99	0.27	35,35,35,35	0
7	CA	E	1508	1/1	0.99	0.19	44,44,44,44	0
7	CA	E	1512	1/1	1.00	0.20	35,35,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.