



Full wwPDB EM Validation Report ⓘ

Apr 4, 2024 – 05:29 pm BST

PDB ID : 8RBZ
EMDB ID : EMD-19040
Title : Structure of Integrator-PP2A-SOSS-CTD post-termination complex
Authors : Fianu, I.; Ochmann, M.; Walshe, J.L.; Cramer, P.
Deposited on : 2023-12-05
Resolution : 3.70 Å (reported)
Based on initial model : ?

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.
















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	9	100%
2	B	213	48% 50%
3	C	106	37% 63%
4	Y	13	100%
5	a	2192	49% 51%
6	b	1204	87% 13%
7	c	1042	44% 56%
8	d	963	85% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	e	1021	 83% 16%
10	f	889	 61% 39%
11	g	964	 91% 8%
12	h	995	 88% 11%
13	i	658	 95% 5%
14	j	710	 15% 94% 6%
15	k	602	 90% 8%
16	n	518	 24% 39% 61%
17	o	451	 81% 19%
18	p	591	 97% ..
19	q	311	 93% 7%
20	r	27	 100%
21	m	706	 5% 95%

2 Entry composition i

There are 23 unique types of molecules in this entry. The entry contains 74102 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UNK-UNK-UNK-UNK-UNK-UNK-UNK-UNK-UNK.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
1	1	9	45	27	9	9	0	0

- Molecule 2 is a protein called SOSS complex subunit B1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	107	806	516	135	151	4	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	SER	-	expression tag	UNP Q9BQ15
B	0	ASN	-	expression tag	UNP Q9BQ15

- Molecule 3 is a protein called SOSS complex subunit C.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
3	C	39	296	190	56	50	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	SER	-	expression tag	UNP Q9NRY2
C	0	ASN	-	expression tag	UNP Q9NRY2

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	Y	13	95	60	13	22	0	0

- Molecule 5 is a protein called Integrator complex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	a	1085	8525	5425	1502	1547	51	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	-1	SER	-	expression tag	UNP Q8N201
a	0	ASN	-	expression tag	UNP Q8N201

- Molecule 6 is a protein called Integrator complex subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	b	1051	7988	5120	1352	1454	62	0	0

- Molecule 7 is a protein called Integrator complex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	c	463	3717	2380	646	662	29	0	0

- Molecule 8 is a protein called Integrator complex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	d	818	6395	4081	1091	1189	34	0	0

- Molecule 9 is a protein called Integrator complex subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	e	855	6253	3974	1139	1114	26	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
e	-1	SER	-	expression tag	UNP Q6P9B9
e	0	ASN	-	expression tag	UNP Q6P9B9

- Molecule 10 is a protein called Integrator complex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	f	544	4137	2653	701	758	25	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
f	-1	SER	-	expression tag	UNP Q9UL03
f	0	ASN	-	expression tag	UNP Q9UL03

- Molecule 11 is a protein called Integrator complex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	g	886	6725	4261	1169	1255	40	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
g	-1	SER	-	expression tag	UNP Q9NVH2
g	0	ASN	-	expression tag	UNP Q9NVH2

- Molecule 12 is a protein called Integrator complex subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	h	887	6748	4333	1156	1221	38	0	0

- Molecule 13 is a protein called Integrator complex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	i	626	4867	3134	792	908	33	0	0

- Molecule 14 is a protein called Integrator complex subunit 10.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
14	j	668	3313	1976	668	669	0	0

- Molecule 15 is a protein called Integrator complex subunit 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
15	k	551	4079	2613	706	734	26	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
k	-1	SER	-	expression tag	UNP Q5TA45
k	0	ASN	-	expression tag	UNP Q5TA45
k	203	GLN	GLU	engineered mutation	UNP Q5TA45
k	487	GLU	ASP	conflict	UNP Q5TA45

- Molecule 16 is a protein called Integrator complex subunit 14.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
16	n	204	1007	599	204	204	0	0

- Molecule 17 is a protein called Integrator complex subunit 15.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
17	o	367	1824	1090	367	367	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
o	-1	SER	-	expression tag	UNP Q96N11
o	0	ASN	-	expression tag	UNP Q96N11

- Molecule 18 is a protein called Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A alpha isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	p	580	4432	2822	753	831	26	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
p	-1	SER	-	expression tag	UNP P30153
p	0	ASN	-	expression tag	UNP P30153

- Molecule 19 is a protein called Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
19	q	290	2322	1467	403	437	15	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
q	-1	SER	-	expression tag	UNP P67775
q	0	ASN	-	expression tag	UNP P67775
q	88	ASN	ASP	engineered mutation	UNP P67775

- Molecule 20 is a protein called DSS1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
20	r	27	237	146	37	53	1	0	0

- Molecule 21 is a protein called Integrator complex subunit 13.


Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
21	m	35	287	175	61	51	0	0

- Molecule 22 is ZINC ION (three-letter code: ZN) (formula: Zn).



Mol	Chain	Residues	Atoms		AltConf
22	k	2	Total	Zn	0
			2	2	

- Molecule 23 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		AltConf
23	q	2	Total	Mn	0
			2	2	



• Molecule 6: Integrator complex subunit 2





• Molecule 7: Integrator complex subunit 3




• Molecule 8: Integrator complex subunit 4


• Molecule 11: Integrator complex subunit 7

Chain g:  91% 8%





• Molecule 12: Integrator complex subunit 8

Chain h:  88% 11%




• Molecule 13: Integrator complex subunit 9

Chain i:  95% 5%




• Molecule 14: Integrator complex subunit 10

Chain j:  15% 94% 6%



• Molecule 15: Integrator complex subunit 11

Chain k:  90% 8%

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	236382	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	39.93	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.195	Depositor
Minimum map value	-0.083	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.00308	Depositor
Map size (\AA)	503.99997, 503.99997, 503.99997	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.05, 1.05, 1.05	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
2	B	0.29	0/820	0.61	0/1109
3	C	0.26	0/304	0.49	0/413
4	Y	0.27	0/100	0.46	0/139
5	a	0.28	0/8682	0.58	2/11763 (0.0%)
6	b	0.31	0/8128	0.59	1/11072 (0.0%)
7	c	0.28	0/3791	0.57	0/5134
8	d	0.29	0/6519	0.57	2/8853 (0.0%)
9	e	0.29	0/6390	0.54	0/8706
10	f	0.28	0/4240	0.58	0/5784
11	g	0.29	0/6835	0.55	1/9271 (0.0%)
12	h	0.28	0/6867	0.52	0/9330
13	i	0.29	0/4986	0.60	1/6798 (0.0%)
14	j	0.24	0/3309	0.36	0/4608
15	k	0.28	0/4168	0.57	0/5665
16	n	0.24	0/1006	0.41	0/1399
17	o	0.24	0/1821	0.36	0/2538
18	p	0.31	0/4506	0.61	1/6130 (0.0%)
19	q	0.28	0/2378	0.59	0/3228
20	r	0.27	0/241	0.48	0/327
21	m	0.25	0/292	0.62	0/393
All	All	0.29	0/75383	0.56	8/102660 (0.0%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
18	p	421	LEU	CA-CB-CG	5.94	128.95	115.30
6	b	54	LEU	CA-CB-CG	5.84	128.73	115.30
5	a	1835	LEU	CA-CB-CG	5.49	127.94	115.30
11	g	235	LEU	CA-CB-CG	5.34	127.59	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	i	309	LEU	CA-CB-CG	5.29	127.46	115.30
5	a	2088	LEU	CA-CB-CG	5.19	127.23	115.30
8	d	493	LEU	CA-CB-CG	5.02	126.84	115.30
8	d	519	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	45	0	12	0	0
2	B	806	0	806	3	0
3	C	296	0	286	0	0
4	Y	95	0	82	0	0
5	a	8525	0	8673	0	0
6	b	7988	0	8032	0	0
7	c	3717	0	3806	0	0
8	d	6395	0	6420	0	0
9	e	6253	0	6063	0	0
10	f	4137	0	3989	0	0
11	g	6725	0	6694	0	0
12	h	6748	0	6613	0	0
13	i	4867	0	4840	0	0
14	j	3313	0	1464	0	0
15	k	4079	0	3873	0	0
16	n	1007	0	439	0	0
17	o	1824	0	802	0	0
18	p	4432	0	4487	0	0
19	q	2322	0	2215	0	0
20	r	237	0	201	0	0
21	m	287	0	275	0	0
22	k	2	0	0	0	0
23	q	2	0	0	0	0
All	All	74102	0	70072	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:91:ASP:N	2:B:91:ASP:OD1	2.49	0.46
2:B:67:ILE:HG22	2:B:95:ILE:HD13	1.99	0.44
2:B:97:GLU:OE1	2:B:98:PHE:N	2.52	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	105/213 (49%)	100 (95%)	5 (5%)	0	100	100
3	C	37/106 (35%)	37 (100%)	0	0	100	100
4	Y	11/13 (85%)	11 (100%)	0	0	100	100
5	a	1061/2192 (48%)	1031 (97%)	29 (3%)	1 (0%)	51	83
6	b	1033/1204 (86%)	981 (95%)	52 (5%)	0	100	100
7	c	461/1042 (44%)	447 (97%)	14 (3%)	0	100	100
8	d	806/963 (84%)	787 (98%)	19 (2%)	0	100	100
9	e	833/1021 (82%)	800 (96%)	33 (4%)	0	100	100
10	f	530/889 (60%)	512 (97%)	18 (3%)	0	100	100
11	g	876/964 (91%)	848 (97%)	27 (3%)	1 (0%)	51	83
12	h	865/995 (87%)	837 (97%)	28 (3%)	0	100	100
13	i	618/658 (94%)	595 (96%)	23 (4%)	0	100	100
14	j	660/710 (93%)	643 (97%)	17 (3%)	0	100	100
15	k	539/602 (90%)	513 (95%)	25 (5%)	1 (0%)	47	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
16	n	202/518 (39%)	197 (98%)	5 (2%)	0	100	100
17	o	361/451 (80%)	343 (95%)	17 (5%)	1 (0%)	41	74
18	p	578/591 (98%)	561 (97%)	17 (3%)	0	100	100
19	q	288/311 (93%)	275 (96%)	13 (4%)	0	100	100
20	r	25/27 (93%)	24 (96%)	1 (4%)	0	100	100
21	m	33/706 (5%)	30 (91%)	3 (9%)	0	100	100
All	All	9922/14176 (70%)	9572 (96%)	346 (4%)	4 (0%)	100	100

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	a	1630	VAL
11	g	885	ASN
15	k	305	ILE
17	o	318	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	87/181 (48%)	86 (99%)	1 (1%)	73	85
3	C	28/87 (32%)	28 (100%)	0	100	100
4	Y	13/13 (100%)	13 (100%)	0	100	100
5	a	946/1909 (50%)	940 (99%)	6 (1%)	86	93
6	b	869/1072 (81%)	861 (99%)	8 (1%)	78	88
7	c	414/927 (45%)	409 (99%)	5 (1%)	71	84
8	d	702/845 (83%)	700 (100%)	2 (0%)	92	96
9	e	616/814 (76%)	612 (99%)	4 (1%)	86	93
10	f	438/798 (55%)	436 (100%)	2 (0%)	88	94
11	g	723/842 (86%)	720 (100%)	3 (0%)	91	95

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
12	h	700/896 (78%)	693 (99%)	7 (1%)	76	86
13	i	554/600 (92%)	551 (100%)	3 (0%)	88	94
15	k	403/522 (77%)	397 (98%)	6 (2%)	65	81
18	p	485/514 (94%)	479 (99%)	6 (1%)	71	84
19	q	251/276 (91%)	249 (99%)	2 (1%)	81	89
20	r	27/27 (100%)	27 (100%)	0	100	100
21	m	30/639 (5%)	30 (100%)	0	100	100
All	All	7286/10962 (66%)	7231 (99%)	55 (1%)	82	89

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	98	PHE
5	a	1088	ASP
5	a	1630	VAL
5	a	1746	ARG
5	a	1842	PHE
5	a	1983	ASP
5	a	2171	MET
6	b	130	PHE
6	b	435	MET
6	b	461	LYS
6	b	505	MET
6	b	537	ARG
6	b	574	LYS
6	b	845	GLN
6	b	1093	GLN
7	c	131	MET
7	c	244	PHE
7	c	397	TYR
7	c	402	PHE
7	c	423	MET
8	d	156	CYS
8	d	412	CYS
9	e	154	GLN
9	e	181	MET
9	e	225	TRP
9	e	407	LEU
10	f	412	PHE
10	f	434	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
11	g	672	LYS
11	g	808	LEU
11	g	826	GLN
12	h	250	CYS
12	h	271	TYR
12	h	295	CYS
12	h	407	PHE
12	h	430	LYS
12	h	447	CYS
12	h	883	ARG
13	i	281	PHE
13	i	394	PHE
13	i	462	LYS
15	k	178	ASP
15	k	204	SER
15	k	261	LEU
15	k	267	PHE
15	k	302	PHE
15	k	576	TRP
18	p	125	GLU
18	p	128	PHE
18	p	245	MET
18	p	434	LEU
18	p	490	SER
18	p	538	PHE
19	q	49	ARG
19	q	86	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
9	e	690	GLN
11	g	145	GLN
12	h	988	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19040. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 240




Y Index: 240




Z Index: 240


6.2.2 Raw map



X Index: 240



Y Index: 240




Z Index: 240

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 230




Y Index: 280




Z Index: 282


6.3.2 Raw map



X Index: 230



Y Index: 240




Z Index: 281


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00308. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 2799 nm^3 ; this corresponds to an approximate mass of 2528 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.270 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.270\AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.70	-	-
Author-provided FSC curve	3.80	5.48	3.86
Unmasked-calculated*	4.37	7.53	4.50

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.37 differs from the reported value 3.7 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-19040 and PDB model 8RBZ. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.00308 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00308).













































9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.00308) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9770	 0.2930
1	 1.0000	 0.3650
B	 1.0000	 0.0960
C	 0.9630	 0.0550
Y	 0.9890	 0.5060
a	 0.9830	 0.2600
b	 0.9930	 0.3820
c	 0.9850	 0.1360
d	 0.9970	 0.3340
e	 0.9940	 0.3530
f	 0.9950	 0.3430
g	 0.9930	 0.4080
h	 0.9950	 0.3840
i	 0.9960	 0.2720
j	 0.8400	 0.0330
k	 0.9950	 0.1930
m	 0.9890	 0.0910
n	 0.3880	 0.0150
o	 0.9750	 0.0730
p	 0.9980	 0.2470
q	 0.9940	 0.4750
r	 0.9910	 0.3220

