



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2023 – 12:38 AM EDT

PDB ID : 7RCJ  
Title : Crystal structure of ZnuA from *Citrobacter koseri*  
Authors : Yukl, E.T.; Yekwa, E.L.  
Deposited on : 2021-07-07  
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1665 (3.20-3.12)
Clashscore	141614	1804 (3.20-3.12)
Ramachandran outliers	138981	1770 (3.20-3.12)
Sidechain outliers	138945	1769 (3.20-3.12)
RSRZ outliers	127900	1616 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	 62% 18% 17%
1	B	316	 60% 20% 16%
1	C	316	 59% 21% 17%
1	D	316	 61% 19% 16%
1	E	316	 64% 15% 16%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	316	 60% 19% 17%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called High-affinity zinc uptake system protein ZnuA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	2020	1289	346	377	8	0	0	0
1	B	264	2028	1293	347	380	8	0	0	0
1	C	263	2020	1289	346	377	8	0	0	0
1	D	264	2028	1293	347	380	8	0	0	0
1	E	264	2028	1293	347	380	8	0	0	0
1	F	263	2020	1289	346	377	8	0	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0
2	D	1	Total 1	Zn 1	0	0
2	E	1	Total 1	Zn 1	0	0
2	F	1	Total 1	Zn 1	0	0

- Molecule 3 is 6-tungstotellurate(VI) (three-letter code: TEW) (formula: O<sub>24</sub>TeW<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	Te	W		
3	B	1	31	24	1	6	0	0
3	D	1	31	24	1	6	0	0
3	F	1	31	24	1	6	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: High-affinity zinc uptake system protein ZnuA

Chain A: 




- Molecule 1: High-affinity zinc uptake system protein ZnuA


Chain B: 




- Molecule 1: High-affinity zinc uptake system protein ZnuA

Chain C: 







- Molecule 1: High-affinity zinc uptake system protein ZnuA



- Molecule 1: High-affinity zinc uptake system protein ZnuA



- Molecule 1: High-affinity zinc uptake system protein ZnuA



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.25Å 81.67Å 126.56Å 90.00° 112.99° 90.00°	Depositor
Resolution (Å)	47.47 – 3.15 47.42 – 3.15	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.47-3.15) 99.7 (47.42-3.15)	Depositor EDS
$R_{merge}$	0.22	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 3.12Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.217 , 0.254 0.209 , 0.252	Depositor DCC
$R_{free}$ test set	2006 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	89.3	Xtrriage
Anisotropy	0.037	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 78.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.125 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12243	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	106.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TEW, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/2062	0.79	1/2796 (0.0%)
1	B	0.60	1/2070 (0.0%)	0.82	1/2807 (0.0%)
1	C	0.61	0/2062	0.80	1/2796 (0.0%)
1	D	0.58	0/2070	0.79	2/2807 (0.1%)
1	E	0.59	0/2070	0.78	1/2807 (0.0%)
1	F	0.58	0/2062	0.79	1/2796 (0.0%)
All	All	0.59	1/12396 (0.0%)	0.80	7/16809 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	153	GLU	CG-CD	5.27	1.59	1.51

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	81	ILE	CG1-CB-CG2	5.98	124.55	111.40
1	C	81	ILE	CG1-CB-CG2	5.95	124.50	111.40
1	A	203	LYS	CD-CE-NZ	5.81	125.06	111.70
1	D	203	LYS	CD-CE-NZ	5.76	124.95	111.70
1	B	203	LYS	CD-CE-NZ	5.75	124.93	111.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	312	LYS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2020	0	2032	33	0
1	B	2028	0	2036	39	0
1	C	2020	0	2032	46	0
1	D	2028	0	2036	39	0
1	E	2028	0	2036	30	0
1	F	2020	0	2032	37	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	B	31	0	0	0	0
3	D	31	0	0	0	0
3	F	31	0	0	0	0
All	All	12243	0	12204	217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:LYS:HE3	1:A:91:LYS:HA	1.70	0.73
1:E:91:LYS:HE3	1:E:91:LYS:HA	1.71	0.72
1:B:91:LYS:HA	1:B:91:LYS:HE3	1.72	0.71
1:C:83:PRO:HB3	1:C:90:GLU:HG3	1.72	0.71
1:D:203:LYS:HG3	1:D:222:GLY:O	1.91	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/316 (82%)	228 (88%)	24 (9%)	7 (3%)	5	27
1	B	260/316 (82%)	230 (88%)	24 (9%)	6 (2%)	6	31
1	C	259/316 (82%)	226 (87%)	25 (10%)	8 (3%)	4	23
1	D	260/316 (82%)	230 (88%)	22 (8%)	8 (3%)	4	23
1	E	260/316 (82%)	231 (89%)	21 (8%)	8 (3%)	4	23
1	F	259/316 (82%)	228 (88%)	24 (9%)	7 (3%)	5	27
All	All	1557/1896 (82%)	1373 (88%)	140 (9%)	44 (3%)	5	26

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	ASP
1	A	266	ALA
1	B	266	ALA
1	C	61	ASP
1	C	266	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/251 (85%)	191 (90%)	22 (10%)	7	26

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	214/251 (85%)	189 (88%)	25 (12%)	5	22
1	C	213/251 (85%)	189 (89%)	24 (11%)	6	23
1	D	214/251 (85%)	191 (89%)	23 (11%)	6	25
1	E	214/251 (85%)	191 (89%)	23 (11%)	6	25
1	F	213/251 (85%)	188 (88%)	25 (12%)	5	22
All	All	1281/1506 (85%)	1139 (89%)	142 (11%)	6	24

5 of 142 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	281	THR
1	F	71	ARG
1	F	230	THR
1	C	71	ARG
1	C	53	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	302	GLN
1	E	101	GLN
1	F	302	GLN
1	F	101	GLN
1	F	220	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TEW	B	402	-	29,42,42	0.84	0	12,129,129	1.92	4 (33%)
3	TEW	D	402	-	29,42,42	0.82	0	12,129,129	1.97	4 (33%)
3	TEW	F	402	-	29,42,42	0.84	0	12,129,129	2.19	5 (41%)

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	B	402	TEW	O12-TE1-O7	-3.72	80.81	85.21
3	F	402	TEW	O12-TE1-O11	-3.61	80.94	85.21
3	F	402	TEW	O13-TE1-O1	3.46	97.77	94.65
3	D	402	TEW	O7-TE1-O1	-3.30	81.31	85.21
3	B	402	TEW	O13-TE1-O1	3.28	97.61	94.65

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/316 (83%)	-0.23	0 <b>100</b> <b>100</b>	66, 92, 120, 152	0
1	B	264/316 (83%)	-0.19	1 (0%) <b>92</b> <b>89</b>	69, 95, 127, 150	0
1	C	263/316 (83%)	-0.21	1 (0%) <b>92</b> <b>89</b>	72, 98, 130, 155	0
1	D	264/316 (83%)	-0.11	4 (1%) <b>73</b> <b>61</b>	69, 102, 162, 236	0
1	E	264/316 (83%)	-0.10	6 (2%) <b>60</b> <b>46</b>	72, 102, 170, 222	0
1	F	263/316 (83%)	-0.21	0 <b>100</b> <b>100</b>	72, 106, 143, 167	0
All	All	1581/1896 (83%)	-0.18	12 (0%) <b>86</b> <b>78</b>	66, 99, 144, 236	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	250	VAL	8.3
1	E	250	VAL	6.3
1	D	252	GLN	4.2
1	E	252	GLN	3.9
1	E	251	GLU	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	TEW	D	402	31/31	0.82	0.18	312,334,349,370	31
3	TEW	F	402	31/31	0.82	0.22	355,376,414,434	31
3	TEW	B	402	31/31	0.91	0.22	324,343,356,382	31
2	ZN	C	401	1/1	0.99	0.20	80,80,80,80	0
2	ZN	D	401	1/1	0.99	0.20	96,96,96,96	0
2	ZN	F	401	1/1	0.99	0.18	91,91,91,91	0
2	ZN	B	401	1/1	1.00	0.21	89,89,89,89	0
2	ZN	E	401	1/1	1.00	0.19	93,93,93,93	0
2	ZN	A	401	1/1	1.00	0.20	84,84,84,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around TEW D 402:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around TEW F 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.