



Full wwPDB EM Validation Report ⓘ

Sep 3, 2024 – 08:03 pm BST

PDB ID : 8RCK
EMDB ID : EMD-19053
Title : CryoEM structure of mTORC1 with a paediatric kidney cancer-associated 1455-EWED-1458 duplication in mTOR, Focused on one protomer copy.
Authors : Anandapadamanaban, M.; Hay, I.M.; Perisic, O.; Williams, R.L.
Deposited on : 2023-12-06
Resolution : 3.40 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev112
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	2549	
2	E	326	
3	Y	1335	
4	X	118	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 54652 atoms, of which 27400 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	B	2036	33036	10476	16667	2863	2926	104	0	0

- Molecule 2 is a protein called Target of rapamycin complex subunit LST8.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	E	317	4798	1526	2342	436	476	18	0	0

- Molecule 3 is a protein called Regulatory-associated protein of mTOR.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
3	Y	1044	16645	5322	8320	1441	1506	56	0	0

- Molecule 4 is a protein called Eukaryotic translation initiation factor 4E-binding protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
4	X	8	127	42	58	9	17	1	0	0

- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf	
			Total	C	H	N	O		P
5	B	1	44	10	13	6	12	3	0


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
6	B	2	2	2	0

3 Residue-property plots



These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serine/threonine-protein kinase mTOR





● Molecule 2: Target of rapamycin complex subunit LST8



- Molecule 4: Eukaryotic translation initiation factor 4E-binding protein 1

Chain X:
7%
93%

MET	GLU	SER	CYS	GLY	ARG	GLY	ASN	SER	ASN	SER	PRO	SER	VAL	SER	THR	CYS	LYS	SER	THR	PRO	PRO	ARG	ARG	ALA	LEU	ILE	PRO	THR	THR	ALA	ILE	PRO	PRO	ARG	GLY	ARG	GLY	VAL	VAL	THR	THR	SER	SER	PRO	PRO	GLY	GLY	ASP	SER	SER	SER	VAL	ASP	GLN	GLU	PRO	PRO	PRO	PRO	PRO	PRO	MET	MET	GLY	GLU	ALA	ALA	SER	SER	GLN	GLN	SER	SER	HIS	HIS	LEU	LEU	ARG	ARG	ASN	ASN	SER	SER	PRO	PRO	LEU	LEU	PRO	PRO	GLU	GLU	ASP	ASP	THR	THR	LYS	LYS	ARG	ARG	ALA	ALA	GLY	GLY	GLY	GLY	GLU	GLU	THR	THR	ARG	ARG	ILE	ILE	TYR	TYR	ASP	ASP	ARG	ARG	LYS	LYS	PHE	PHE	LEU	LEU	MET	MET		
GLU	GLU	CYS	CYS	ARG	ARG	ASN	ASN	SER	SER	PRO	PRO	VAL	VAL	THR	THR	LYS	LYS	THR	THR	PRO	PRO	ARG	ARG	ASP	ASP	LEU	LEU	THR	THR	ILE	ILE	PRO	PRO	ARG	ARG	GLY	GLY	VAL	VAL	THR	THR	SER	SER	PRO	PRO	GLY	GLY	ASP	ASP	SER	SER	VAL	VAL	GLN	GLN	GLU	GLU	PRO	PRO	PRO	PRO	PRO	PRO	MET	MET	GLU	GLU	ALA	ALA	SER	SER	GLN	GLN	SER	SER	HIS	HIS	LEU	LEU	ARG	ARG	ASN	ASN	SER	SER	PRO	PRO	LEU	LEU	PRO	PRO	GLU	GLU	ASP	ASP	THR	THR	LYS	LYS	ARG	ARG	ALA	ALA	GLY	GLY	GLY	GLY	GLU	GLU	THR	THR	ARG	ARG	ILE	ILE	TYR	TYR	ASP	ASP	ARG	ARG	LYS	LYS	PHE	PHE	LEU	LEU	MET	MET

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	177138	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.046	Depositor
Minimum map value	-0.021	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.005	Depositor
Map size (\AA)	440.32, 440.32, 440.32	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.86, 0.86, 0.86	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.31	0/16696	0.64	3/22592 (0.0%)
2	E	0.32	0/2514	0.70	2/3426 (0.1%)
3	Y	0.31	0/8524	0.63	1/11596 (0.0%)
4	X	0.40	0/69	0.48	0/89
All	All	0.31	0/27803	0.64	6/37703 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	12
2	E	0	3
3	Y	0	5
All	All	0	20

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	164	ARG	C-N-CA	10.32	147.51	121.70
1	B	2133	CYS	C-N-CA	8.79	143.66	121.70
1	B	2372	PRO	C-N-CA	7.24	139.81	121.70
3	Y	1205	MET	C-N-CA	6.68	138.41	121.70
2	E	222	TYR	C-N-CA	5.78	136.14	121.70
2	E	185	MET	C-N-CA	5.32	134.99	121.70

There are no chirality outliers.

All (20) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	104	ILE	Peptide
1	B	1271	ALA	Peptide
1	B	1522	GLY	Peptide
1	B	1580	GLY	Peptide
1	B	165	ARG	Peptide
1	B	1939	ILE	Peptide
1	B	287	GLU	Peptide
1	B	444	VAL	Peptide
1	B	459	ILE	Peptide
1	B	835	ALA	Peptide
1	B	837	ARG	Peptide
1	B	937	ASN	Peptide
2	E	113	ASP	Peptide
2	E	191	THR	Peptide
2	E	292	ASN	Peptide
3	Y	1098	ASN	Peptide
3	Y	1171	SER	Peptide
3	Y	1183	ALA	Peptide
3	Y	24	TRP	Peptide
3	Y	591	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	16369	16667	16646	29	0
2	E	2456	2342	2341	9	0
3	Y	8325	8320	8314	14	0
4	X	69	58	57	0	0
5	B	31	13	13	1	0
6	B	2	0	0	0	0
All	All	27252	27400	27371	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:245:PHE:O	1:B:264:ALA:N	2.26	0.69
3:Y:574:ARG:NH1	3:Y:610:ASP:OD2	2.29	0.66
1:B:1377:ASP:OD1	1:B:1406:LYS:NZ	2.29	0.66
1:B:2339:ARG:O	1:B:2545:TRP:NE1	2.30	0.64
1:B:719:SER:O	1:B:766:ARG:NH2	2.32	0.63
1:B:481:THR:O	1:B:483:PHE:N	2.33	0.61
1:B:1427:GLU:OE2	1:B:2218:LYS:NZ	2.22	0.61
1:B:91:ALA:O	1:B:132:ARG:NH1	2.35	0.59
3:Y:527:MET:O	3:Y:532:ARG:NH1	2.35	0.59
1:B:1362:GLU:OE1	1:B:1395:LYS:NZ	2.36	0.58
3:Y:585:TRP:O	3:Y:629:ASN:ND2	2.37	0.57
1:B:502:LYS:NZ	1:B:534:ASP:OD2	2.41	0.54
3:Y:33:LYS:NZ	3:Y:1155:GLU:OE2	2.25	0.52
1:B:1004:CYS:O	1:B:1009:ARG:NH2	2.44	0.51
2:E:106:ASP:O	2:E:108:THR:N	2.44	0.50
1:B:1438:LYS:NZ	1:B:2393:ASP:OD2	2.32	0.48
2:E:102:THR:HG1	2:E:112:TRP:HZ3	1.60	0.48
3:Y:1178:ARG:NH2	3:Y:1223:PRO:O	2.45	0.47
1:B:870:ASN:O	1:B:874:THR:N	2.47	0.47
2:E:304:ILE:HB	2:E:305:LYS:HB2	1.97	0.47
1:B:2546:CYS:HG	1:B:2549:TRP:HE3	1.63	0.47
1:B:2133:CYS:HA	1:B:2134:ARG:HB2	1.98	0.46
2:E:87:ASN:ND2	2:E:105:GLU:OE1	2.46	0.45
2:E:185:MET:HB3	2:E:186:ALA:CA	2.47	0.45
1:B:203:GLN:OE1	1:B:206:ARG:NE	2.46	0.44
2:E:113:ASP:OD2	2:E:122:GLN:NE2	2.50	0.44
2:E:185:MET:HB3	2:E:186:ALA:HA	1.99	0.44
3:Y:198:ASN:ND2	3:Y:248:ALA:O	2.51	0.43
3:Y:1088:THR:OG1	3:Y:1090:ASP:OD1	2.34	0.43
1:B:878:GLN:OE1	1:B:881:ARG:NH2	2.51	0.43
1:B:886:ARG:NH2	1:B:1565:ASP:OD1	2.47	0.43
1:B:2240:VAL:HG23	5:B:3000:ANP:HN62	1.83	0.43
2:E:192:GLY:N	2:E:222:TYR:O	2.47	0.43
3:Y:823:TYR:O	3:Y:826:VAL:N	2.52	0.43
3:Y:79:THR:HB	3:Y:80:PRO:HD3	2.01	0.42
3:Y:1235:ASP:OD1	3:Y:1254:LYS:NZ	2.50	0.42
1:B:947:TYR:HB2	1:B:948:PRO:HD3	2.01	0.42
1:B:1548:ALA:O	1:B:1550:LEU:N	2.53	0.42
3:Y:271:ILE:HG22	3:Y:272:LYS:H	1.84	0.41
1:B:719:SER:OG	1:B:763:ASN:O	2.37	0.41
1:B:775:ILE:O	1:B:779:LEU:N	2.38	0.41
2:E:62:TYR:O	2:E:64:HIS:ND1	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:Y:616:ARG:NE	3:Y:654:ASP:OD2	2.51	0.41
1:B:460:ARG:HE	1:B:504:LEU:HD21	1.85	0.41
1:B:123:MET:HG3	1:B:170:LEU:HD12	2.03	0.41
1:B:69:HIS:NE2	1:B:73:GLU:OE2	2.54	0.41
1:B:849:ALA:O	1:B:1611:ARG:NH2	2.47	0.41
1:B:2368:ARG:NE	1:B:2373:GLU:OE2	2.40	0.41
1:B:1528:SER:OG	1:B:1529:MET:N	2.55	0.41
3:Y:1205:MET:CB	3:Y:1206:THR:HA	2.51	0.41
1:B:2189:HIS:N	1:B:2233:ASN:O	2.54	0.40
3:Y:35:HIS:O	3:Y:962:TRP:NE1	2.45	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	1994/2549 (78%)	1928 (97%)	56 (3%)	10 (0%)	25	54
2	E	315/326 (97%)	297 (94%)	17 (5%)	1 (0%)	37	66
3	Y	1032/1335 (77%)	996 (96%)	33 (3%)	3 (0%)	37	66
4	X	6/118 (5%)	6 (100%)	0	0	100	100
All	All	3347/4328 (77%)	3227 (96%)	106 (3%)	14 (0%)	32	60

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	165	ARG
1	B	1311	ASN
1	B	1572	ASP
3	Y	271	ILE
3	Y	634	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1028	ARG
1	B	1116	PRO
1	B	2134	ARG
1	B	1580	GLY
2	E	292	ASN
3	Y	272	LYS
1	B	1939	ILE
1	B	482	VAL
1	B	1115	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	1789/2220 (81%)	1789 (100%)	0	100	100
2	E	269/276 (98%)	269 (100%)	0	100	100
3	Y	921/1163 (79%)	921 (100%)	0	100	100
4	X	8/102 (8%)	8 (100%)	0	100	100
All	All	2987/3761 (79%)	2987 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	536	GLN
1	B	1264	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ANP	B	3000	6	29,33,33	4.64	15 (51%)	31,52,52	2.95	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ANP	B	3000	6	-	5/14/38/38	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	3000	ANP	C2'-C1'	-16.39	1.28	1.53
5	B	3000	ANP	O4'-C1'	10.62	1.55	1.41
5	B	3000	ANP	C3'-C4'	-9.94	1.27	1.53
5	B	3000	ANP	C2'-C3'	4.58	1.65	1.53
5	B	3000	ANP	O4'-C4'	4.57	1.55	1.45
5	B	3000	ANP	PG-O1G	3.89	1.52	1.46
5	B	3000	ANP	PB-O3A	3.71	1.63	1.59
5	B	3000	ANP	PB-N3B	3.32	1.72	1.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	3000	ANP	C6-N6	3.03	1.45	1.34
5	B	3000	ANP	PB-O1B	2.64	1.50	1.46
5	B	3000	ANP	C5-C4	-2.53	1.34	1.40
5	B	3000	ANP	PG-N3B	2.45	1.69	1.63
5	B	3000	ANP	PB-O2B	-2.14	1.51	1.56
5	B	3000	ANP	C5'-C4'	2.11	1.58	1.51
5	B	3000	ANP	PG-O3G	-2.06	1.51	1.56

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	3000	ANP	C5-C6-N6	9.96	135.49	120.35
5	B	3000	ANP	N6-C6-N1	-7.00	104.04	118.57
5	B	3000	ANP	C1'-N9-C4	-6.66	114.93	126.64
5	B	3000	ANP	N3-C2-N1	-5.66	119.84	128.68
5	B	3000	ANP	C3'-C2'-C1'	4.15	107.22	100.98
5	B	3000	ANP	PB-O3A-PA	-2.25	124.69	132.62

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	3000	ANP	PG-N3B-PB-O1B
5	B	3000	ANP	O4'-C4'-C5'-O5'
5	B	3000	ANP	C3'-C4'-C5'-O5'
5	B	3000	ANP	PB-O3A-PA-O1A
5	B	3000	ANP	PB-O3A-PA-O2A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	3000	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19053. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256




Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256



Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 271




Y Index: 226




Z Index: 237


6.3.2 Raw map



X Index: 256



Y Index: 226




Z Index: 231


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.005. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_19053_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 273 nm³; this corresponds to an approximate mass of 247 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.37	3.97	3.40
Unmasked-calculated*	4.05	6.86	4.11

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.05 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-19053 and PDB model 8RCK. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.005 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.005).











9.4 Atom inclusion [i](#)



At the recommended contour level, 66% of all backbone atoms, 66% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.005) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6620	 0.3050
B	 0.6630	 0.3100
E	 0.7330	 0.2760
X	 0.8410	 0.4630
Y	 0.6610	 0.3030

