



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2024 – 08:01 PM EST

PDB ID : 3RF6
Title : Crystal structure of glycerol-3 phosphate bound HAD-like phosphatase from *Saccharomyces cerevisiae*
Authors : Nocek, B.; Kuznetsova, K.; Evdokimova, E.; Savchenko, A.; Iakunine, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2011-04-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	 6% 91% 5% .
1	B	355	 5% 95% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	G3P	A	501	X	-	-	-
3	G3P	B	501	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Uncharacterized protein YKR070W.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	340	2659	1700	450	504	5	0	4	0
1	B	352	2764	1775	469	514	6	0	3	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP P36151
A	-1	ASN	-	expression tag	UNP P36151
A	0	ALA	-	expression tag	UNP P36151
A	19	ALA	ASP	engineered mutation	UNP P36151
B	-2	SER	-	expression tag	UNP P36151
B	-1	ASN	-	expression tag	UNP P36151
B	0	ALA	-	expression tag	UNP P36151
B	19	ALA	ASP	engineered mutation	UNP P36151

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is SN-GLYCEROL-3-PHOSPHATE (three-letter code: G3P) (formula: $C_3H_9O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		
4	B	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	261	Total	O	0	0
			261	261		
5	B	284	Total	O	0	0
			284	284		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein YKR070W



- Molecule 1: Uncharacterized protein YKR070W



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.91Å 71.65Å 195.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.24 – 1.70 35.24 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.4 (35.24-1.70) 99.5 (35.24-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.70Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.164 , 0.194 0.164 , 0.194	Depositor DCC
R_{free} test set	4701 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtrriage
Anisotropy	0.392	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6017	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G3P, FLC, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/2732	0.75	1/3717 (0.0%)
1	B	0.75	2/2836 (0.1%)	0.75	0/3850
All	All	0.72	2/5568 (0.0%)	0.75	1/7567 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	331	CYS	CB-SG	-10.54	1.64	1.82
1	B	143	TYR	CD1-CE1	5.93	1.48	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	115	ASP	CB-CG-OD1	5.83	123.55	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2659	0	2607	17	0
1	B	2764	0	2750	9	0
2	A	13	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	13	0	5	0	0
3	A	10	0	7	0	0
3	B	10	0	7	0	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	A	261	0	0	1	0
5	B	284	0	0	2	0
All	All	6017	0	5381	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:LYS:HG2	1:A:184:MET:SD	2.17	0.85
1:A:154:LYS:CG	1:A:184:MET:SD	2.81	0.69
1:B:70:LEU:O	1:B:72[A]:VAL:HG23	1.96	0.65
1:B:39:LYS:HG2	1:B:72[B]:VAL:HG12	1.85	0.57
1:A:260:ILE:HD11	1:A:289:PRO:HD2	1.89	0.53
1:B:323:ASN:ND2	5:B:455:HOH:O	2.36	0.52
1:A:260:ILE:HD12	1:A:289:PRO:CG	2.39	0.52
1:A:60:ARG:O	1:A:64:GLU:HG3	2.10	0.51
1:B:350:LYS:HD3	1:B:351:TYR:CE2	2.47	0.49
1:A:260:ILE:CD1	1:A:289:PRO:HD2	2.47	0.45
1:B:141:MET:HG2	5:B:581:HOH:O	2.17	0.45
1:A:280:LEU:O	1:A:283:THR:HG22	2.16	0.45
1:A:334:THR:HG22	1:A:335:LEU:HD12	1.99	0.44
1:A:260:ILE:HD13	1:A:290:PHE:HE2	1.81	0.44
1:A:154:LYS:HG3	1:A:184:MET:SD	2.58	0.44
1:A:211:ASN:CB	1:A:212:PRO:CD	2.96	0.43
1:A:259:LEU:HG	1:A:290:PHE:CE2	2.53	0.43
1:A:334:THR:C	1:A:335:LEU:HD12	2.39	0.43
1:A:211:ASN:HB2	1:A:212:PRO:CD	2.48	0.42
1:A:260:ILE:HD13	1:A:290:PHE:CE2	2.54	0.42
1:B:281:LEU:HD12	1:B:281:LEU:HA	1.87	0.42
1:B:39:LYS:CG	1:B:72[B]:VAL:HG12	2.48	0.41
1:A:181:GLU:HG2	5:A:523:HOH:O	2.19	0.41
1:B:39:LYS:HG3	1:B:72[B]:VAL:CG1	2.50	0.41
1:B:191:GLU:N	1:B:191:GLU:OE1	2.54	0.41
1:A:335:LEU:HD12	1:A:335:LEU:N	2.36	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/355 (96%)	331 (97%)	10 (3%)	1 (0%)	41	24
1	B	353/355 (99%)	346 (98%)	6 (2%)	1 (0%)	41	24
All	All	695/710 (98%)	677 (97%)	16 (2%)	2 (0%)	41	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	211	ASN
1	A	211	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/306 (94%)	285 (99%)	2 (1%)	84	77
1	B	298/306 (97%)	297 (100%)	1 (0%)	92	89
All	All	585/612 (96%)	582 (100%)	3 (0%)	88	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	272	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	281	LEU
1	B	114	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	G3P	A	501	4	9,9,9	0.61	0	11,12,12	0.63	0
2	FLC	A	401	-	12,12,12	2.12	3 (25%)	17,17,17	1.61	3 (17%)
3	G3P	B	501	4	9,9,9	1.59	1 (11%)	11,12,12	0.89	0
2	FLC	B	402	-	12,12,12	1.75	4 (33%)	17,17,17	1.46	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G3P	A	501	4	1/1/2/2	0/8/8/8	-
3	G3P	B	501	4	1/1/2/2	4/8/8/8	-
2	FLC	B	402	-	-	11/16/16/16	-
2	FLC	A	401	-	-	0/16/16/16	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FLC	CA-CB	-4.32	1.48	1.53
3	B	501	G3P	P-O2P	-3.73	1.38	1.50
2	A	401	FLC	CB-CBC	-3.45	1.49	1.53
2	B	402	FLC	CG-CB	-3.35	1.49	1.53
2	A	401	FLC	CG-CB	-3.05	1.50	1.53
2	B	402	FLC	CB-CBC	-2.25	1.51	1.53
2	B	402	FLC	CA-CB	-2.19	1.51	1.53
2	B	402	FLC	OHB-CB	-2.06	1.39	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FLC	OB2-CBC-CB	3.60	119.31	113.05
2	B	402	FLC	OB2-CBC-CB	3.34	118.85	113.05
2	B	402	FLC	CB-CG-CGC	-2.45	107.88	113.81
2	A	401	FLC	CB-CA-CAC	-2.35	108.12	113.81
2	A	401	FLC	OB1-CBC-CB	-2.11	119.27	122.25

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	501	G3P	C2
3	B	501	G3P	C2

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	402	FLC	CAC-CA-CB-CG
2	B	402	FLC	CAC-CA-CB-OHB
2	B	402	FLC	OHB-CB-CBC-OB1
3	B	501	G3P	O2-C2-C3-O1P
2	B	402	FLC	CAC-CA-CB-CBC

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	501	G3P	C1-C2-C3-O1P
2	B	402	FLC	OHB-CB-CBC-OB2
2	B	402	FLC	CG-CB-CBC-OB1
2	B	402	FLC	CG-CB-CBC-OB2
3	B	501	G3P	O1-C1-C2-O2
2	B	402	FLC	CA-CB-CBC-OB2
2	B	402	FLC	CA-CB-CBC-OB1
3	B	501	G3P	C3-O1P-P-O4P
2	B	402	FLC	CB-CG-CGC-OG1
2	B	402	FLC	CB-CG-CGC-OG2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/355 (95%)	0.14	23 (6%) 17 19	16, 31, 74, 107	0
1	B	352/355 (99%)	0.01	16 (4%) 33 37	17, 28, 69, 111	0
All	All	692/710 (97%)	0.07	39 (5%) 24 27	16, 29, 73, 111	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	270	ILE	6.7
1	A	274	VAL	6.0
1	B	273	SER	5.7
1	A	287	THR	5.7
1	A	268	GLY	4.6
1	A	286	SER	4.5
1	A	271	GLY	4.4
1	A	272	GLN	4.0
1	B	274	VAL	3.9
1	B	268	GLY	3.9
1	B	272	GLN	3.9
1	A	267	SER	3.9
1	A	273	SER	3.7
1	B	352	ALA	3.6
1	B	10	THR	3.5
1	B	270	ILE	3.4
1	A	269	LYS	3.4
1	A	278	LEU	3.3
1	B	351	TYR	3.2
1	B	9	THR	3.1
1	B	267	SER	3.1
1	A	288	SER	3.0
1	A	280	LEU	2.9
1	A	266	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	269	LYS	2.8
1	B	7	PHE	2.7
1	A	327	ASP	2.6
1	A	275	LYS	2.6
1	B	271	GLY	2.5
1	A	276	GLN	2.4
1	B	11	SER	2.4
1	B	223	PHE	2.4
1	B	6	PHE	2.3
1	A	291	HIS	2.3
1	A	40	LEU	2.2
1	A	283	THR	2.2
1	A	289	PRO	2.1
1	A	329	LYS	2.1
1	A	285	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	A	701	1/1	0.77	0.13	32,32,32,32	1
2	FLC	B	402	13/13	0.88	0.18	26,60,75,77	0
4	MG	B	701	1/1	0.88	0.11	31,31,31,31	1
3	G3P	A	501	10/10	0.91	0.20	27,33,42,46	10
2	FLC	A	401	13/13	0.94	0.15	27,45,56,58	0
3	G3P	B	501	10/10	0.95	0.18	23,28,41,48	10
4	MG	A	353	1/1	0.99	0.07	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.