



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2023 – 02:32 PM EDT

PDB ID : 4RJV  
Title : Crystal Structure of a De Novo Designed Ferredoxin Fold, Northeast Structural Genomics Consortium (NESG) Target OR461  
Authors : O'Connell, P.T.; Lin, Y.-R.; Guan, R.; Koga, N.; Koga, R.; Seetharaman, J.; Janjua, H.; Xiao, R.; Maglaqui, M.; Everett, J.K.; Acton, T.B.; Baker, D.; Montelione, G.T.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2014-10-09  
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	87	
1	B	87	
1	C	87	
1	D	87	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called OR461.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	83	657	408	121	125	3	0	0	0
1	B	80	628	391	114	120	3	0	0	0
1	C	79	620	385	113	119	3	0	0	0
1	D	83	657	408	121	125	3	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	54	Total	O	0	0
			54	54		
2	B	50	Total	O	0	0
			50	50		
2	C	45	Total	O	0	0
			45	45		
2	D	47	Total	O	0	0
			47	47		





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	37.67Å 46.69Å 48.07Å 86.49° 72.89° 72.33°	Depositor
Resolution (Å)	30.17 – 1.52 30.17 – 1.52	Depositor EDS
% Data completeness (in resolution range)	93.2 (30.17-1.52) 93.5 (30.17-1.52)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.72 (at 1.52Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.199 , 0.231 0.199 , 0.231	Depositor DCC
$R_{free}$ test set	2012 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.2	Xtrriage
Anisotropy	0.438	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.030 for -h,-l,-k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2758	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5610e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/663	0.52	0/888
1	B	0.38	0/632	0.63	1/846 (0.1%)
1	C	0.37	0/624	0.53	0/835
1	D	0.34	0/663	0.53	0/888
All	All	0.36	0/2582	0.55	1/3457 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3	MET	CG-SD-CE	-9.54	84.94	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	657	0	668	15	0
1	B	628	0	648	10	0
1	C	620	0	637	7	0
1	D	657	0	668	6	0
2	A	54	0	0	5	1
2	B	50	0	0	1	1
2	C	45	0	0	2	0
2	D	47	0	0	4	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2758	0	2621	35	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:25:ARG:NH1	2:D:144:HOH:O	1.98	0.92
1:A:22:MET:SD	2:A:142:HOH:O	2.31	0.88
1:A:34:ARG:HG2	1:B:34:ARG:HG2	1.54	0.88
1:C:45:ARG:NH1	2:C:127:HOH:O	2.18	0.76
1:C:25:ARG:NH2	1:C:56:GLU:OE1	2.28	0.67
1:A:4:ASP:OD2	1:A:77:ARG:NH1	2.26	0.67
1:D:6:ARG:NH1	2:D:126:HOH:O	2.30	0.65
1:A:71:THR:OG1	2:A:112:HOH:O	2.13	0.64
1:B:3:MET:HG3	1:B:4:ASP:N	2.13	0.64
1:A:6:ARG:NH1	1:A:41:ASP:OD2	2.32	0.62
1:A:56:GLU:HB2	2:A:121:HOH:O	2.01	0.60
1:B:6:ARG:NH1	1:B:38:ASP:OD2	2.38	0.56
1:C:36:THR:OG1	1:C:43:GLU:OE1	2.23	0.56
1:B:66:LYS:NZ	2:B:143:HOH:O	2.42	0.52
1:C:6:ARG:NE	2:C:124:HOH:O	2.33	0.51
1:B:3:MET:CE	1:B:58:ALA:HB2	2.43	0.49
1:D:63:ARG:HA	1:D:66:LYS:HD3	1.94	0.49
1:C:45:ARG:HB3	1:C:45:ARG:CZ	2.43	0.48
1:A:17:ARG:HD2	1:B:13:GLU:HG3	1.96	0.47
1:B:3:MET:HE1	1:B:58:ALA:HB2	1.94	0.47
1:A:58:ALA:O	1:A:74:ARG:NH2	2.48	0.47
1:A:34:ARG:HG2	1:B:34:ARG:CG	2.36	0.46
1:B:21:LYS:HD2	1:B:68:PHE:CZ	2.50	0.46
1:A:51:GLU:OE1	2:A:146:HOH:O	2.20	0.45
1:A:22:MET:HA	2:A:142:HOH:O	2.16	0.45
1:A:3:MET:HB3	1:A:54:LEU:HD22	2.00	0.44
1:D:6:ARG:HE	1:D:6:ARG:HB2	1.31	0.44
1:A:65:ALA:HB2	1:A:72:VAL:HG12	2.00	0.43
1:D:8:ARG:NH1	2:D:140:HOH:O	2.51	0.43
1:B:21:LYS:HD2	1:B:68:PHE:HZ	1.82	0.43
1:A:21:LYS:HZ3	1:A:68:PHE:HZ	1.67	0.42
1:D:37:ASP:OD2	2:D:121:HOH:O	2.22	0.41
1:C:5:ILE:CD1	1:C:74:ARG:HG2	2.50	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:ASP:O	1:A:15:LEU:HD23	2.20	0.41
1:C:2:GLU:OE2	1:C:45:ARG:NH2	2.53	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:145:HOH:O	2:D:143:HOH:O[1_644]	1.84	0.36
2:B:118:HOH:O	2:D:107:HOH:O[1_554]	2.13	0.07

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	81/87 (93%)	79 (98%)	2 (2%)	0	100	100
1	B	78/87 (90%)	78 (100%)	0	0	100	100
1	C	77/87 (88%)	76 (99%)	1 (1%)	0	100	100
1	D	81/87 (93%)	81 (100%)	0	0	100	100
All	All	317/348 (91%)	314 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	69/73 (94%)	68 (99%)	1 (1%)	67	41
1	B	66/73 (90%)	64 (97%)	2 (3%)	41	12
1	C	65/73 (89%)	65 (100%)	0	100	100
1	D	69/73 (94%)	67 (97%)	2 (3%)	42	13
All	All	269/292 (92%)	264 (98%)	5 (2%)	57	26

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	ARG
1	B	10	ASP
1	B	74	ARG
1	D	1	MET
1	D	19	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	83/87 (95%)	2.59	41 (49%) 0 0	20, 37, 64, 89	0
1	B	80/87 (91%)	2.24	40 (50%) 0 0	17, 29, 57, 79	0
1	C	79/87 (90%)	2.34	39 (49%) 0 0	20, 32, 59, 77	0
1	D	83/87 (95%)	2.63	43 (51%) 0 0	17, 35, 66, 84	0
All	All	325/348 (93%)	2.46	163 (50%) 0 0	17, 34, 64, 89	0

All (163) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	40	ASN	12.7
1	A	39	GLY	11.9
1	D	40	ASN	11.3
1	D	38	ASP	11.0
1	A	83	HIS	10.3
1	D	83	HIS	10.3
1	C	40	ASN	9.5
1	B	40	ASN	8.7
1	C	39	GLY	8.7
1	A	38	ASP	8.7
1	D	39	GLY	8.4
1	A	37	ASP	7.3
1	C	79	SER	6.7
1	D	41	ASP	6.7
1	B	38	ASP	6.7
1	B	39	GLY	5.9
1	A	41	ASP	5.6
1	C	10	ASP	5.5
1	C	38	ASP	5.5
1	D	37	ASP	5.2
1	B	41	ASP	4.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	44	ILE	4.8
1	B	80	LEU	4.7
1	D	65	ALA	4.6
1	D	70	ILE	4.4
1	A	10	ASP	4.4
1	B	5	ILE	4.4
1	A	46	ILE	4.3
1	D	46	ILE	4.3
1	D	82	HIS	4.2
1	D	72	VAL	4.2
1	C	46	ILE	4.2
1	D	10	ASP	4.0
1	B	37	ASP	4.0
1	B	10	ASP	4.0
1	D	42	LEU	3.9
1	A	80	LEU	3.9
1	B	19	LEU	3.9
1	C	33	VAL	3.8
1	D	71	THR	3.8
1	A	33	VAL	3.8
1	D	33	VAL	3.8
1	C	5	ILE	3.8
1	C	37	ASP	3.7
1	B	46	ILE	3.7
1	A	8	ARG	3.6
1	D	19	LEU	3.6
1	A	44	ILE	3.5
1	C	19	LEU	3.5
1	D	23	VAL	3.5
1	B	44	ILE	3.5
1	D	8	ARG	3.4
1	D	53	VAL	3.4
1	D	5	ILE	3.4
1	A	49	VAL	3.4
1	D	35	PHE	3.4
1	C	64	LEU	3.4
1	C	12	ASP	3.3
1	A	82	HIS	3.3
1	C	73	THR	3.3
1	D	50	PRO	3.3
1	B	23	VAL	3.3
1	C	41	ASP	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	71	THR	3.3
1	D	44	ILE	3.2
1	C	72	VAL	3.2
1	C	35	PHE	3.2
1	A	23	VAL	3.2
1	A	76	ILE	3.2
1	A	61	ALA	3.2
1	A	15	LEU	3.2
1	D	76	ILE	3.2
1	C	11	ASP	3.1
1	B	42	LEU	3.1
1	D	28	SER	3.1
1	B	72	VAL	3.1
1	A	5	ILE	3.1
1	D	15	LEU	3.1
1	A	7	PHE	3.1
1	B	51	GLU	3.1
1	B	33	VAL	3.0
1	D	3	MET	3.0
1	A	57	LEU	3.0
1	D	27	VAL	3.0
1	B	35	PHE	2.9
1	B	61	ALA	2.9
1	A	70	ILE	2.9
1	A	53	VAL	2.9
1	D	14	ALA	2.9
1	C	23	VAL	2.9
1	B	7	PHE	2.9
1	D	68	PHE	2.9
1	D	66	LYS	2.9
1	A	71	THR	2.9
1	C	30	GLY	2.9
1	C	18	ALA	2.9
1	A	72	VAL	2.8
1	C	3	MET	2.8
1	B	18	ALA	2.8
1	C	22	MET	2.8
1	A	52	GLN	2.8
1	B	12	ASP	2.8
1	B	70	ILE	2.8
1	B	68	PHE	2.8
1	D	54	LEU	2.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	22	MET	2.7
1	C	7	PHE	2.7
1	C	42	LEU	2.7
1	D	20	LEU	2.7
1	B	15	LEU	2.7
1	D	12	ASP	2.6
1	B	79	SER	2.6
1	C	65	ALA	2.6
1	D	9	GLY	2.6
1	A	30	GLY	2.6
1	A	65	ALA	2.6
1	D	52	GLN	2.6
1	A	3	MET	2.5
1	B	65	ALA	2.5
1	A	54	LEU	2.5
1	C	57	LEU	2.5
1	D	64	LEU	2.5
1	C	61	ALA	2.5
1	A	66	LYS	2.5
1	B	75	THR	2.4
1	D	67	GLU	2.4
1	B	16	ALA	2.4
1	A	36	THR	2.4
1	B	29	PHE	2.4
1	C	55	LYS	2.4
1	B	53	VAL	2.3
1	A	18	ALA	2.3
1	D	11	ASP	2.3
1	A	50	PRO	2.3
1	B	11	ASP	2.3
1	A	19	LEU	2.3
1	A	6	ARG	2.3
1	D	22	MET	2.2
1	B	55	LYS	2.2
1	C	8	ARG	2.2
1	B	64	LEU	2.2
1	C	51	GLU	2.2
1	B	76	ILE	2.2
1	D	36	THR	2.2
1	A	20	LEU	2.2
1	C	76	ILE	2.1
1	B	54	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	75	THR	2.1
1	A	58	ALA	2.1
1	A	35	PHE	2.1
1	B	57	LEU	2.1
1	B	73	THR	2.1
1	B	27	VAL	2.1
1	C	24	VAL	2.1
1	C	15	LEU	2.1
1	C	16	ALA	2.1
1	C	54	LEU	2.1
1	D	81	GLU	2.0
1	C	53	VAL	2.0
1	C	78	GLY	2.0
1	A	11	ASP	2.0
1	A	75	THR	2.0
1	B	1	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.