



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 04:11 pm BST

PDB ID : 1RN1
Title : THREE-DIMENSIONAL STRUCTURE OF GLN 25-RIBONUCLEASE T1 AT 1.84 ANGSTROMS RESOLUTION: STRUCTURAL VARIATIONS AT THE BASE RECOGNITION AND CATALYTIC SITES
Authors : Arni, R.K.; Pal, G.P.; Ravichandran, K.G.; Tulinsky, A.; Walz Junior, F.G.; Metcalf, P.
Deposited on : 1991-11-22
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	104	76% 18% 5% •
1	B	104	78% 19% ••
1	C	104	82% 17% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	105	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE T1 ISOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	103	Total 753	C 462	N 124	O 163	S 4	0	0	0
1	B	103	Total 771	C 474	N 125	O 168	S 4	0	0	0
1	C	104	Total 777	C 478	N 126	O 169	S 4	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0
2	C	1	Total 5	O 4	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	50	Total 50	O 50	0	0
3	B	67	Total 67	O 67	0	0
3	C	62	Total 62	O 62	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: RIBONUCLEASE T1 ISOZYME

Chain A: 




- Molecule 1: RIBONUCLEASE T1 ISOZYME

Chain B: 



- Molecule 1: RIBONUCLEASE T1 ISOZYME

Chain C: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.71Å 37.54Å 77.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.84	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-1.84)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, X-PLOR	Depositor
R, R_{free}	0.144 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2495	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.16	3/772 (0.4%)	1.82	17/1050 (1.6%)
1	B	1.19	2/792 (0.3%)	1.67	10/1079 (0.9%)
1	C	1.26	7/798 (0.9%)	1.68	8/1088 (0.7%)
All	All	1.20	12/2362 (0.5%)	1.73	35/3217 (1.1%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	31	GLU	CD-OE1	7.12	1.33	1.25
1	A	46	GLU	CD-OE1	6.45	1.32	1.25
1	C	71	GLY	N-CA	6.45	1.55	1.46
1	B	28	GLU	CD-OE2	6.29	1.32	1.25
1	C	28	GLU	CD-OE1	6.18	1.32	1.25
1	C	46	GLU	CD-OE2	6.02	1.32	1.25
1	C	102	GLU	CD-OE1	5.92	1.32	1.25
1	A	82	GLU	CD-OE2	5.78	1.32	1.25
1	B	82	GLU	CD-OE2	5.64	1.31	1.25
1	C	58	GLU	CD-OE1	5.62	1.31	1.25
1	A	28	GLU	CD-OE1	5.52	1.31	1.25
1	C	82	GLU	CD-OE2	5.24	1.31	1.25

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	GLU	OE1-CD-OE2	11.22	136.77	123.30
1	C	4	TYR	CB-CG-CD2	-10.72	114.57	121.00
1	A	3	ASP	CB-CG-OD2	9.53	126.88	118.30
1	A	77	ARG	NE-CZ-NH2	-9.16	115.72	120.30
1	A	66	ASP	CB-CG-OD1	-8.87	110.32	118.30
1	A	24	TYR	CB-CG-CD1	-8.57	115.86	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	68	TYR	CB-CG-CD2	-8.16	116.10	121.00
1	A	56	TYR	CB-CG-CD2	-7.52	116.49	121.00
1	A	102	GLU	OE1-CD-OE2	7.40	132.18	123.30
1	C	77	ARG	NE-CZ-NH1	7.17	123.89	120.30
1	A	56	TYR	CB-CG-CD1	6.60	124.96	121.00
1	C	33	VAL	CB-CA-C	-6.48	99.10	111.40
1	C	75	ALA	N-CA-CB	-6.47	101.04	110.10
1	B	2	CYS	O-C-N	6.28	132.75	122.70
1	B	28	GLU	OE1-CD-OE2	6.14	130.66	123.30
1	B	28	GLU	CG-CD-OE2	-6.12	106.07	118.30
1	A	103	CYS	CA-CB-SG	-5.95	103.30	114.00
1	B	68	TYR	CG-CD2-CE2	-5.76	116.69	121.30
1	B	49	ASP	CB-CG-OD2	5.63	123.37	118.30
1	A	44	ASN	CA-C-O	-5.59	108.36	120.10
1	C	102	GLU	OE1-CD-OE2	5.56	129.97	123.30
1	B	46	GLU	CG-CD-OE1	5.48	129.27	118.30
1	B	102	GLU	CB-CA-C	-5.45	99.50	110.40
1	A	38	TYR	CB-CG-CD2	-5.43	117.74	121.00
1	B	88	GLY	O-C-N	5.26	131.11	122.70
1	C	70	GLY	C-N-CA	-5.23	111.31	122.30
1	A	25	GLN	CB-CG-CD	5.19	125.09	111.60
1	A	24	TYR	CG-CD1-CE1	-5.16	117.17	121.30
1	B	82	GLU	CA-CB-CG	5.16	124.75	113.40
1	C	4	TYR	CB-CG-CD1	5.08	124.05	121.00
1	A	3	ASP	CB-CA-C	5.08	120.56	110.40
1	A	54	SER	N-CA-CB	-5.05	102.93	110.50
1	A	82	GLU	CG-CD-OE2	-5.02	108.26	118.30
1	C	103	CYS	CA-CB-SG	-5.02	104.97	114.00
1	A	100	PHE	CB-CA-C	-5.01	100.39	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	753	0	640	8	0
1	B	771	0	657	10	0
1	C	777	0	662	3	0
2	A	5	0	0	2	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	50	0	0	1	0
3	B	67	0	0	4	0
3	C	62	0	0	2	0
All	All	2495	0	1959	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:SER:O	1:A:99:ASN:HB2	1.76	0.85
1:B:27:HIS:HD2	3:B:120:HOH:O	1.58	0.84
1:B:55:PRO:HG2	1:B:82:GLU:HG2	1.63	0.79
1:B:42:TYR:CE1	3:B:126:HOH:O	2.45	0.69
1:B:102:GLU:HG3	3:B:129:HOH:O	1.96	0.65
1:B:45:TYR:OH	3:B:154:HOH:O	2.16	0.62
1:C:55:PRO:HD2	3:C:129:HOH:O	2.01	0.61
1:B:81:ASN:HD21	1:B:85:GLN:NE2	1.99	0.60
1:A:20:GLN:NE2	3:A:126:HOH:O	2.37	0.57
1:C:25:GLN:NE2	3:C:164:HOH:O	2.41	0.54
1:A:92:HIS:CE1	1:A:100:PHE:CZ	2.96	0.53
1:B:81:ASN:HD21	1:B:85:GLN:HE21	1.56	0.53
1:B:36:ASN:HB2	1:B:38:TYR:CE2	2.43	0.53
1:A:92:HIS:CE1	1:A:100:PHE:CE1	3.03	0.46
1:A:22:ALA:O	1:A:26:LEU:HD13	2.15	0.45
1:B:12:SER:O	1:B:15:ASP:HB2	2.16	0.45
1:A:77:ARG:NH2	2:A:105:SO4:O1	2.48	0.43
1:C:59:TRP:HA	1:C:60:PRO:HD3	1.91	0.42
1:A:38:TYR:OH	2:A:105:SO4:O3	2.29	0.41
1:A:54:SER:HB3	1:A:56:TYR:CE2	2.55	0.41
1:B:79:VAL:HG12	1:B:87:ALA:HB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/104 (95%)	98 (99%)	1 (1%)	0	100	100
1	B	101/104 (97%)	99 (98%)	2 (2%)	0	100	100
1	C	102/104 (98%)	101 (99%)	1 (1%)	0	100	100
All	All	302/312 (97%)	298 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/85 (93%)	76 (96%)	3 (4%)	33	15
1	B	83/85 (98%)	81 (98%)	2 (2%)	49	32
1	C	83/85 (98%)	80 (96%)	3 (4%)	35	17
All	All	245/255 (96%)	237 (97%)	8 (3%)	38	20

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	VAL
1	A	84	ASN
1	A	98	ASN
1	B	84	ASN
1	B	86	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	14	SER
1	C	33	VAL
1	C	54	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	83	ASN
1	A	84	ASN
1	A	98	ASN
1	B	20	GLN
1	B	27	HIS
1	B	43	ASN
1	B	84	ASN
1	B	85	GLN
1	C	25	GLN
1	C	83	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	C	107	-	4,4,4	0.80	0	6,6,6	0.72	0
2	SO4	B	106	-	4,4,4	0.67	0	6,6,6	1.12	1 (16%)
2	SO4	A	105	-	4,4,4	0.46	0	6,6,6	0.64	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	106	SO4	O4-S-O3	2.16	118.29	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	105	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.