



wwPDB X-ray Structure Validation Summary Report ⓘ

May 18, 2020 – 11:29 pm BST

PDB ID : 3RQ1
Title : Crystal Structure of Aminotransferase Class I and II from *Veillonella parvula*
Authors : Kim, Y.; Hatzos-Skintges, C.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2011-04-27
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	 74% 24% ..
1	B	418	 2% 63% 33% ..
1	C	418	 9% 65% 30% ..
1	D	418	 9% 60% 29% 5% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	424	-	-	X	-
2	GOL	C	424	-	-	X	-
2	GOL	C	426	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminotransferase class I and II.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	414	3310	2090	572	630	11	7	0	9	0
1	B	414	3239	2051	554	617	10	7	0	1	0
1	C	413	3291	2078	565	631	10	7	0	8	0
1	D	392	3109	1974	527	593	9	6	0	5	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP D1BN29
A	-1	ASN	-	EXPRESSION TAG	UNP D1BN29
A	0	ALA	-	EXPRESSION TAG	UNP D1BN29
B	-2	SER	-	EXPRESSION TAG	UNP D1BN29
B	-1	ASN	-	EXPRESSION TAG	UNP D1BN29
B	0	ALA	-	EXPRESSION TAG	UNP D1BN29
C	-2	SER	-	EXPRESSION TAG	UNP D1BN29
C	-1	ASN	-	EXPRESSION TAG	UNP D1BN29
C	0	ALA	-	EXPRESSION TAG	UNP D1BN29
D	-2	SER	-	EXPRESSION TAG	UNP D1BN29
D	-1	ASN	-	EXPRESSION TAG	UNP D1BN29
D	0	ALA	-	EXPRESSION TAG	UNP D1BN29

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 12 6 6	0	1

- Molecule 3 is OXALOACETATE ION (three-letter code: OAA) (formula: $C_4H_3O_5$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 9 4 5	0	0
3	B	1	Total C O 9 4 5	0	0
3	C	1	Total C O 9 4 5	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Cl 1 1	0	0

- Molecule 5 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula: C₅H₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			10	5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	135	Total	O	0	0
			135	135		
6	B	82	Total	O	0	0
			82	82		
6	C	121	Total	O	0	0
			121	121		
6	D	104	Total	O	0	0
			104	104		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aminotransferase class I and II




- Molecule 1: Aminotransferase class I and II




- Molecule 1: Aminotransferase class I and II





• Molecule 1: Aminotransferase class I and II



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.99Å 81.06Å 106.39Å 102.40° 101.37° 102.95°	Depositor
Resolution (Å)	42.40 – 2.20 42.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.4 (42.40-2.20) 96.5 (42.40-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.190 , 0.263 0.183 , 0.253	Depositor DCC
R_{free} test set	4226 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	39.8	Xtrriage
Anisotropy	0.553	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13495	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, AKG, OAA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/3371	0.81	0/4548
1	B	0.63	0/3300	0.78	0/4454
1	C	0.68	0/3353	0.80	2/4527 (0.0%)
1	D	0.67	0/3169	0.82	5/4280 (0.1%)
All	All	0.67	0/13193	0.80	7/17809 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	95	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	D	95	ARG	NE-CZ-NH2	-5.94	117.33	120.30
1	D	387	ARG	NE-CZ-NH1	-5.67	117.47	120.30
1	C	188	ASN	C-N-CD	-5.48	108.55	120.60
1	D	119	TYR	N-CA-C	5.47	125.78	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	228	TYR	Mainchain
1	D	228	TYR	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3310	0	3224	95	0
1	B	3239	0	3163	159	0
1	C	3291	0	3194	155	0
1	D	3109	0	3016	153	0
2	A	18	0	24	10	0
2	B	6	0	8	1	0
2	C	18	0	24	17	0
2	D	24	0	32	5	0
3	A	9	0	2	0	0
3	B	9	0	2	0	0
3	C	9	0	2	1	0
4	C	1	0	0	0	0
5	C	10	0	4	3	0
6	A	135	0	0	8	0
6	B	82	0	0	15	0
6	C	121	0	0	10	0
6	D	104	0	0	5	0
All	All	13495	0	12695	531	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 531 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:LYS:HG2	1:A:349:MSE:HE1	1.25	1.14
1:C:189:PRO:HB2	1:C:359[B]:THR:HG21	1.37	1.07
1:C:29:ALA:HB2	1:C:37:VAL:HG11	1.30	1.06
1:C:267:ARG:NH2	2:C:426:GOL:H31	1.72	1.03
1:D:7:LYS:HA	1:D:10:LYS:HE2	1.43	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	421/418 (101%)	414 (98%)	7 (2%)	0	100	100
1	B	413/418 (99%)	404 (98%)	9 (2%)	0	100	100
1	C	419/418 (100%)	409 (98%)	10 (2%)	0	100	100
1	D	391/418 (94%)	377 (96%)	14 (4%)	0	100	100
All	All	1644/1672 (98%)	1604 (98%)	40 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/335 (104%)	330 (95%)	19 (5%)	22	26
1	B	341/335 (102%)	322 (94%)	19 (6%)	21	25
1	C	347/335 (104%)	314 (90%)	33 (10%)	8	8
1	D	327/335 (98%)	291 (89%)	36 (11%)	6	5
All	All	1364/1340 (102%)	1257 (92%)	107 (8%)	13	13

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	56	LYS
1	C	329	GLN
1	D	366	ASN
1	C	59	LYS
1	C	167	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	215	ASN
1	C	51	ASN
1	D	240	ASN
1	B	325	ASN
1	C	70	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 1 is monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	423	-	5,5,5	0.36	0	5,5,5	0.41	0
3	OAA	A	422	-	2,8,8	1.30	0	2,10,10	1.53	1 (50%)
2	GOL	D	421	-	5,5,5	0.32	0	5,5,5	0.38	0
5	AKG	C	423	-	3,9,9	0.92	0	4,11,11	3.91	1 (25%)
3	OAA	B	422	-	2,8,8	2.37	1 (50%)	2,10,10	1.95	1 (50%)
3	OAA	C	425	-	2,8,8	1.97	1 (50%)	2,10,10	0.71	0
2	GOL	B	416	-	5,5,5	0.31	0	5,5,5	0.47	0
2	GOL	C	426	-	5,5,5	0.42	0	5,5,5	0.54	0
2	GOL	A	421	-	5,5,5	0.35	0	5,5,5	0.64	0
2	GOL	D	416	-	5,5,5	0.37	0	5,5,5	0.37	0
2	GOL	C	421	-	5,5,5	0.38	0	5,5,5	0.40	0
2	GOL	D	423[A]	-	5,5,5	0.38	0	5,5,5	0.37	0
2	GOL	D	423[B]	-	5,5,5	0.31	0	5,5,5	0.37	0
2	GOL	A	424	-	5,5,5	0.30	0	5,5,5	0.49	0
2	GOL	C	424	-	5,5,5	0.46	0	5,5,5	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	423	-	-	3/4/4/4	-
3	OAA	A	422	-	-	2/2/8/8	-
2	GOL	D	421	-	-	2/4/4/4	-
5	AKG	C	423	-	-	0/3/9/9	-
3	OAA	B	422	-	-	2/2/8/8	-
3	OAA	C	425	-	-	2/2/8/8	-
2	GOL	B	416	-	-	2/4/4/4	-
2	GOL	C	426	-	-	0/4/4/4	-
2	GOL	A	421	-	-	3/4/4/4	-
2	GOL	D	416	-	-	0/4/4/4	-
2	GOL	C	421	-	-	4/4/4/4	-
2	GOL	D	423[A]	-	-	3/4/4/4	-
2	GOL	D	423[B]	-	-	2/4/4/4	-
2	GOL	A	424	-	-	2/4/4/4	-
2	GOL	C	424	-	-	1/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	422	OAA	C2-C3	-3.35	1.48	1.51
3	C	425	OAA	C2-C3	-2.75	1.49	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	423	AKG	C3-C4-C5	7.54	125.33	112.67
3	B	422	OAA	C1-C2-C3	-2.67	110.75	115.51
3	A	422	OAA	C1-C2-C3	2.07	119.20	115.51

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	422	OAA	C1-C2-C3-O3
3	A	422	OAA	C1-C2-C3-C4
3	C	425	OAA	C1-C2-C3-O3
3	C	425	OAA	C1-C2-C3-C4
2	B	416	GOL	O1-C1-C2-C3

There are no ring outliers.

10 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	423	GOL	2	0
5	C	423	AKG	3	0
3	C	425	OAA	1	0
2	B	416	GOL	1	0
2	C	426	GOL	4	0
2	A	421	GOL	1	0
2	D	423[A]	GOL	3	0
2	D	423[B]	GOL	2	0
2	A	424	GOL	7	0
2	C	424	GOL	13	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	407/418 (97%)	-0.34	1 (0%) 95 94	30, 44, 63, 85	0
1	B	407/418 (97%)	-0.05	7 (1%) 70 68	36, 58, 85, 110	0
1	C	406/418 (97%)	0.33	37 (9%) 9 8	29, 48, 101, 131	0
1	D	386/418 (92%)	0.32	37 (9%) 8 6	31, 53, 115, 166	0
All	All	1606/1672 (96%)	0.06	82 (5%) 28 26	29, 51, 95, 166	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	346	GLY	7.0
1	C	371	GLU	6.2
1	D	343	ALA	6.0
1	B	213	GLY	5.9
1	D	335	ALA	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	C	421	6/6	0.68	0.13	74,80,81,81	0
5	AKG	C	423	10/10	0.81	0.22	53,65,76,76	0
2	GOL	A	423	6/6	0.82	0.24	66,77,79,79	0
3	OAA	C	425	9/9	0.82	0.30	77,81,94,94	0
2	GOL	C	426	6/6	0.85	0.44	57,63,72,73	0
3	OAA	A	422	9/9	0.87	0.31	52,62,74,82	0
2	GOL	A	424	6/6	0.89	0.23	72,79,84,87	0
2	GOL	D	421	6/6	0.90	0.24	80,83,85,87	0
3	OAA	B	422	9/9	0.90	0.16	46,60,71,72	0
2	GOL	B	416	6/6	0.91	0.42	64,70,74,74	0
2	GOL	C	424	6/6	0.91	0.16	38,49,60,62	0
2	GOL	D	423[B]	6/6	0.92	0.46	61,62,64,65	6
2	GOL	D	423[A]	6/6	0.92	0.46	59,62,63,64	6
2	GOL	D	416	6/6	0.92	0.41	70,73,79,80	0
2	GOL	A	421	6/6	0.96	0.15	51,57,59,60	0
4	CL	C	422	1/1	0.97	0.07	75,75,75,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.