



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2024 – 04:21 pm BST

PDB ID : 8RSJ  
Title : Crystal structure of Methanobrevibacter oralis macrodomain in complex with ADPr in open conformation  
Authors : Ariza, A.  
Deposited on : 2024-01-24  
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	 94% 6%
1	B	262	 93% 6%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5164 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-acetyl-ADP-ribose deacetylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	2256	1430	374	431	21	0	33	0
1	B	261	2206	1395	369	424	18	0	27	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A166ACJ5
A	0	PRO	-	expression tag	UNP A0A166ACJ5
B	-1	GLY	-	expression tag	UNP A0A166ACJ5
B	0	PRO	-	expression tag	UNP A0A166ACJ5

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is [(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl [(2 {R},3 {S},4 {S})-2,3,4,5-tetrakis(oxidanyl)pentyl] hydrogen phosphate (three-letter code: JNT) (formula: C<sub>15</sub>H<sub>25</sub>N<sub>5</sub>O<sub>14</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	0	0
			36	15	5	14	2		
3	B	1	Total	C	N	O	P	0	0
			36	15	5	14	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	297	Total 297	O 297	0	0
5	B	325	Total 325	O 325	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: O-acetyl-ADP-ribose deacetylase

Chain A:  94% 6%



- Molecule 1: O-acetyl-ADP-ribose deacetylase

Chain B:  93% 6%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.99Å 58.10Å 76.27Å 90.00° 117.66° 90.00°	Depositor
Resolution (Å)	67.39 – 1.66 67.39 – 1.66	Depositor EDS
% Data completeness (in resolution range)	98.3 (67.39-1.66) 87.8 (67.39-1.66)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 1.66Å)	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
R, $R_{free}$	0.192 , 0.232 0.201 , 0.237	Depositor DCC
$R_{free}$ test set	3294 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.1	Xtriage
Anisotropy	0.709	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 33.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.044 for -h-l,k,h 0.044 for l,k,-h-l 0.114 for h,-k,-h-l 0.049 for -h-l,-k,l 0.046 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5164	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JNT, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/2387	0.80	0/3206
1	B	0.50	0/2316	0.78	0/3112
All	All	0.51	0/4703	0.79	0/6318

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	173[A]	ARG	Sidechain
1	A	173[B]	ARG	Sidechain
1	B	184	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	0	2347	17	0
1	B	2206	0	2273	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	36	0	0	0	0
3	B	36	0	0	0	0
4	B	6	0	8	0	0
5	A	297	0	0	3	1
5	B	325	0	0	2	1
All	All	5164	0	4628	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136[B]:MET:HE3	1:A:136[B]:MET:HA	1.36	1.04
1:B:217[B]:LYS:HA	1:B:217[B]:LYS:HE3	1.44	0.95
1:A:136[B]:MET:HA	1:A:136[B]:MET:CE	2.04	0.87
1:B:231[B]:ASN:OD1	5:B:401:HOH:O	1.95	0.82
1:B:217[B]:LYS:HE3	1:B:217[B]:LYS:CA	2.15	0.73
1:A:170[A]:ARG:HH11	1:A:170[A]:ARG:HB3	1.59	0.67
1:A:170[A]:ARG:HH11	1:A:170[A]:ARG:CB	2.13	0.60
1:B:237:LEU:HD13	1:B:240:ILE:HD11	1.84	0.59
1:A:170[A]:ARG:HB3	1:A:170[A]:ARG:NH1	2.20	0.55
1:A:170[B]:ARG:O	1:A:170[B]:ARG:HD3	2.06	0.55
1:B:237:LEU:HD13	1:B:240:ILE:CD1	2.38	0.54
1:A:136[B]:MET:CE	1:A:136[B]:MET:CA	2.83	0.52
1:A:53[B]:ARG:NH2	5:A:404:HOH:O	2.45	0.50
1:B:36:ARG:HG2	1:B:111[A]:MET:HE1	1.95	0.49
1:B:132:ASN:HD21	1:B:136:MET:HE2	1.78	0.48
1:A:25:PRO:HB2	1:A:30:GLU:CB	2.44	0.47
1:A:25:PRO:CB	1:A:30:GLU:HB3	2.45	0.47
1:B:132:ASN:HD21	1:B:136:MET:CE	2.27	0.47
1:A:118:ILE:HD13	5:A:623:HOH:O	2.15	0.46
1:A:25:PRO:HB2	1:A:30:GLU:HB3	1.97	0.46
1:B:120:HIS:HD2	5:B:542:HOH:O	1.98	0.46
1:A:36:ARG:HB3	1:A:111[A]:MET:HE1	2.00	0.44
1:A:111[A]:MET:HA	1:A:111[A]:MET:HE2	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LEU:HD22	1:B:126:GLN:HB3	1.99	0.44
1:A:120:HIS:HD2	5:A:570:HOH:O	2.01	0.43
1:B:177[B]:ASP:OD2	1:B:181[B]:LYS:HD2	2.19	0.42
1:B:111[A]:MET:CE	1:B:111[A]:MET:HA	2.50	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:447:HOH:O	5:B:501:HOH:O[2_546]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/262 (113%)	286 (97%)	9 (3%)	0	100	100
1	B	286/262 (109%)	284 (99%)	2 (1%)	0	100	100
All	All	581/524 (111%)	570 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/237 (115%)	268 (98%)	4 (2%)	60	41
1	B	262/237 (110%)	257 (98%)	5 (2%)	52	30
All	All	534/474 (113%)	525 (98%)	9 (2%)	60	35

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26[A]	LYS
1	A	26[B]	LYS
1	A	55	GLN
1	A	260	ASN
1	B	55	GLN
1	B	134	MET
1	B	141	ASP
1	B	172[A]	MET
1	B	172[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	HIS
1	A	120	HIS
1	A	154	ASN
1	B	120	HIS
1	B	132	ASN
1	B	151	ASN
1	B	154	ASN
1	B	212	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	JNT	A	302	-	33,38,38	2.67	9 (27%)	38,57,57	2.32	10 (26%)
4	GOL	B	303	-	5,5,5	0.11	0	5,5,5	0.42	0
3	JNT	B	302	-	33,38,38	2.25	9 (27%)	38,57,57	2.16	9 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JNT	A	302	-	-	7/28/48/48	0/3/3/3
4	GOL	B	303	-	-	0/4/4/4	-
3	JNT	B	302	-	-	6/28/48/48	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	JNT	O4'-C1'	7.62	1.51	1.41
3	A	302	JNT	C2'-C1'	-6.83	1.43	1.53
3	B	302	JNT	C2-N3	6.40	1.42	1.32
3	A	302	JNT	O2C-C2A	-6.11	1.30	1.43
3	B	302	JNT	O4C-C4A	6.08	1.56	1.43
3	A	302	JNT	C2-N3	4.21	1.38	1.32
3	A	302	JNT	O1C-C1A	-3.82	1.26	1.42
3	B	302	JNT	C2'-C1'	-3.50	1.48	1.53

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	JNT	O1C-C1A	-3.33	1.28	1.42
3	A	302	JNT	O4C-C4A	3.26	1.50	1.43
3	A	302	JNT	C5'-C4'	3.18	1.61	1.51
3	B	302	JNT	O4'-C1'	3.10	1.45	1.41
3	A	302	JNT	C4A-C3A	2.79	1.58	1.53
3	B	302	JNT	C5-C4	-2.69	1.33	1.40
3	B	302	JNT	C8-N7	-2.62	1.30	1.34
3	A	302	JNT	C3'-C4'	2.19	1.58	1.53
3	B	302	JNT	O5'-C5'	-2.18	1.36	1.44
3	B	302	JNT	O2C-C2A	-2.15	1.38	1.43

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	JNT	N3-C2-N1	-6.10	119.15	128.68
3	A	302	JNT	O1C-C1A-C2A	6.04	124.23	111.07
3	B	302	JNT	O4'-C1'-C2'	-5.64	98.68	106.93
3	A	302	JNT	C5A-C4A-C3A	-5.38	101.82	112.20
3	B	302	JNT	N3-C2-N1	-5.17	120.59	128.68
3	B	302	JNT	C5A-C4A-C3A	-4.32	103.87	112.20
3	A	302	JNT	C1'-N9-C4	-4.19	119.29	126.64
3	B	302	JNT	C1A-C2A-C3A	4.03	121.16	112.41
3	B	302	JNT	O1C-C1A-C2A	4.01	119.81	111.07
3	B	302	JNT	O2C-C2A-C1A	3.75	117.95	109.14
3	B	302	JNT	O4C-C4A-C5A	3.73	118.30	109.92
3	A	302	JNT	C1A-C2A-C3A	3.68	120.39	112.41
3	A	302	JNT	O4'-C1'-C2'	-3.67	101.56	106.93
3	B	302	JNT	O2C-C2A-C3A	3.62	117.90	109.10
3	A	302	JNT	O2C-C2A-C1A	3.19	116.64	109.14
3	A	302	JNT	O2C-C2A-C3A	2.77	115.83	109.10
3	A	302	JNT	O3C-C3A-C2A	-2.53	102.69	108.81
3	B	302	JNT	C1'-N9-C4	-2.47	122.29	126.64
3	A	302	JNT	O3C-C3A-C4A	2.09	113.87	108.81

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	JNT	C5'-O5'-PA-O3A
3	B	302	JNT	C5'-O5'-PA-O3A
3	B	302	JNT	O1C-C1A-C2A-O2C
3	A	302	JNT	C1A-C2A-C3A-O3C

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	302	JNT	C1A-C2A-C3A-O3C
3	B	302	JNT	C1A-C2A-C3A-C4A
3	A	302	JNT	C1A-C2A-C3A-C4A
3	A	302	JNT	O1C-C1A-C2A-O2C
3	A	302	JNT	PA-O3A-PB-O1B
3	B	302	JNT	PA-O3A-PB-O1B
3	A	302	JNT	PA-O3A-PB-O2B
3	A	302	JNT	C5'-O5'-PA-O1A
3	B	302	JNT	C5'-O5'-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/262 (100%)	-0.87	0 100 100	11, 24, 39, 58	33 (12%)
1	B	261/262 (99%)	-0.91	0 100 100	10, 23, 39, 59	27 (10%)
All	All	523/524 (99%)	-0.89	0 100 100	10, 24, 39, 59	60 (11%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	B	303	6/6	0.96	0.08	37,43,45,45	0
3	JNT	B	302	36/36	0.99	0.03	12,17,19,19	0
3	JNT	A	302	36/36	0.99	0.03	13,17,20,21	0
2	ZN	B	301	1/1	1.00	0.01	15,15,15,15	0
2	ZN	A	301	1/1	1.00	0.01	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.