



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 2, 2026 – 01:15 pm BST

PDB ID : 9RV7 / pdb\_00009rv7  
Title : Streptococcus pneumoniae StkP catalytic domain T167A/T169A double mutant in complex with AMP-PNP and Mg<sup>2+</sup>  
Authors : Hamidi, M.; Ravaud, S.; Grangeasse, C.  
Deposited on : 2025-07-07  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	<div> <div>7%</div> <div>68%</div> <div>14%</div> <div>•</div> <div>17%</div> </div>
1	B	309	<div> <div>5%</div> <div>78%</div> <div>13%</div> <div>•</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	B	303	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4888 atoms, of which 182 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase StkP.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	256	Total	C	H	N	O	S	0	0	0
			2115	1277	101	356	374	7			
1	B	286	Total	C	H	N	O	S	0	0	0
			2316	1413	81	390	423	9			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP Q8DNS0
A	-20	GLY	-	expression tag	UNP Q8DNS0
A	-19	GLY	-	expression tag	UNP Q8DNS0
A	-18	HIS	-	expression tag	UNP Q8DNS0
A	-17	HIS	-	expression tag	UNP Q8DNS0
A	-16	HIS	-	expression tag	UNP Q8DNS0
A	-15	HIS	-	expression tag	UNP Q8DNS0
A	-14	HIS	-	expression tag	UNP Q8DNS0
A	-13	HIS	-	expression tag	UNP Q8DNS0
A	-12	HIS	-	expression tag	UNP Q8DNS0
A	-11	GLY	-	expression tag	UNP Q8DNS0
A	-10	GLY	-	expression tag	UNP Q8DNS0
A	-9	GLY	-	expression tag	UNP Q8DNS0
A	-8	ARG	-	expression tag	UNP Q8DNS0
A	-7	GLU	-	expression tag	UNP Q8DNS0
A	-6	ASN	-	expression tag	UNP Q8DNS0
A	-5	LEU	-	expression tag	UNP Q8DNS0
A	-4	TYR	-	expression tag	UNP Q8DNS0
A	-3	PHE	-	expression tag	UNP Q8DNS0
A	-2	GLN	-	expression tag	UNP Q8DNS0
A	-1	GLY	-	expression tag	UNP Q8DNS0
A	0	HIS	-	expression tag	UNP Q8DNS0
A	167	ALA	THR	engineered mutation	UNP Q8DNS0
A	169	ALA	THR	engineered mutation	UNP Q8DNS0
B	-21	MET	-	initiating methionine	UNP Q8DNS0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	GLY	-	expression tag	UNP Q8DNS0
B	-19	GLY	-	expression tag	UNP Q8DNS0
B	-18	HIS	-	expression tag	UNP Q8DNS0
B	-17	HIS	-	expression tag	UNP Q8DNS0
B	-16	HIS	-	expression tag	UNP Q8DNS0
B	-15	HIS	-	expression tag	UNP Q8DNS0
B	-14	HIS	-	expression tag	UNP Q8DNS0
B	-13	HIS	-	expression tag	UNP Q8DNS0
B	-12	HIS	-	expression tag	UNP Q8DNS0
B	-11	GLY	-	expression tag	UNP Q8DNS0
B	-10	GLY	-	expression tag	UNP Q8DNS0
B	-9	GLY	-	expression tag	UNP Q8DNS0
B	-8	ARG	-	expression tag	UNP Q8DNS0
B	-7	GLU	-	expression tag	UNP Q8DNS0
B	-6	ASN	-	expression tag	UNP Q8DNS0
B	-5	LEU	-	expression tag	UNP Q8DNS0
B	-4	TYR	-	expression tag	UNP Q8DNS0
B	-3	PHE	-	expression tag	UNP Q8DNS0
B	-2	GLN	-	expression tag	UNP Q8DNS0
B	-1	GLY	-	expression tag	UNP Q8DNS0
B	0	HIS	-	expression tag	UNP Q8DNS0
B	167	ALA	THR	engineered mutation	UNP Q8DNS0
B	169	ALA	THR	engineered mutation	UNP Q8DNS0

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	120	Total	O	0	0
			120	120		
6	B	259	Total	O	0	0
			259	259		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Serine/threonine-protein kinase StkP



#### • Molecule 1: Serine/threonine-protein kinase StkP





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.28Å 51.73Å 81.14Å 90.00° 103.77° 90.00°	Depositor
Resolution (Å)	39.40 – 1.80 39.40 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.40-1.80) 99.9 (39.40-1.80)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.48 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.190 , 0.238 0.190 , 0.238	Depositor DCC
$R_{free}$ test set	2622 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.0	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4888	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MPD, GOL, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/2047	0.72	1/2764 (0.0%)
1	B	0.78	0/2273	0.87	1/3076 (0.0%)
All	All	0.67	0/4320	0.81	2/5840 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	254	LYS	CD-CE-NZ	5.73	130.25	111.90
1	A	269	LEU	CA-CB-CG	-5.69	96.39	116.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2014	101	2040	46	0
1	B	2235	81	2250	39	1
2	A	31	0	13	0	0
2	B	31	0	13	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	8	0	14	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	6	0	8	6	0
6	A	120	0	0	4	0
6	B	259	0	0	3	0
All	All	4706	182	4338	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:170:ASN:HB3	6:B:481:HOH:O	1.60	1.01
4:A:303:MPD:H12	6:B:475:HOH:O	1.66	0.94
1:B:188:LYS:HA	5:B:303:GOL:H32	1.51	0.89
1:B:188:LYS:N	1:B:188:LYS:HD2	1.92	0.84
1:B:189:ALA:HB2	5:B:303:GOL:H11	1.62	0.79
1:A:125:ARG:HD3	6:A:404:HOH:O	1.84	0.78
1:A:177:HIS:ND1	1:A:210:ILE:HD12	2.00	0.76
1:A:58:PHE:CZ	1:A:88:LEU:HD21	2.23	0.73
1:B:188:LYS:HD2	1:B:188:LYS:H	1.51	0.73
1:B:159:VAL:HG21	5:B:303:GOL:H12	1.69	0.73
1:A:79:ILE:CD1	1:A:88:LEU:HD22	2.19	0.71
1:A:57:ARG:HH21	1:A:60:ARG:HD3	1.55	0.71
1:A:57:ARG:HG2	1:A:60:ARG:HB2	1.77	0.67
1:A:62:ALA:HB1	1:A:79:ILE:HD12	1.77	0.66
1:A:57:ARG:NE	1:A:60:ARG:HH11	1.94	0.66
1:A:100:ARG:HH21	1:A:103:LYS:CB	2.10	0.65
1:A:125:ARG:O	1:A:129:THR:HG23	1.98	0.64
1:A:57:ARG:HG2	1:A:60:ARG:CB	2.27	0.63
1:A:57:ARG:HE	1:A:60:ARG:HB2	1.64	0.62
1:A:137:LEU:HD13	1:A:198:MET:HG3	1.82	0.62
1:B:106:TYR:CD1	1:B:107:PRO:HA	2.36	0.61
1:B:57:ARG:HD2	1:B:60:ARG:NH2	2.16	0.60
4:A:303:MPD:H11	6:B:637:HOH:O	2.02	0.59
4:A:303:MPD:C5	1:B:119:GLN:HE21	2.15	0.59
1:A:122:LEU:HD13	1:B:122:LEU:HD13	1.84	0.59
4:A:303:MPD:H53	1:B:119:GLN:HE21	1.66	0.59
1:B:189:ALA:CB	5:B:303:GOL:H11	2.32	0.58
1:A:57:ARG:NH2	1:A:60:ARG:HD3	2.20	0.55
1:A:177:HIS:HD1	1:A:210:ILE:HD12	1.70	0.54
1:B:76:ILE:HD13	1:B:90:MET:HG2	1.88	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:188:LYS:HA	5:B:303:GOL:C3	2.33	0.53
1:A:269:LEU:O	1:A:269:LEU:HG	1.91	0.52
1:A:57:ARG:HE	1:A:60:ARG:HD3	1.75	0.51
1:A:98:LEU:O	1:A:102:ILE:HG13	2.10	0.51
1:A:284:ILE:O	1:A:285:PHE:C	2.53	0.51
1:A:276:ASN:CA	6:A:403:HOH:O	2.59	0.51
1:A:7:ILE:HD12	1:A:32:LEU:HD21	1.92	0.50
1:B:11:ARG:HG3	1:B:33:ILE:HG12	1.93	0.50
1:A:121:LEU:HD12	1:A:269:LEU:HD22	1.94	0.48
1:B:113:ALA:O	1:B:117:MET:HG3	2.12	0.48
1:A:100:ARG:HH21	1:A:103:LYS:CG	2.26	0.48
1:A:65:MET:O	1:A:68:LEU:HB2	2.14	0.48
1:A:259:TYR:CE2	1:A:265:MET:HG3	2.49	0.48
1:B:11:ARG:HG3	1:B:33:ILE:CG1	2.43	0.48
1:A:276:ASN:HA	6:A:403:HOH:O	2.14	0.46
1:A:100:ARG:NE	1:A:100:ARG:HA	2.31	0.46
1:B:74:VAL:HG21	1:B:153:THR:HB	1.98	0.46
1:B:37:GLU:HA	1:B:37:GLU:OE1	2.15	0.46
1:A:176:VAL:N	6:A:407:HOH:O	2.49	0.46
1:B:188:LYS:CA	5:B:303:GOL:H32	2.34	0.46
1:B:44:LEU:HD22	1:B:161:PHE:CE2	2.51	0.46
1:A:106:TYR:CD1	1:A:107:PRO:HA	2.51	0.45
1:B:144:LEU:HD23	1:B:150:ALA:HA	1.98	0.45
1:A:58:PHE:CE2	1:A:86:GLN:HB3	2.52	0.45
1:A:59:GLN:O	1:A:63:ARG:HG2	2.17	0.45
1:B:70:HIS:ND1	1:B:71:PRO:HD2	2.32	0.45
1:A:5:GLY:O	1:A:6:LYS:HD3	2.18	0.44
1:A:100:ARG:HH21	1:A:103:LYS:HG2	1.82	0.44
1:A:57:ARG:HH21	1:A:60:ARG:CD	2.26	0.44
1:A:79:ILE:HD12	1:A:88:LEU:HD22	1.95	0.43
1:B:182:GLU:HG2	1:B:183:GLN:N	2.33	0.43
1:A:100:ARG:NH2	1:A:103:LYS:HG2	2.34	0.43
1:B:74:VAL:HG13	1:B:91:GLU:HG2	2.00	0.43
1:B:76:ILE:HD12	1:B:88:LEU:HD12	2.00	0.43
1:B:3:GLN:HG3	1:B:6:LYS:HG2	2.01	0.43
1:A:71:PRO:HD3	1:B:125:ARG:CZ	2.49	0.42
1:B:74:VAL:HG21	1:B:143:LEU:HD12	2.01	0.42
1:B:70:HIS:CE1	1:B:71:PRO:HD2	2.55	0.42
1:A:209:HIS:HE2	1:A:235:GLU:CD	2.27	0.42
1:A:57:ARG:NE	1:A:60:ARG:HB2	2.31	0.42
1:B:172:MET:O	1:B:173:LEU:HB2	2.20	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:PHE:HZ	1:A:88:LEU:CD2	2.33	0.42
1:B:135:ARG:HG2	1:B:189:ALA:HB3	2.01	0.42
1:A:125:ARG:NE	1:A:263:SER:OG	2.49	0.41
1:B:3:GLN:HG3	1:B:6:LYS:CG	2.50	0.41
1:B:70:HIS:CG	1:B:71:PRO:HD2	2.55	0.41
1:A:134:HIS:O	1:A:135:ARG:HB2	2.21	0.41
1:B:58:PHE:CE2	1:B:86:GLN:HB3	2.56	0.41
1:A:57:ARG:HG2	1:A:60:ARG:HB3	2.02	0.41
1:A:130:ARG:O	1:A:132:ILE:HG13	2.20	0.41
1:B:136:ASP:OD2	2:B:301:ANP:O2G	2.38	0.41
1:A:124:MET:HG3	1:A:198:MET:SD	2.61	0.40
1:B:137:LEU:HD23	1:B:137:LEU:HA	1.94	0.40
1:B:171:SER:C	1:B:172:MET:SD	3.04	0.40
1:B:269:LEU:HD13	1:B:269:LEU:C	2.47	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:GLY:O	1:B:172:MET:O[2_658]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/309 (81%)	239 (96%)	11 (4%)	0	100	100
1	B	284/309 (92%)	271 (95%)	11 (4%)	2 (1%)	18	8
All	All	534/618 (86%)	510 (96%)	22 (4%)	2 (0%)	30	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	167	ALA
1	B	172	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/256 (84%)	211 (99%)	3 (1%)	59	52
1	B	238/256 (93%)	234 (98%)	4 (2%)	53	45
All	All	452/512 (88%)	445 (98%)	7 (2%)	57	49

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LEU
1	A	55	VAL
1	A	58	PHE
1	B	44	LEU
1	B	88	LEU
1	B	159	VAL
1	B	269	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	HIS
1	A	183	GLN
1	A	241	GLN
1	B	49	GLN
1	B	119	GLN
1	B	168	GLN
1	B	170	ASN
1	B	223	GLN
1	B	227	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ANP	B	301	3	32,33,33	2.05	5 (15%)	44,52,52	1.67	5 (11%)
2	ANP	A	301	3	32,33,33	2.54	5 (15%)	44,52,52	1.18	4 (9%)
4	MPD	A	303	-	7,7,7	0.50	0	9,10,10	1.24	1 (11%)
5	GOL	B	303	-	5,5,5	0.97	0	5,5,5	1.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	B	301	3	-	4/18/38/38	0/3/3/3
2	ANP	A	301	3	-	2/18/38/38	0/3/3/3
4	MPD	A	303	-	-	1/5/5/5	-
5	GOL	B	303	-	-	2/4/4/4	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	ANP	PB-O3A	11.28	1.73	1.59
2	B	301	ANP	PB-O3A	6.92	1.67	1.59
2	A	301	ANP	PG-N3B	5.97	1.79	1.63
2	B	301	ANP	PG-N3B	5.39	1.77	1.63
2	B	301	ANP	PG-O1G	4.29	1.53	1.46
2	A	301	ANP	PG-O1G	4.22	1.52	1.46
2	B	301	ANP	PB-O1B	3.65	1.51	1.46
2	A	301	ANP	PB-O1B	2.47	1.50	1.46
2	A	301	ANP	PB-O2B	-2.09	1.51	1.56
2	B	301	ANP	PG-O2G	-2.06	1.51	1.56

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	ANP	O2B-PB-O1B	6.90	124.39	109.92
2	B	301	ANP	C2'-C3'-C4'	-4.92	93.08	102.64
2	A	301	ANP	O2B-PB-O1B	3.81	117.91	109.92
2	B	301	ANP	O1G-PG-N3B	-3.48	106.65	111.77
2	A	301	ANP	O1G-PG-N3B	-3.07	107.25	111.77
2	A	301	ANP	C2'-C3'-C4'	2.40	107.30	102.64
2	B	301	ANP	O1B-PB-N3B	2.17	114.96	111.77
2	A	301	ANP	O2G-PG-O3G	2.13	113.31	107.64
2	B	301	ANP	C3'-C2'-C1'	2.10	105.42	101.43
4	A	303	MPD	C1-C2-C3	-2.09	100.22	109.96

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	ANP	PB-N3B-PG-O1G
4	A	303	MPD	C2-C3-C4-O4
5	B	303	GOL	C1-C2-C3-O3
5	B	303	GOL	O2-C2-C3-O3
2	B	301	ANP	PB-O3A-PA-O1A
2	B	301	ANP	PG-N3B-PB-O3A
2	B	301	ANP	PB-O3A-PA-O2A
2	A	301	ANP	PB-O3A-PA-O1A
2	B	301	ANP	C5'-O5'-PA-O1A


There are no ring outliers.



3 monomers are involved in 11 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	ANP	1	0
4	A	303	MPD	4	0
5	B	303	GOL	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	256/309 (82%)	0.67	21 (8%) 17 15	26, 40, 65, 95	0
1	B	286/309 (92%)	0.14	14 (4%) 35 33	18, 28, 56, 84	0
All	All	542/618 (87%)	0.39	35 (6%) 25 23	18, 34, 63, 95	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	169	ALA	6.6
1	B	170	ASN	6.2
1	A	55	VAL	6.2
1	B	173	LEU	5.6
1	B	166	LEU	4.8
1	B	167	ALA	4.3
1	A	56	ALA	4.2
1	B	171	SER	4.0
1	B	2	ILE	3.9
1	A	285	PHE	3.9
1	B	172	MET	3.8
1	A	156	GLY	3.5
1	B	165	SER	3.4
1	B	162	ALA	3.3
1	A	176	VAL	3.0
1	B	164	THR	2.8
1	A	275	TYR	2.8
1	A	284	ILE	2.8
1	A	157	ILE	2.8
1	A	57	ARG	2.7
1	A	100	ARG	2.7
1	B	163	GLU	2.7
1	A	213	ASP	2.6
1	A	44	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	22	GLY	2.6
1	A	155	PHE	2.5
1	A	58	PHE	2.4
1	A	256	THR	2.4
1	B	159	VAL	2.3
1	B	168	GLN	2.3
1	A	60	ARG	2.2
1	A	88	LEU	2.1
1	A	106	TYR	2.1
1	A	132	ILE	2.0
1	A	4	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	B	303	6/6	0.75	0.13	38,40,41,44	0
3	MG	A	302	1/1	0.79	0.12	51,51,51,51	0
4	MPD	A	303	8/8	0.89	0.11	23,33,38,41	0
2	ANP	A	301	31/31	0.91	0.09	36,50,57,61	0
3	MG	B	302	1/1	0.94	0.05	25,25,25,25	0
2	ANP	B	301	31/31	0.97	0.05	21,25,29,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around GOL B 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MG A 302:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around MPD A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ANP A 301:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.