



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 02:50 AM EDT

PDB ID : 7RX8  
Title : Structure of METTL3-METTL14(R298H) mutant methyltransferase complex  
Authors : Wang, P.; Nam, Y.  
Deposited on : 2021-08-21  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	
2	B	349	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase 70 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	201	1674	1074	298	290	12	0	7	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	MET	-	initiating methionine	UNP Q86U44

- Molecule 2 is a protein called N6-adenosine-methyltransferase non-catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	284	2331	1486	402	431	12	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	108	GLY	-	expression tag	UNP Q9HCE5
B	109	SER	-	expression tag	UNP Q9HCE5
B	110	GLY	-	expression tag	UNP Q9HCE5
B	298	HIS	ARG	engineered mutation	UNP Q9HCE5


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	111	Total	O	0	0
			111	111		
3	B	271	Total	O	0	0
			271	271		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N6-adenosine-methyltransferase 70 kDa subunit



- Molecule 2: N6-adenosine-methyltransferase non-catalytic subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.48Å 101.48Å 117.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.57 – 1.85 46.57 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.8 (46.57-1.85) 97.8 (46.57-1.85)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 1.86Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.171 , 0.202 0.170 , 0.201	Depositor DCC
$R_{free}$ test set	2514 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtrriage
Anisotropy	0.024	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4387	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/1740	0.64	0/2363
2	B	0.47	0/2396	0.65	0/3247
All	All	0.44	0/4136	0.65	0/5610

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1674	0	1673	19	0
2	B	2331	0	2293	18	0
3	A	111	0	0	1	0
3	B	271	0	0	4	0
All	All	4387	0	3966	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:297:LYS:HB2	2:B:300:THR:HG22	1.68	0.73

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:LYS:HD3	2:B:313:LEU:HD21	1.81	0.62
1:A:437[B]:MET:SD	3:B:532:HOH:O	2.56	0.62
1:A:373:TRP:HB2	1:A:551:LEU:HD13	1.83	0.60
2:B:162:LYS:NZ	3:B:501:HOH:O	2.30	0.59
1:A:471[A]:ARG:NH2	2:B:399:SER:HA	2.19	0.58
1:A:437[B]:MET:SD	3:B:514:HOH:O	2.57	0.57
1:A:389:PHE:O	1:A:423:GLN:HG2	2.06	0.55
2:B:390:GLU:OE1	2:B:394:ARG:NH1	2.40	0.55
1:A:564:ARG:NH2	3:A:603:HOH:O	2.41	0.54
2:B:272:THR:H	2:B:282:GLN:NE2	2.04	0.54
1:A:555:HIS:CE1	1:A:557:LEU:HD21	2.45	0.52
1:A:444[A]:LEU:HG	1:A:449:TYR:HB2	1.92	0.51
2:B:200:ARG:O	2:B:275:LEU:HD22	2.13	0.49
2:B:311:ILE:HD12	2:B:313:LEU:HB2	1.96	0.48
2:B:297:LYS:CB	2:B:300:THR:HG22	2.41	0.47
2:B:263:LYS:HB3	2:B:318:GLU:HG2	1.98	0.46
1:A:434:GLY:O	1:A:437[A]:MET:HG2	2.16	0.45
1:A:404:LEU:HD13	1:A:406:TYR:CZ	2.52	0.45
1:A:466:ILE:HG21	2:B:311:ILE:HD13	1.98	0.45
2:B:324:ILE:H	2:B:324:ILE:HG13	1.67	0.44
1:A:465:ARG:NE	2:B:136:ASP:OD2	2.45	0.44
2:B:209:LYS:HD3	2:B:211:TRP:NE1	2.33	0.44
1:A:496:GLN:HG2	2:B:280:VAL:HA	2.01	0.43
1:A:541:GLN:HB2	1:A:544:TRP:CD1	2.54	0.42
1:A:560:ASP:O	1:A:564:ARG:HG3	2.20	0.41
1:A:369:PHE:CZ	1:A:558:ASP:HA	2.55	0.41
1:A:376:CYS:SG	1:A:381:LEU:HD22	2.61	0.41
1:A:378:ILE:HD13	1:A:534:PHE:CD1	2.57	0.40
2:B:118:ASP:OD1	2:B:120:CYS:HB2	2.21	0.40
2:B:200:ARG:NH1	2:B:201:GLU:OE2	2.54	0.40
2:B:282:GLN:NE2	3:B:510:HOH:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/225 (92%)	202 (98%)	4 (2%)	0	100	100
2	B	284/349 (81%)	277 (98%)	7 (2%)	0	100	100
All	All	490/574 (85%)	479 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/198 (93%)	185 (100%)	0	100	100
2	B	256/295 (87%)	255 (100%)	1 (0%)	91	89
All	All	441/493 (90%)	440 (100%)	1 (0%)	93	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	390	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	423	GLN
1	A	555	HIS
2	B	282	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	201/225 (89%)	0.34	19 (9%) 8 7	9, 32, 59, 74	0
2	B	284/349 (81%)	0.04	15 (5%) 26 25	8, 20, 56, 82	0
All	All	485/574 (84%)	0.16	34 (7%) 16 15	8, 24, 58, 82	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	369	PHE	6.6
2	B	299	SER	4.9
1	A	557	LEU	4.6
2	B	300	THR	4.4
2	B	116	HIS	4.3
2	B	202	THR	4.3
1	A	561	VAL	4.2
1	A	565	PHE	4.0
2	B	275	LEU	3.8
2	B	207	ASN	3.7
1	A	559	PRO	3.6
1	A	370	PRO	3.4
2	B	298	HIS	3.3
2	B	204	ILE	3.2
1	A	406	TYR	3.1
1	A	492	GLN	2.9
1	A	554	ILE	2.9
2	B	117	ASN	2.8
2	B	203	GLY	2.8
1	A	567	GLN	2.8
1	A	403	GLU	2.8
1	A	563	ALA	2.6
2	B	321	ILE	2.6
1	A	560	ASP	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	494	PHE	2.5
1	A	556	LEU	2.5
1	A	527	GLY	2.4
1	A	558	ASP	2.2
2	B	139[A]	LEU	2.1
2	B	199	TYR	2.1
1	A	490	ASN	2.1
2	B	200	ARG	2.1
2	B	297	LYS	2.1
1	A	445[A]	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.