



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 4, 2024 – 12:43 AM EDT

PDB ID : 8SC9
Title : Structure of PPARG in complex with MTX-531
Authors : Whitehead, C.E.; Leopold, J.
Deposited on : 2023-04-05
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	 10% 88% 5% 6%
1	B	278	 13% 89% 5% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1005	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8985 atoms, of which 4384 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	261	4288	1359	2180	345	393	11	2180	5	0
1	B	262	4338	1378	2204	350	395	11	2204	6	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	200	GLY	-	expression tag	UNP P37231
A	201	SER	-	expression tag	UNP P37231
A	202	HIS	-	expression tag	UNP P37231
A	203	MET	-	expression tag	UNP P37231
B	200	GLY	-	expression tag	UNP P37231
B	201	SER	-	expression tag	UNP P37231
B	202	HIS	-	expression tag	UNP P37231
B	203	MET	-	expression tag	UNP P37231

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: $C_7H_{17}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	11	7	1	3	0	0

- Molecule 4 is N-[(5P)-2-chloro-5-(4-[(1R)-1-phenylethyl]amino)quinazolin-6-yl]pyridin-3-yl]methanesulfonamide (three-letter code: D0D) (formula: C₂₂H₂₀ClN₅O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Cl	N	O			S
4	B	1	40	30	1	6	2	1	0	1


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	126	Total 128	O 128	0	2
5	B	129	Total 130	O 130	0	3


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.20Å 61.38Å 118.74Å 90.00° 102.52° 90.00°	Depositor
Resolution (Å)	33.22 – 1.85 33.22 – 1.86	Depositor EDS
% Data completeness (in resolution range)	63.4 (33.22-1.85) 63.4 (33.22-1.86)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.85Å)	Xtrriage
Refinement program	BUSTER 2.11.8 (8-JUN-2022)	Depositor
R, R_{free}	0.212 , 0.241 0.207 , 0.232	Depositor DCC
R_{free} test set	1762 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	42.5	Xtrriage
Anisotropy	0.078	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 60.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8985	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: D0D, SO4, TAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2160	0.38	0/2910
1	B	0.26	0/2186	0.38	0/2942
All	All	0.26	0/4346	0.38	0/5852

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	2180	2171	12	0
1	B	2134	2204	2192	11	1
2	A	25	0	0	3	0
2	B	25	0	0	0	0
3	A	11	0	17	2	0
4	B	40	0	0	4	0
5	A	128	0	0	0	0
5	B	130	0	0	1	0
All	All	4601	4384	4380	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:323:HIS:NE2	1:B:476:LEU:HD12	2.07	0.70
1:B:323:HIS:CD2	4:B:1006[B]:D0D:C18	2.83	0.62
1:A:205:ASN:OD1	1:A:206:PRO:HD2	2.01	0.60
1:A:244:LYS:HD3	3:A:1006:TAM:H61	1.84	0.58
1:B:453:LEU:O	1:B:457:LYS:HG2	2.05	0.56
1:A:402[B]:ASN:OD1	2:A:1003:SO4:O2	2.27	0.51
1:B:453:LEU:O	1:B:456:ILE:HG12	2.11	0.51
1:B:323:HIS:HD2	4:B:1006[B]:D0D:C18	2.23	0.50
1:A:290:VAL:O	1:A:294:GLN:HG3	2.13	0.49
1:B:234[A]:ARG:HD3	5:B:1101[A]:HOH:O	2.11	0.49
1:B:326:ILE:HG21	4:B:1006[B]:D0D:C16	2.45	0.47
1:A:343:GLU:O	1:A:343:GLU:HG3	2.15	0.47
1:A:290:VAL:HG21	1:A:466:HIS:CG	2.51	0.44
1:B:449:HIS:NE2	4:B:1006[B]:D0D:C19	2.81	0.44
1:A:443:ARG:HG3	1:B:440:THR:CG2	2.49	0.43
1:A:323:HIS:CE1	1:A:472:ILE:HG21	2.53	0.42
1:A:216:LYS:NZ	2:A:1002:SO4:O4	2.43	0.42
1:A:244:LYS:HA	3:A:1006:TAM:H61	2.02	0.41
1:A:381:ASP:HB2	2:A:1004:SO4:O4	2.19	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:ASN:HD22	1:B:337:ASP:OD1[2_656]	1.44	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/278 (94%)	262 (100%)	0	0	100	100
1	B	262/278 (94%)	260 (99%)	2 (1%)	0	100	100
All	All	524/556 (94%)	522 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/250 (96%)	238 (100%)	1 (0%)	91	89
1	B	241/250 (96%)	240 (100%)	1 (0%)	91	89
All	All	480/500 (96%)	478 (100%)	2 (0%)	91	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	444	GLN
1	B	210	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	430	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TAM	A	1006	-	7,10,10	0.55	0	9,12,12	0.84	1 (11%)
2	SO4	A	1002	-	4,4,4	0.18	0	6,6,6	0.12	0
2	SO4	B	1003	-	4,4,4	0.16	0	6,6,6	0.14	0
2	SO4	B	1005	-	4,4,4	0.22	0	6,6,6	0.11	0
4	D0D	B	1006[A]	-	33,34,34	1.33	4 (12%)	46,49,49	1.17	4 (8%)
2	SO4	A	1004	-	4,4,4	0.19	0	6,6,6	0.17	0
4	D0D	B	1006[B]	-	33,34,34	1.30	4 (12%)	46,49,49	1.19	5 (10%)
2	SO4	A	1005	-	4,4,4	0.24	0	6,6,6	0.15	0
2	SO4	A	1001	-	4,4,4	0.17	0	6,6,6	0.08	0
2	SO4	A	1003	-	4,4,4	0.23	0	6,6,6	0.13	0
2	SO4	B	1001	-	4,4,4	0.19	0	6,6,6	0.20	0
2	SO4	B	1002	-	4,4,4	0.23	0	6,6,6	0.20	0
2	SO4	B	1004	-	4,4,4	0.17	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TAM	A	1006	-	-	3/12/12/12	-
4	D0D	B	1006[B]	-	-	2/17/17/17	0/4/4/4
4	D0D	B	1006[A]	-	-	2/17/17/17	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1006[A]	D0D	S-N1	3.42	1.67	1.63
4	B	1006[B]	D0D	S-N1	3.42	1.67	1.63
4	B	1006[A]	D0D	C11-S	-2.97	1.68	1.75
4	B	1006[B]	D0D	C11-S	-2.97	1.68	1.75
4	B	1006[A]	D0D	C13-N3	2.54	1.37	1.34
4	B	1006[B]	D0D	C13-N3	2.54	1.37	1.34
4	B	1006[A]	D0D	C12-N3	2.48	1.38	1.33
4	B	1006[B]	D0D	C12-N3	2.48	1.38	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1006[A]	D0D	C11-S-N1	-3.87	102.22	106.63
4	B	1006[B]	D0D	C11-S-N1	-3.87	102.22	106.63
4	B	1006[A]	D0D	C7-C6-C4	-2.96	116.54	121.69
4	B	1006[B]	D0D	C7-C6-C4	-2.96	116.54	121.69
4	B	1006[A]	D0D	C3-C4-C6	-2.56	116.93	121.36
4	B	1006[B]	D0D	C3-C4-C6	-2.56	116.93	121.36
4	B	1006[B]	D0D	C21-C14-N4	2.33	113.37	108.95
3	A	1006	TAM	C3-C-C2	-2.05	106.89	110.50
4	B	1006[A]	D0D	C3-C4-C5	2.04	121.29	118.09
4	B	1006[B]	D0D	C3-C4-C5	2.04	121.29	118.09

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1006	TAM	C2-C-C1-C4
3	A	1006	TAM	C3-C-C1-C4
3	A	1006	TAM	N-C-C1-C4
4	B	1006[A]	D0D	C9-N1-S-O1
4	B	1006[A]	D0D	C9-N1-S-C11
4	B	1006[B]	D0D	C9-N1-S-O1
4	B	1006[B]	D0D	C9-N1-S-C11

There are no ring outliers.

5 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1006	TAM	2	0
2	A	1002	SO4	1	0
2	A	1004	SO4	1	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1006[B]	D0D	4	0
2	A	1003	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/278 (93%)	0.72	29 (11%) 5 5	34, 54, 87, 112	0
1	B	262/278 (94%)	0.75	37 (14%) 2 3	30, 51, 101, 149	0
All	All	523/556 (94%)	0.73	66 (12%) 3 3	30, 53, 94, 149	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	459	THR	12.0
1	B	456	ILE	8.3
1	B	458	LYS	7.4
1	B	242	THR	7.3
1	B	455	VAL	6.6
1	A	206	PRO	6.5
1	A	204	LEU	6.1
1	A	475	ASP	6.0
1	A	262	ILE	5.8
1	B	461	THR	5.8
1	A	261	LYS	5.7
1	B	275	LYS	5.6
1	B	452	LEU	5.5
1	B	269	PRO	5.3
1	B	462	ASP	5.2
1	B	240	LYS	5.0
1	A	241	THR	4.7
1	A	260	ASP	4.5
1	B	363	PHE	4.5
1	B	454	GLN	4.5
1	A	474	LYS	4.1
1	B	243	ASP	4.0
1	B	460	GLU	3.9
1	B	239	GLY	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	359	PRO	3.6
1	B	207	GLU	3.5
1	B	473	TYR	3.5
1	B	241	THR	3.4
1	A	207	GLU	3.1
1	B	267	ILE	3.1
1	A	362	ASP	3.0
1	A	256	MET	3.0
1	B	360	PHE	3.0
1	B	474	LYS	3.0
1	B	263	LYS	2.9
1	B	476	LEU	2.9
1	A	205	ASN	2.8
1	A	391	ILE	2.8
1	B	451	GLN	2.8
1	A	325	ILE	2.7
1	B	285[A]	CYS	2.7
1	B	470	GLN	2.7
1	B	466	HIS	2.6
1	B	427	GLU	2.6
1	A	275	LYS	2.6
1	A	388	ILE	2.5
1	B	268	THR	2.5
1	B	357	ARG	2.5
1	B	247	PHE	2.4
1	B	339	VAL	2.4
1	A	442	LEU	2.4
1	A	328	THR	2.4
1	B	341	ILE	2.4
1	B	340	LEU	2.3
1	A	327	TYR	2.3
1	A	387	PHE	2.2
1	A	467	PRO	2.2
1	A	244	LYS	2.2
1	A	426	PRO	2.2
1	A	476	LEU	2.2
1	A	390	VAL	2.1
1	A	444	GLN	2.1
1	B	358	LYS	2.1
1	A	427	GLU	2.1
1	B	327	TYR	2.1
1	A	296	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	1005	5/5	0.75	0.47	127,127,127,127	0
4	D0D	B	1006[A]	31/31	0.84	0.20	68,70,71,72	9
4	D0D	B	1006[B]	31/31	0.84	0.20	68,71,71,72	9
2	SO4	B	1003	5/5	0.85	0.29	110,110,110,110	0
2	SO4	A	1003	5/5	0.88	0.29	127,127,127,128	0
3	TAM	A	1006	11/11	0.88	0.17	72,74,74,75	0
2	SO4	B	1005	5/5	0.91	0.27	119,119,119,119	0
2	SO4	B	1002	5/5	0.92	0.09	103,104,104,104	0
2	SO4	A	1002	5/5	0.92	0.27	102,103,103,103	0
2	SO4	B	1004	5/5	0.93	0.24	125,125,125,125	0
2	SO4	A	1001	5/5	0.95	0.20	119,119,119,119	0
2	SO4	A	1004	5/5	0.95	0.29	89,89,89,89	0
2	SO4	B	1001	5/5	0.96	0.16	72,73,73,74	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around D0D B 1006 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around D0D B 1006 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.