



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 17, 2024 – 03:37 PM EST

PDB ID : 3SDS
Title : Crystal structure of a mitochondrial ornithine carbamoyltransferase from *Coccidioides immitis*
Authors : Edwards, T.E.; Abendroth, J.; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2011-06-09
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	 84% 9% 11%
1	B	353	 80% 9% 11%
1	C	353	 81% 8% 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine carbamoyltransferase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2323	1474	398	440	11	0	0	0
1	B	315	2313	1466	398	438	11	0	0	0
1	C	315	2292	1453	391	437	11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P0CL21
A	-2	PRO	-	expression tag	UNP P0CL21
A	-1	GLY	-	expression tag	UNP P0CL21
A	0	SER	-	expression tag	UNP P0CL21
B	-3	GLY	-	expression tag	UNP P0CL21
B	-2	PRO	-	expression tag	UNP P0CL21
B	-1	GLY	-	expression tag	UNP P0CL21
B	0	SER	-	expression tag	UNP P0CL21
C	-3	GLY	-	expression tag	UNP P0CL21
C	-2	PRO	-	expression tag	UNP P0CL21
C	-1	GLY	-	expression tag	UNP P0CL21
C	0	SER	-	expression tag	UNP P0CL21

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cl	0	0
			2	2		
2	B	2	Total	Cl	0	0
			2	2		
2	C	2	Total	Cl	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total 28	O 28	0	0
3	B	14	Total 14	O 14	0	0
3	C	22	Total 22	O 22	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.36Å 150.36Å 92.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.80 19.86 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-2.80) 99.6 (19.86-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.79Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.193 , 0.239 0.192 , 0.237	Depositor DCC
R_{free} test set	1504 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	57.1	Xtrriage
Anisotropy	0.045	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6998	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/2368	0.63	0/3223
1	B	0.56	0/2357	0.62	0/3211
1	C	0.56	0/2337	0.61	0/3190
All	All	0.58	0/7062	0.62	0/9624

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2323	0	2265	5	0
1	B	2313	0	2235	15	0
1	C	2292	0	2187	12	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	28	0	0	0	0
3	B	14	0	0	0	0
3	C	22	0	0	0	0
All	All	6998	0	6687	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:145:ALA:O	1:B:146:LEU:HB2	1.94	0.67
1:B:250:VAL:HG22	1:C:176:HIS:HB2	1.77	0.66
1:C:298:TRP:CZ2	1:C:320:ARG:HD3	2.36	0.60
1:C:303:CYS:O	1:C:304:LEU:HB2	2.05	0.56
1:B:28:LEU:O	1:B:206:LYS:NZ	2.29	0.56
1:C:105:GLU:OE2	1:C:110:THR:OG1	2.20	0.56
1:B:307:HIS:HB3	1:B:308:PRO:HD2	1.90	0.54
1:C:81:GLU:HG3	1:C:91:PRO:HG2	1.90	0.53
1:A:145:ALA:O	1:A:146:LEU:HB2	2.09	0.52
1:A:307:HIS:HB3	1:A:308:PRO:HD2	1.93	0.50
1:B:304:LEU:HB3	1:B:305:PRO:HA	1.95	0.49
1:C:44:LYS:HD2	1:C:332:TRP:CD1	2.48	0.48
1:B:250:VAL:CG2	1:C:176:HIS:HB2	2.42	0.47
1:A:69:PHE:HE1	1:A:77:ARG:HG3	1.80	0.47
1:B:246:THR:O	1:B:246:THR:HG22	2.15	0.46
1:A:74:THR:O	1:A:78:VAL:HG23	2.16	0.46
1:B:95:GLY:O	1:B:98:ASP:N	2.47	0.46
1:C:322:LEU:O	1:C:325:PRO:HD2	2.16	0.46
1:C:283:LYS:HD3	1:C:309:GLU:O	2.15	0.45
1:C:61:SER:HA	1:C:88:GLY:O	2.15	0.45
1:A:44:LYS:HD2	1:A:332:TRP:CD1	2.52	0.44
1:B:176:HIS:HA	1:B:177:PRO:HD3	1.71	0.44
1:C:245:THR:OG1	1:C:246:THR:N	2.49	0.43
1:C:298:TRP:CE2	1:C:320:ARG:HD3	2.53	0.43
1:B:314:GLU:O	1:B:318:SER:HB3	2.18	0.42
1:B:25:ILE:HG21	1:B:203:ALA:HB2	2.00	0.42
1:B:261:ASP:O	1:B:310:GLU:HG2	2.19	0.42
1:B:246:THR:O	1:B:246:THR:CG2	2.68	0.41
1:B:77:ARG:O	1:B:81:GLU:HB2	2.21	0.40
1:B:189:TRP:CD2	1:B:197:LEU:HD13	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/353 (87%)	292 (95%)	13 (4%)	1 (0%)	41	72
1	B	307/353 (87%)	294 (96%)	11 (4%)	2 (1%)	22	53
1	C	307/353 (87%)	289 (94%)	17 (6%)	1 (0%)	41	72
All	All	920/1059 (87%)	875 (95%)	41 (4%)	4 (0%)	34	66

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	LEU
1	B	146	LEU
1	C	148	ASP
1	B	96	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/295 (82%)	236 (97%)	7 (3%)	42	76
1	B	239/295 (81%)	230 (96%)	9 (4%)	33	67
1	C	235/295 (80%)	226 (96%)	9 (4%)	33	67
All	All	717/885 (81%)	692 (96%)	25 (4%)	36	70

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	SER
1	A	77	ARG
1	A	125	VAL
1	A	147	CYS
1	A	186	LYS
1	A	262	THR
1	A	323	VAL
1	B	76	THR
1	B	105	GLU
1	B	147	CYS
1	B	151	HIS
1	B	220	GLU
1	B	233	ARG
1	B	286	SER
1	B	306	ARG
1	B	320	ARG
1	C	101	LEU
1	C	107	LEU
1	C	175	THR
1	C	185	LEU
1	C	205	THR
1	C	262	THR
1	C	281	ASP
1	C	291	ARG
1	C	306	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/353 (88%)	-0.71	2 (0%) 89 86	34, 49, 73, 97	0
1	B	315/353 (89%)	-0.65	5 (1%) 72 66	34, 55, 80, 104	0
1	C	315/353 (89%)	-0.62	0 100 100	34, 57, 88, 115	0
All	All	944/1059 (89%)	-0.66	7 (0%) 87 84	34, 53, 82, 115	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	128	HIS	2.8
1	A	127	PRO	2.7
1	B	17	SER	2.5
1	B	167	ALA	2.5
1	B	129	SER	2.4
1	B	103	VAL	2.4
1	A	102	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	350	1/1	0.86	0.23	78,78,78,78	0
2	CL	A	351	1/1	0.95	0.12	71,71,71,71	1
2	CL	B	351	1/1	0.95	0.15	65,65,65,65	1
2	CL	B	350	1/1	0.97	0.07	66,66,66,66	0
2	CL	C	350	1/1	0.98	0.09	54,54,54,54	0
2	CL	C	351	1/1	0.99	0.16	64,64,64,64	1

6.5 Other polymers [i](#)

There are no such residues in this entry.