



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 15, 2023 – 03:07 PM EDT

PDB ID : 1SEJ  
Title : Crystal Structure of Dihydrofolate Reductase-Thymidylate Synthase from *Cryptosporidium hominis* Bound to 1843U89/NADPH/dUMP  
Authors : Anderson, A.C.  
Deposited on : 2004-02-17  
Resolution : 2.87 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	 2% 60% 33% 6%
1	B	521	 2% 60% 32% 7%
1	C	521	 2% 60% 32% 6%
1	D	521	 2% 61% 32% 6%
1	E	521	 3% 62% 31% 6%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 22194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called bifunctional dihydrofolate reductase-thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	519	Total 4223	C 2694	N 713	O 793	S 23	0	0	0
1	B	519	Total 4223	C 2694	N 713	O 793	S 23	0	0	0
1	C	519	Total 4223	C 2694	N 713	O 793	S 23	0	0	0
1	D	519	Total 4223	C 2694	N 713	O 793	S 23	0	0	0
1	E	519	Total 4223	C 2694	N 713	O 793	S 23	0	0	0

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: C<sub>9</sub>H<sub>13</sub>N<sub>2</sub>O<sub>8</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 20	C 9	N 2	O 8	P 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	C	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	D	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	E	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is S)-2-(5(((1,2-DIHYDRO-3-METHYL-1-OXOBENZO(F)QUINAZOLIN-9-YL)METHYL)AMINO)1-OXO-2-ISOINDOLINYL)GLUTARIC ACID (three-letter code: F89) (formula: C<sub>27</sub>H<sub>24</sub>N<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total	C	N	O	0	0
			37	27	4	6		
3	A	1	Total	C	N	O	0	0
			37	27	4	6		
3	B	1	Total	C	N	O	0	0
			37	27	4	6		
3	B	1	Total	C	N	O	0	0
			37	27	4	6		
3	C	1	Total	C	N	O	0	0
			37	27	4	6		
3	C	1	Total	C	N	O	0	0
			37	27	4	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O	0	0
			37	27	4	6		
3	D	1	Total	C	N	O	0	0
			37	27	4	6		
3	E	1	Total	C	N	O	0	0
			37	27	4	6		
3	E	1	Total	C	N	O	0	0
			37	27	4	6		

- Molecule 4 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
4	E	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	93	Total O 93 93	0	0
5	B	101	Total O 101 101	0	0
5	C	77	Total O 77 77	0	0
5	D	67	Total O 67 67	0	0
5	E	31	Total O 31 31	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: bifunctional dihydrofolate reductase-thymidylate synthase




- Molecule 1: bifunctional dihydrofolate reductase-thymidylate synthase







• Molecule 1: bifunctional dihydrofolate reductase-thymidylate synthase




• Molecule 1: bifunctional dihydrofolate reductase-thymidylate synthase







● Molecule 1: bifunctional dihydrofolate reductase-thymidylate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	214.90Å 116.30Å 219.70Å 90.00° 95.23° 90.00°	Depositor
Resolution (Å)	45.15 – 2.87 45.15 – 2.87	Depositor EDS
% Data completeness (in resolution range)	90.4 (45.15-2.87) 90.5 (45.15-2.87)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 2.86Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.220 , 0.237 0.215 , 0.233	Depositor DCC
$R_{free}$ test set	11185 reflections (9.40%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.8	Xtrriage
Anisotropy	0.267	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	22194	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F89, UMP, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/4320	0.70	5/5838 (0.1%)
1	B	0.41	0/4320	0.73	6/5838 (0.1%)
1	C	0.40	0/4320	0.70	5/5838 (0.1%)
1	D	0.40	0/4320	0.70	4/5838 (0.1%)
1	E	0.42	0/4320	0.73	5/5838 (0.1%)
All	All	0.41	0/21600	0.71	25/29190 (0.1%)

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	E	257	ARG	NE-CZ-NH2	-14.20	113.20	120.30
1	E	257	ARG	NE-CZ-NH1	13.77	127.18	120.30
1	B	257	ARG	NE-CZ-NH2	-13.71	113.44	120.30
1	B	257	ARG	NE-CZ-NH1	13.56	127.08	120.30
1	A	257	ARG	NE-CZ-NH1	-9.18	115.71	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4223	0	4159	218	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4223	0	4159	211	0
1	C	4223	0	4159	207	0
1	D	4223	0	4159	206	0
1	E	4223	0	4159	197	0
2	A	20	0	11	2	0
2	B	20	0	11	3	0
2	C	20	0	11	3	0
2	D	20	0	11	2	0
2	E	20	0	11	2	0
3	A	74	0	44	16	0
3	B	74	0	44	13	0
3	C	74	0	44	15	0
3	D	74	0	44	17	0
3	E	74	0	44	17	0
4	A	48	0	26	10	0
4	B	48	0	26	8	0
4	C	48	0	26	8	0
4	D	48	0	26	8	0
4	E	48	0	26	8	0
5	A	93	0	0	2	0
5	B	101	0	0	3	0
5	C	77	0	0	4	0
5	D	67	0	0	4	0
5	E	31	0	0	1	0
All	All	22194	0	21200	1028	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 1028 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:341:ILE:HA	1:E:397:MET:HE3	1.32	1.10
1:C:341:ILE:HA	1:C:397:MET:HE3	1.35	1.09
1:A:341:ILE:HA	1:A:397:MET:HE3	1.34	1.07
1:D:341:ILE:HA	1:D:397:MET:HE3	1.34	1.06
1:B:341:ILE:HA	1:B:397:MET:HE3	1.34	1.04

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/521 (99%)	472 (91%)	36 (7%)	9 (2%)	9	28
1	B	517/521 (99%)	472 (91%)	36 (7%)	9 (2%)	9	28
1	C	517/521 (99%)	470 (91%)	38 (7%)	9 (2%)	9	28
1	D	517/521 (99%)	473 (92%)	35 (7%)	9 (2%)	9	28
1	E	517/521 (99%)	469 (91%)	39 (8%)	9 (2%)	9	28
All	All	2585/2605 (99%)	2356 (91%)	184 (7%)	45 (2%)	9	28

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	ALA
1	A	102	MET
1	A	179	GLU
1	A	182	THR
1	B	84	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	468/470 (100%)	432 (92%)	36 (8%)	13	34
1	B	468/470 (100%)	431 (92%)	37 (8%)	12	32
1	C	468/470 (100%)	432 (92%)	36 (8%)	13	34
1	D	468/470 (100%)	433 (92%)	35 (8%)	13	35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	468/470 (100%)	434 (93%)	34 (7%)	14	36
All	All	2340/2350 (100%)	2162 (92%)	178 (8%)	13	35

5 of 178 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	103	ASN
1	D	516	LYS
1	D	133	LEU
1	D	295	LEU
1	E	103	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 104 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	412	ASN
1	D	256	ASN
1	E	412	ASN
1	C	422	GLN
1	D	81	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NDP	C	614	-	45,52,52	1.79	10 (22%)	53,80,80	1.54	11 (20%)
3	F89	D	616	-	40,41,41	3.91	9 (22%)	53,60,60	3.92	17 (32%)
2	UMP	D	615	-	21,21,21	2.87	8 (38%)	31,31,31	2.19	9 (29%)
3	F89	B	608	-	40,41,41	3.91	9 (22%)	53,60,60	3.90	18 (33%)
3	F89	E	621	-	40,41,41	3.90	9 (22%)	53,60,60	3.92	17 (32%)
2	UMP	A	603	-	21,21,21	2.84	8 (38%)	31,31,31	2.19	9 (29%)
4	NDP	D	618	-	45,52,52	1.78	8 (17%)	53,80,80	1.54	11 (20%)
3	F89	D	617	-	40,41,41	3.90	9 (22%)	53,60,60	3.92	17 (32%)
2	UMP	C	611	-	21,21,21	2.81	7 (33%)	31,31,31	2.19	9 (29%)
3	F89	C	613	-	40,41,41	3.90	9 (22%)	53,60,60	3.92	16 (30%)
4	NDP	E	622	-	45,52,52	1.79	10 (22%)	53,80,80	1.54	11 (20%)
4	NDP	B	610	-	45,52,52	1.76	9 (20%)	53,80,80	1.55	11 (20%)
2	UMP	B	607	-	21,21,21	2.83	7 (33%)	31,31,31	2.19	9 (29%)
3	F89	B	609	-	40,41,41	3.90	9 (22%)	53,60,60	3.94	17 (32%)
2	UMP	E	619	-	21,21,21	2.84	7 (33%)	31,31,31	2.18	9 (29%)
3	F89	A	604	-	40,41,41	3.90	9 (22%)	53,60,60	3.88	17 (32%)
4	NDP	A	606	-	45,52,52	1.79	9 (20%)	53,80,80	1.54	12 (22%)
3	F89	C	612	-	40,41,41	3.91	9 (22%)	53,60,60	3.90	17 (32%)
3	F89	A	605	-	40,41,41	3.91	9 (22%)	53,60,60	3.93	17 (32%)
3	F89	E	620	-	40,41,41	3.90	9 (22%)	53,60,60	3.85	16 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NDP	C	614	-	-	5/30/77/77	0/5/5/5
3	F89	D	616	-	-	4/18/30/30	0/5/5/5
2	UMP	D	615	-	-	1/10/22/22	0/2/2/2
3	F89	B	608	-	-	4/18/30/30	0/5/5/5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	F89	E	621	-	-	4/18/30/30	0/5/5/5
2	UMP	A	603	-	-	1/10/22/22	0/2/2/2
4	NDP	D	618	-	-	5/30/77/77	0/5/5/5
3	F89	D	617	-	-	4/18/30/30	0/5/5/5
2	UMP	C	611	-	-	1/10/22/22	0/2/2/2
3	F89	C	613	-	-	4/18/30/30	0/5/5/5
4	NDP	E	622	-	-	5/30/77/77	0/5/5/5
4	NDP	B	610	-	-	5/30/77/77	0/5/5/5
2	UMP	B	607	-	-	1/10/22/22	0/2/2/2
3	F89	B	609	-	-	4/18/30/30	0/5/5/5
2	UMP	E	619	-	-	1/10/22/22	0/2/2/2
3	F89	A	604	-	-	4/18/30/30	0/5/5/5
4	NDP	A	606	-	-	5/30/77/77	0/5/5/5
3	F89	C	612	-	-	4/18/30/30	0/5/5/5
3	F89	A	605	-	-	4/18/30/30	0/5/5/5
3	F89	E	620	-	-	4/18/30/30	0/5/5/5

The worst 5 of 173 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	612	F89	C19-N	-17.63	1.33	1.46
3	B	608	F89	C19-N	-17.59	1.33	1.46
3	E	620	F89	C19-N	-17.54	1.33	1.46
3	D	616	F89	C19-N	-17.54	1.33	1.46
3	A	604	F89	C19-N	-17.54	1.33	1.46

The worst 5 of 270 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	621	F89	C15-C19-N	21.81	109.45	102.18
3	B	609	F89	C15-C19-N	21.73	109.42	102.18
3	D	617	F89	C15-C19-N	21.70	109.41	102.18
3	C	613	F89	C15-C19-N	21.65	109.39	102.18
3	A	605	F89	C15-C19-N	21.64	109.39	102.18

There are no chirality outliers.

5 of 70 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
3	A	604	F89	CB-CA-N-C
3	A	605	F89	CB-CA-N-C19
3	B	608	F89	CB-CA-N-C
3	B	609	F89	CB-CA-N-C19
3	C	612	F89	CB-CA-N-C


There are no ring outliers.


20 monomers are involved in 122 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	614	NDP	8	0
3	D	616	F89	8	0
2	D	615	UMP	2	0
3	B	608	F89	6	0
3	E	621	F89	10	0
2	A	603	UMP	2	0
4	D	618	NDP	8	0
3	D	617	F89	9	0
2	C	611	UMP	3	0
3	C	613	F89	8	0
4	E	622	NDP	8	0
4	B	610	NDP	8	0
2	B	607	UMP	3	0
3	B	609	F89	7	0
2	E	619	UMP	2	0
3	A	604	F89	7	0
4	A	606	NDP	10	0
3	C	612	F89	7	0
3	A	605	F89	9	0
3	E	620	F89	7	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

































## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	519/521 (99%)	-0.28	10 (1%) 66 65	16, 30, 54, 76	0
1	B	519/521 (99%)	-0.25	9 (1%) 70 70	16, 29, 51, 76	0
1	C	519/521 (99%)	-0.31	9 (1%) 70 70	17, 30, 52, 76	0
1	D	519/521 (99%)	-0.29	13 (2%) 57 55	18, 31, 55, 76	0
1	E	519/521 (99%)	-0.19	15 (2%) 51 48	19, 32, 55, 76	0
All	All	2595/2605 (99%)	-0.26	56 (2%) 62 60	16, 31, 54, 76	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	521	VAL	6.1
1	D	521	VAL	6.0
1	B	180	LYS	5.1
1	D	182	THR	5.0
1	B	521	VAL	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	F89	D	616	37/37	0.83	0.28	42,50,56,56	0
3	F89	A	604	37/37	0.84	0.27	42,50,56,56	0
3	F89	E	620	37/37	0.84	0.36	42,50,56,56	0
3	F89	C	612	37/37	0.85	0.27	42,50,56,56	0
3	F89	B	608	37/37	0.86	0.28	42,50,56,56	0
3	F89	D	617	37/37	0.91	0.23	42,50,56,56	0
3	F89	C	613	37/37	0.91	0.23	42,50,56,56	0
3	F89	E	621	37/37	0.91	0.21	42,50,56,56	0
3	F89	B	609	37/37	0.92	0.24	42,50,56,56	0
3	F89	A	605	37/37	0.93	0.24	42,50,56,56	0
2	UMP	D	615	20/20	0.94	0.20	52,64,69,69	0
2	UMP	B	607	20/20	0.95	0.22	47,61,67,67	0
2	UMP	E	619	20/20	0.95	0.17	54,64,69,69	0
2	UMP	C	611	20/20	0.95	0.21	49,61,68,68	0
4	NDP	C	614	48/48	0.95	0.17	31,43,53,54	0
2	UMP	A	603	20/20	0.96	0.24	50,63,68,69	0
4	NDP	D	618	48/48	0.96	0.17	33,42,52,52	0
4	NDP	E	622	48/48	0.96	0.14	33,43,52,53	0
4	NDP	B	610	48/48	0.98	0.17	29,38,49,50	0
4	NDP	A	606	48/48	0.98	0.17	29,38,48,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around F89 D 616:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around F89 A 604:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around F89 E 620:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around F89 C 612:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around F89 B 608:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around F89 D 617:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around F89 C 613:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around F89 E 621:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around F89 B 609:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around F89 A 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UMP D 615:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UMP B 607:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UMP E 619:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around UMP C 611:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NDP C 614:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UMP A 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around NDP D 618:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around NDP E 622:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NDP B 610:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.