



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 01:05 AM EDT

PDB ID : 7SF9  
Title : Branchiostoma Floridae Violet Fluorescent Protein  
Authors : Pletnev, S.; Pletneva, N.; Pletnev, V.Z.; Muslinkina, L.  
Deposited on : 2021-10-03  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	
1	B	238	
1	C	238	
1	D	238	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Violet Fluorescent Protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	228	Total 1857	C 1183	N 310	O 353	S 11	0	6	0
1	B	218	Total 1784	C 1142	N 300	O 332	S 10	0	5	0
1	C	218	Total 1765	C 1130	N 295	O 330	S 10	0	3	0
1	D	219	Total 1750	C 1122	N 290	O 328	S 10	0	1	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	108	Total 108	O 108	0	0
2	B	137	Total 137	O 137	0	0
2	C	93	Total 93	O 93	0	0
2	D	65	Total 65	O 65	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Violet Fluorescent Protein




- Molecule 1: Violet Fluorescent Protein





- Molecule 1: Violet Fluorescent Protein



- Molecule 1: Violet Fluorescent Protein





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	177.67Å 177.67Å 52.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	153.87 – 1.80 29.61 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.7 (153.87-1.80) 95.7 (29.61-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.209 , 0.263 0.219 , 0.272	Depositor DCC
$R_{free}$ test set	1647 reflections (1.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtrriage
Anisotropy	0.041	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.018 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7559	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PQ4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	0/1895	0.96	2/2557 (0.1%)
1	B	1.02	0/1822	0.99	2/2461 (0.1%)
1	C	0.96	1/1803 (0.1%)	0.97	3/2436 (0.1%)
1	D	0.88	0/1788	0.99	3/2417 (0.1%)
All	All	0.95	1/7308 (0.0%)	0.98	10/9871 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	106	SER	CB-OG	-6.67	1.33	1.42

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	204	ARG	NE-CZ-NH2	-6.49	117.06	120.30
1	D	97	ARG	CB-CG-CD	-5.80	96.51	111.60
1	A	226	ASP	CB-CG-OD1	5.67	123.40	118.30
1	D	157	ASP	CB-CG-OD1	5.58	123.32	118.30
1	C	204	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	226	ASP	CB-CG-OD2	-5.51	113.34	118.30
1	B	204	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	D	180	MET	CG-SD-CE	-5.35	91.64	100.20

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	226	ASP	CB-CG-OD1	5.25	123.03	118.30
1	C	165	ASP	CB-CG-OD2	5.18	122.96	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	215	LEU	Mainchain
1	B	97	ARG	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1857	0	1777	15	0
1	B	1784	0	1718	15	0
1	C	1765	0	1693	24	0
1	D	1750	0	1679	37	0
2	A	108	0	0	1	0
2	B	137	0	0	0	0
2	C	93	0	0	5	0
2	D	65	0	0	1	0
All	All	7559	0	6867	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:69:ALA:CB	1:D:124:PHE:HE2	1.79	0.95
1:D:31:MET:CE	1:D:48:LEU:HD11	2.01	0.91
1:B:125:GLN:HE22	1:C:125:GLN:HE22	1.22	0.88
1:D:17:GLU:HG2	1:D:121:LYS:HG2	1.57	0.85
1:D:69:ALA:CB	1:D:124:PHE:CE2	2.60	0.84

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:GLN:HE21	1:A:120:ILE:HD13	1.41	0.84
1:D:31:MET:HE2	1:D:48:LEU:HD11	1.60	0.83
1:D:69:ALA:HB1	1:D:124:PHE:CE2	2.15	0.82
1:C:33:GLY:HA3	1:C:48:LEU:HD23	1.61	0.79
1:C:37:GLY:HA3	1:C:44:GLU:OE1	1.85	0.77
1:D:69:ALA:HB2	1:D:124:PHE:HE2	1.48	0.77
1:D:17:GLU:CG	1:D:121:LYS:HG2	2.16	0.75
1:C:37:GLY:CA	1:C:44:GLU:OE1	2.35	0.75
1:B:98[A]:THR:HG21	2:C:303:HOH:O	1.90	0.72
1:A:174:LYS:HE2	2:A:302:HOH:O	1.91	0.70
1:B:94:GLN:HG2	1:B:112[A]:ARG:HE	1.57	0.70
1:D:33:GLY:HA3	1:D:48:LEU:HB3	1.76	0.67
1:B:98[B]:THR:HG21	2:C:305:HOH:O	1.93	0.67
1:A:72:GLN:NE2	1:A:120:ILE:HD13	2.09	0.67
1:B:27:VAL:HG11	1:B:56:LYS:HE3	1.75	0.66
1:D:31:MET:HE3	1:D:48:LEU:HD11	1.79	0.63
1:C:33:GLY:CA	1:C:48:LEU:HD23	2.28	0.63
1:D:32:VAL:O	1:D:48:LEU:HB2	2.01	0.61
1:D:31:MET:CE	1:D:48:LEU:CD1	2.76	0.60
1:D:68:PQ4:CB	1:D:97:ARG:HH12	2.14	0.59
1:D:31:MET:HE3	1:D:48:LEU:CD1	2.33	0.58
1:D:17:GLU:CD	1:D:121:LYS:HG2	2.23	0.58
1:C:97:ARG:CB	1:C:182:GLU:HB2	2.34	0.57
1:C:207:GLU:HG3	1:C:220:TRP:HZ3	1.67	0.57
1:C:100[A]:GLN:HE22	1:C:106:SER:HB2	1.69	0.57
1:D:16:HIS:CE1	1:D:36:THR:N	2.73	0.56
1:D:19:HIS:ND1	1:D:123:GLU:OE1	2.38	0.56
1:C:37:GLY:HA2	1:C:44:GLU:OE1	2.06	0.55
1:C:97:ARG:HB3	1:C:182:GLU:HB2	1.88	0.55
1:D:69:ALA:HB1	1:D:124:PHE:CZ	2.42	0.55
1:B:181[B]:ARG:HH11	1:B:181[B]:ARG:HG2	1.72	0.54
1:A:34:GLU:OE1	1:A:34:GLU:N	2.41	0.54
1:C:207:GLU:HG3	1:C:220:TRP:CZ3	2.41	0.54
1:D:31:MET:HA	1:D:51:THR:HG23	1.90	0.53
1:D:38:ASN:ND2	1:D:41:GLU:OE2	2.41	0.53
1:B:181[B]:ARG:HG2	1:B:181[B]:ARG:NH1	2.24	0.53
1:A:95:VAL:HB	1:A:111:TYR:HB3	1.91	0.52
1:A:96:HIS:CE1	1:A:110:HIS:CD2	2.98	0.52
1:C:17:GLU:OE2	1:C:119:HIS:NE2	2.30	0.51
1:D:31:MET:HB2	1:D:48:LEU:HD13	1.91	0.51
1:B:97:ARG:HB3	1:B:182:GLU:HB2	1.93	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:16:HIS:HE1	1:D:36:THR:N	2.09	0.50
1:B:98[B]:THR:HG23	1:B:98[B]:THR:O	2.11	0.50
1:D:17:GLU:OE2	1:D:121:LYS:HE2	2.11	0.50
1:D:42:GLY:O	1:D:221:GLN:HG3	2.12	0.50
1:B:45:GLU:HG2	1:B:218:LYS:HG2	1.93	0.50
1:C:33:GLY:HA3	1:C:47:LYS:O	2.12	0.50
1:C:95:VAL:HB	1:C:111:TYR:HB3	1.94	0.49
1:A:70:PHE:HB3	1:A:73:TYR:HD2	1.78	0.49
1:D:68:PQ4:CG	1:D:97:ARG:HH12	2.25	0.49
1:D:29:PHE:HA	1:D:52:ASN:HD21	1.77	0.48
1:D:31:MET:HG3	1:D:48:LEU:HD13	1.96	0.48
1:B:27:VAL:HG11	1:B:56:LYS:CE	2.44	0.47
1:A:38:ASN:ND2	1:A:41:GLU:HG3	2.30	0.47
1:C:100[A]:GLN:NE2	2:C:303:HOH:O	2.47	0.47
1:A:61:ILE:HD11	1:A:101:TYR:CZ	2.50	0.47
1:A:150:THR:O	1:A:201:PHE:HA	2.16	0.46
1:D:54:PRO:HB3	1:D:212:LYS:HG3	1.98	0.46
1:B:47:LYS:O	1:B:47:LYS:HG3	2.14	0.46
1:D:78:ASP:HB2	2:D:315:HOH:O	2.16	0.46
1:B:95:VAL:HB	1:B:111:TYR:HB3	1.97	0.46
1:D:17:GLU:CD	1:D:121:LYS:HE2	2.36	0.46
1:B:33:GLY:HA3	1:B:47:LYS:O	2.14	0.45
1:A:13:PRO:CG	1:A:120:ILE:HD11	2.47	0.44
1:D:95:VAL:HG22	1:D:184:ASN:OD1	2.17	0.44
1:C:48:LEU:HB2	1:C:215:LEU:HG	1.99	0.44
1:D:42:GLY:HA2	1:D:73:TYR:O	2.17	0.44
1:D:207:GLU:HG2	1:D:209:GLN:HE22	1.83	0.43
1:C:100[A]:GLN:NE2	2:C:305:HOH:O	2.50	0.43
1:C:33:GLY:N	1:C:48:LEU:HD23	2.33	0.43
1:A:42:GLY:HA2	1:A:73:TYR:O	2.19	0.43
1:B:94:GLN:HG2	1:B:112[A]:ARG:NE	2.30	0.43
1:D:99:ILE:HG12	1:D:180:MET:HG3	2.01	0.42
1:C:132:PRO:HA	1:C:133:PRO:HD3	1.91	0.42
1:C:212:LYS:HA	2:C:311:HOH:O	2.20	0.42
1:D:31:MET:CG	1:D:48:LEU:HD13	2.50	0.42
1:C:17:GLU:CG	1:C:121:LYS:HG2	2.49	0.42
1:D:72:GLN:OE1	1:D:72:GLN:N	2.46	0.42
1:C:68:PQ4:CG	1:C:97:ARG:HH12	2.32	0.41
1:C:213:THR:O	1:C:214:GLU:HB3	2.20	0.41
1:D:123:GLU:C	1:D:124:PHE:HD1	2.24	0.41
1:C:109:ALA:HA	1:C:125:GLN:O	2.19	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:PRO:HG2	1:A:120:ILE:HD11	2.02	0.41
1:A:59:PRO:HG2	1:A:210:HIS:CD2	2.57	0.40
1:A:215:LEU:HD12	1:A:215:LEU:C	2.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/238 (97%)	226 (98%)	5 (2%)	0	100	100
1	B	220/238 (92%)	217 (99%)	2 (1%)	1 (0%)	29	15
1	C	218/238 (92%)	213 (98%)	4 (2%)	1 (0%)	29	15
1	D	217/238 (91%)	215 (99%)	2 (1%)	0	100	100
All	All	886/952 (93%)	871 (98%)	13 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	42	GLY
1	C	214	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/203 (98%)	197 (98%)	3 (2%)	65	56
1	B	192/203 (95%)	186 (97%)	6 (3%)	40	25
1	C	190/203 (94%)	186 (98%)	4 (2%)	53	42
1	D	188/203 (93%)	184 (98%)	4 (2%)	53	42
All	All	770/812 (95%)	753 (98%)	17 (2%)	50	39

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	ARG
1	A	124	PHE
1	A	129	THR
1	B	14	LYS
1	B	41	GLU
1	B	97	ARG
1	B	124	PHE
1	B	140	ASN
1	B	182	GLU
1	C	97	ARG
1	C	124	PHE
1	C	140	ASN
1	C	182	GLU
1	D	31	MET
1	D	41	GLU
1	D	71	TYR
1	D	221	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	HIS
1	A	110	HIS
1	A	210	HIS
1	B	88	HIS
1	B	140	ASN
1	C	125	GLN
1	D	110	HIS
1	D	177	GLN
1	D	209	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PQ4	A	68	1	12,12,13	1.83	4 (33%)	12,15,17	1.16	0
1	PQ4	C	68	1	12,12,13	1.02	1 (8%)	12,15,17	1.38	2 (16%)
1	PQ4	B	68	1	12,12,13	1.02	1 (8%)	12,15,17	1.74	3 (25%)
1	PQ4	D	68	1	12,12,13	1.55	2 (16%)	12,15,17	1.64	3 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PQ4	A	68	1	-	1/4/6/8	0/1/1/1
1	PQ4	C	68	1	-	1/4/6/8	0/1/1/1
1	PQ4	B	68	1	-	0/4/6/8	0/1/1/1
1	PQ4	D	68	1	-	1/4/6/8	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	68	PQ4	C-CA	3.13	1.50	1.45
1	A	68	PQ4	CE1-CD1	2.94	1.44	1.38
1	D	68	PQ4	CA-N	2.72	1.42	1.35
1	A	68	PQ4	CD2-CG	2.60	1.44	1.39
1	C	68	PQ4	C-CA	2.44	1.49	1.45
1	D	68	PQ4	OH-CZ	-2.38	1.31	1.37

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	68	PQ4	CG-CB	2.18	1.51	1.46
1	B	68	PQ4	C-CA	2.02	1.48	1.45

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	68	PQ4	CD1-CE1-CZ	3.61	123.84	119.88
1	D	68	PQ4	CD2-CE2-CZ	-3.24	116.33	119.88
1	C	68	PQ4	O-C-CA	-3.07	121.48	125.39
1	D	68	PQ4	CE1-CD1-CG	-2.88	117.50	121.25
1	D	68	PQ4	CD2-CG-CD1	2.49	121.32	117.64
1	B	68	PQ4	CE2-CD2-CG	2.27	124.20	121.25
1	B	68	PQ4	CD1-CG-CB	2.19	128.70	121.22
1	C	68	PQ4	CE1-CD1-CG	-2.15	118.45	121.25

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	68	PQ4	O-C-CA-CB
1	C	68	PQ4	O-C-CA-CB
1	D	68	PQ4	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	68	PQ4	1	0
1	D	68	PQ4	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	227/238 (95%)	0.04	8 (3%) 44 38	8, 23, 43, 63	0
1	B	217/238 (91%)	0.04	8 (3%) 41 36	8, 17, 42, 79	0
1	C	217/238 (91%)	0.37	20 (9%) 9 6	9, 25, 62, 95	0
1	D	218/238 (91%)	0.67	25 (11%) 4 3	13, 32, 60, 111	0
All	All	879/952 (92%)	0.28	61 (6%) 16 13	8, 24, 54, 111	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	213	THR	8.4
1	D	210	HIS	8.1
1	C	40	ASN	8.0
1	D	212	LYS	7.5
1	C	39	PRO	6.8
1	C	212	LYS	6.6
1	D	214	GLU	6.4
1	D	211	SER	6.0
1	B	40	ASN	5.6
1	C	213	THR	5.3
1	D	27	VAL	5.2
1	B	11	SER	5.0
1	B	39	PRO	4.8
1	C	116	GLU	4.7
1	C	41	GLU	4.6
1	C	42	GLY	4.5
1	C	38	ASN	4.4
1	C	11	SER	4.3
1	C	117	GLY	4.3
1	C	52	ASN	4.2
1	B	41	GLU	4.1

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	4	LYS	4.1
1	D	124	PHE	3.9
1	C	228	MET	3.8
1	D	115	TYR	3.8
1	D	36	THR	3.6
1	C	220	TRP	3.5
1	A	7	ASP	3.4
1	C	115	TYR	3.2
1	D	69	ALA	3.2
1	D	40	ASN	3.1
1	D	228	MET	3.1
1	C	37	GLY	3.0
1	C	211	SER	3.0
1	B	12	LEU	3.0
1	D	116	GLU	3.0
1	C	43	SER	2.9
1	C	36	THR	2.8
1	A	116	GLU	2.7
1	D	118	SER	2.7
1	A	52	ASN	2.7
1	B	36	THR	2.7
1	D	70	PHE	2.7
1	D	117	GLY	2.6
1	A	5	GLU	2.6
1	D	42	GLY	2.6
1	C	118	SER	2.5
1	C	196	GLN	2.5
1	D	209	GLN	2.5
1	A	118	SER	2.5
1	B	174	LYS	2.4
1	D	193	ASP	2.3
1	D	13	PRO	2.3
1	D	45	GLU	2.3
1	D	14	LYS	2.3
1	A	6	GLU	2.2
1	D	220	TRP	2.2
1	D	12	LEU	2.1
1	D	38	ASN	2.1
1	A	134	ASP	2.1
1	B	52	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PQ4	D	68	12/13	0.77	0.21	28,34,45,55	0
1	PQ4	C	68	12/13	0.95	0.08	16,20,21,24	0
1	PQ4	A	68	12/13	0.96	0.13	11,14,18,19	0
1	PQ4	B	68	12/13	0.97	0.10	7,10,13,14	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.