



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2022 – 02:33 PM EDT

PDB ID : 5SKC  
Title : Crystal Structure of human phosphodiesterase 10 in complex with 6-methyl-N-(1-propan-2-ylpyrazol-3-yl)-3-(pyrimidin-5-ylamino)pyridine-2-carboxamide  
Authors : Joseph, C.; Benz, J.; Flohr, A.; Vieira, E.; Rudolph, M.G.  
Deposited on : 2022-02-01  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	 81% 9% 9%
1	B	343	 82% 9% 8%
1	C	343	 83% 8% 9%
1	D	343	 7% 82% 9% 9%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	Total 2549	C 1629	N 435	O 461	S 24	0	1	0
1	B	315	Total 2551	C 1630	N 434	O 463	S 24	0	0	0
1	C	313	Total 2549	C 1629	N 435	O 461	S 24	0	1	0
1	D	313	Total 2537	C 1622	N 433	O 458	S 24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233
B	447	GLY	-	expression tag	UNP Q9Y233
B	448	SER	-	expression tag	UNP Q9Y233
C	447	GLY	-	expression tag	UNP Q9Y233
C	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0
2	D	1	Total 1	Zn 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

- Molecule 4 is 6-methyl-N-[1-(propan-2-yl)-1H-pyrazol-3-yl]-3-[(pyrimidin-5-yl)amino]pyridine-2-carboxamide (three-letter code: KE6) (formula: C<sub>17</sub>H<sub>19</sub>N<sub>7</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 25 17 7 1	0	0
4	B	1	Total C N O 25 17 7 1	0	0
4	C	1	Total C N O 25 17 7 1	0	0
4	D	1	Total C N O 25 17 7 1	0	0



- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	33	Total 33	O 33	0	0
5	B	41	Total 41	O 41	0	0
5	C	41	Total 41	O 41	0	0
5	D	11	Total 11	O 11	0	0



### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A





- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A





## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	136.69Å 136.69Å 236.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.87 – 2.60 43.87 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.87-2.60) 98.7 (43.87-2.60)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.20 (at 2.61Å)	Xtrriage
Refinement program	BUSTER 2.11.7 (19-MAR-2020), REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.216 , 0.242 0.207 , 0.234	Depositor DCC
$R_{free}$ test set	2468 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.1	Xtrriage
Anisotropy	0.044	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10420	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KE6, CME, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/2603	0.55	0/3521
1	B	0.44	0/2602	0.56	0/3521
1	C	0.45	0/2603	0.56	0/3521
1	D	0.37	0/2588	0.53	0/3502
All	All	0.43	0/10396	0.55	0/14065

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2549	0	2526	16	0
1	B	2551	0	2515	9	0
1	C	2549	0	2524	12	0
1	D	2537	0	2511	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	25	0	0	1	0
4	B	25	0	0	0	0
4	C	25	0	0	0	0
4	D	25	0	0	1	0
5	A	33	0	0	0	0
5	B	41	0	0	0	0
5	C	41	0	0	0	0
5	D	11	0	0	0	0
All	All	10420	0	10076	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:520[A]:ARG:CZ	1:A:568:ARG:HD3	2.26	0.65
1:A:525:HIS:HD2	1:A:564:ASP:OD2	1.82	0.62
1:B:646:GLU:OE1	1:B:743:GLN:NE2	2.32	0.61
1:A:520[A]:ARG:NE	1:A:568:ARG:HD3	2.18	0.58
1:A:498:CYS:SG	1:A:554:LYS:HG3	2.45	0.56
1:C:752:LEU:O	1:C:756[B]:ARG:HG2	2.05	0.56
1:D:525:HIS:HD2	1:D:564:ASP:OD2	1.88	0.56
1:D:677:SER:HB3	4:D:803:KE6:N13	2.21	0.55
1:B:644:GLN:O	1:B:648:MET:HG3	2.08	0.53
1:A:727:LEU:HD21	1:A:763:GLU:HG3	1.89	0.53
1:B:515:VAL:HG13	1:B:565:LEU:HD21	1.90	0.53
1:C:493:MET:O	1:C:497:SER:HB2	2.08	0.52
1:B:492:TYR:CZ	1:B:496:ARG:HD2	2.45	0.52
1:C:498:CYS:SG	1:C:554:LYS:HG3	2.50	0.52
1:A:493:MET:O	1:A:497:SER:HB2	2.11	0.51
1:D:492:TYR:CZ	1:D:496:ARG:HD2	2.46	0.51
1:C:459:LEU:HD23	1:C:459:LEU:H	1.75	0.50
1:A:492:TYR:CZ	1:A:496:ARG:HD2	2.46	0.50
1:D:493:MET:O	1:D:497:SER:HB2	2.12	0.50
1:C:492:TYR:CZ	1:C:496:ARG:HD2	2.47	0.49
1:B:493:MET:O	1:B:497:SER:HB2	2.12	0.48
1:A:712:PRO:HA	1:A:718:LYS:HD2	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502:CYS:HB2	1:A:554:LYS:HZ3	1.80	0.47
1:C:712:PRO:HA	1:C:718:LYS:HD2	1.97	0.47
1:C:735:ILE:HD12	1:C:756[A]:ARG:HG2	1.96	0.46
1:D:712:PRO:HA	1:D:718:LYS:HD2	1.97	0.46
1:C:723:PRO:HB2	1:C:766:ILE:HG13	1.98	0.46
1:B:712:PRO:HA	1:B:718:LYS:HD2	1.98	0.45
1:D:479:ILE:HD11	1:D:528:LYS:HG3	2.00	0.44
1:D:544:ASN:HB3	1:D:547:LEU:HD13	2.00	0.44
1:A:510:ARG:HG2	1:A:608:HIS:CE1	2.53	0.43
1:C:544:ASN:HB3	1:C:547:LEU:HD13	2.00	0.43
1:D:510:ARG:HD2	1:D:607:GLY:O	2.18	0.43
1:A:697:TRP:HB3	1:A:716:ARG:HB3	2.00	0.43
1:B:525:HIS:HD2	1:B:564:ASP:OD2	2.01	0.43
1:D:730:TYR:HA	1:D:734:ALA:HB3	2.01	0.43
1:C:697:TRP:HB3	1:C:716:ARG:HB3	2.01	0.43
1:D:697:TRP:HB3	1:D:716:ARG:HB3	2.00	0.42
1:C:477:PHE:HB3	1:C:535:HIS:CE1	2.54	0.42
1:B:697:TRP:HB3	1:B:716:ARG:HB3	2.02	0.42
1:A:502:CYS:HB2	1:A:554:LYS:NZ	2.35	0.41
1:C:459:LEU:H	1:C:459:LEU:CD2	2.33	0.41
1:B:544:ASN:HB3	1:B:547:LEU:HD13	2.02	0.41
1:A:677:SER:HB3	4:A:803:KE6:N13	2.35	0.41
1:A:544:ASN:HB3	1:A:547:LEU:HD13	2.02	0.41
1:A:716:ARG:O	1:A:719:LYS:HG3	2.20	0.40
1:D:716:ARG:O	1:D:719:LYS:HG3	2.22	0.40
1:A:467:ARG:O	1:A:471:GLU:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/343 (91%)	301 (97%)	10 (3%)	0	100	100
1	B	312/343 (91%)	301 (96%)	11 (4%)	0	100	100
1	C	311/343 (91%)	300 (96%)	11 (4%)	0	100	100
1	D	310/343 (90%)	299 (96%)	10 (3%)	1 (0%)	41	64
All	All	1244/1372 (91%)	1201 (96%)	42 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	652	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/305 (92%)	269 (95%)	13 (5%)	27	51
1	B	281/305 (92%)	266 (95%)	15 (5%)	22	45
1	C	282/305 (92%)	269 (95%)	13 (5%)	27	51
1	D	280/305 (92%)	267 (95%)	13 (5%)	27	51
All	All	1125/1220 (92%)	1071 (95%)	54 (5%)	25	49

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	471	GLU
1	A	497	SER
1	A	504	GLU
1	A	517	LYS
1	A	554	LYS
1	A	573	SER
1	A	575	LEU
1	A	576	GLN
1	A	605	LEU
1	A	606	GLU

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	650	GLN
1	A	709	GLN
1	A	760	SER
1	B	459	LEU
1	B	471	GLU
1	B	497	SER
1	B	502	CYS
1	B	504	GLU
1	B	554	LYS
1	B	573	SER
1	B	575	LEU
1	B	576	GLN
1	B	605	LEU
1	B	606	GLU
1	B	650	GLN
1	B	709	GLN
1	B	727	LEU
1	B	760	SER
1	C	470	LYS
1	C	497	SER
1	C	504	GLU
1	C	517	LYS
1	C	554	LYS
1	C	573	SER
1	C	575	LEU
1	C	576	GLN
1	C	605	LEU
1	C	606	GLU
1	C	709	GLN
1	C	760	SER
1	C	766	ILE
1	D	460	MET
1	D	461	GLN
1	D	471	GLU
1	D	497	SER
1	D	504	GLU
1	D	554	LYS
1	D	573	SER
1	D	575	LEU
1	D	576	GLN
1	D	605	LEU
1	D	606	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	709	GLN
1	D	760	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	576	GLN
1	A	604	GLN
1	A	644	GLN
1	A	743	GLN
1	A	761	GLN
1	B	576	GLN
1	B	604	GLN
1	B	761	GLN
1	C	576	GLN
1	C	604	GLN
1	C	743	GLN
1	C	761	GLN
1	D	495	HIS
1	D	576	GLN
1	D	604	GLN
1	D	761	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CME	C	509	1	8,9,10	1.05	0	5,9,11	1.14	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CME	B	509	1	8,9,10	1.27	1 (12%)	5,9,11	0.93	0
1	CME	A	509	1	8,9,10	0.92	0	5,9,11	0.88	0
1	CME	D	509	1	8,9,10	1.00	0	5,9,11	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	C	509	1	-	3/5/8/10	-
1	CME	B	509	1	-	3/5/8/10	-
1	CME	A	509	1	-	3/5/8/10	-
1	CME	D	509	1	-	2/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	509	CME	CB-SG	2.31	1.88	1.81

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	509	CME	CE-SD-SG-CB
1	A	509	CME	SD-CE-CZ-OH
1	B	509	CME	CE-SD-SG-CB
1	C	509	CME	CE-SD-SG-CB
1	C	509	CME	SD-CE-CZ-OH
1	D	509	CME	CE-SD-SG-CB
1	D	509	CME	SD-CE-CZ-OH
1	B	509	CME	SD-CE-CZ-OH
1	A	509	CME	CZ-CE-SD-SG
1	B	509	CME	CZ-CE-SD-SG
1	C	509	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	KE6	A	803	-	25,27,27	0.34	0	27,37,37	1.03	3 (11%)
4	KE6	C	803	-	25,27,27	0.41	0	27,37,37	1.05	2 (7%)
4	KE6	D	803	-	25,27,27	0.53	0	27,37,37	0.98	2 (7%)
4	KE6	B	803	-	25,27,27	0.40	0	27,37,37	0.99	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	KE6	A	803	-	-	3/14/16/16	0/3/3/3
4	KE6	C	803	-	-	1/14/16/16	0/3/3/3
4	KE6	D	803	-	-	2/14/16/16	0/3/3/3
4	KE6	B	803	-	-	2/14/16/16	0/3/3/3

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	803	KE6	C7-C5-N2	3.13	112.65	110.92
4	B	803	KE6	C7-C5-N2	2.89	112.52	110.92
4	A	803	KE6	C7-C5-N2	2.82	112.48	110.92
4	D	803	KE6	C7-C5-N2	2.71	112.42	110.92

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	803	KE6	C10-N4-C17	2.52	127.89	125.40
4	D	803	KE6	C10-N4-C17	2.47	127.84	125.40
4	C	803	KE6	C10-N4-C17	2.34	127.71	125.40
4	A	803	KE6	C10-N4-C17	2.33	127.70	125.40
4	A	803	KE6	C22-C15-N11	2.21	127.80	120.19

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	803	KE6	C24-C17-N4-N2
4	A	803	KE6	C23-C17-N4-C10
4	A	803	KE6	C23-C17-N4-N2
4	B	803	KE6	C23-C17-N4-C10
4	C	803	KE6	C23-C17-N4-C10
4	D	803	KE6	C23-C17-N4-C10
4	B	803	KE6	C24-C17-N4-C10
4	D	803	KE6	C24-C17-N4-C10


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	803	KE6	1	0
4	D	803	KE6	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	312/343 (90%)	-0.09	2 (0%) 89 88	44, 65, 94, 115	0
1	B	314/343 (91%)	-0.14	4 (1%) 77 73	44, 65, 93, 120	0
1	C	312/343 (90%)	-0.28	1 (0%) 94 93	43, 62, 86, 107	0
1	D	312/343 (90%)	0.17	23 (7%) 14 10	69, 91, 114, 132	0
All	All	1250/1372 (91%)	-0.08	30 (2%) 59 53	43, 71, 107, 132	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	720	ASP	4.9
1	D	656	LEU	4.7
1	A	460	MET	4.2
1	C	770	GLU	4.1
1	D	719	LYS	4.0
1	D	456	TRP	3.9
1	D	718	LYS	3.5
1	D	457	GLN	3.3
1	B	770	GLU	3.2
1	D	505	LEU	3.1
1	D	654	LEU	2.9
1	D	649	TYR	2.7
1	D	550	ASP	2.7
1	A	458	GLY	2.5
1	D	712	PRO	2.5
1	D	767	ARG	2.5
1	D	553	ARG	2.4
1	D	765	VAL	2.4
1	D	665	VAL	2.4
1	D	711	ILE	2.3
1	B	460	MET	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	475	PHE	2.3
1	D	709	GLN	2.3
1	D	545	HIS	2.2
1	B	727	LEU	2.1
1	D	603	LEU	2.1
1	D	710	PRO	2.1
1	D	721	GLU	2.1
1	B	769	GLU	2.1
1	D	708	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CME	B	509	10/11	0.78	0.23	73,78,100,101	0
1	CME	D	509	10/11	0.83	0.27	91,96,146,146	0
1	CME	C	509	10/11	0.87	0.20	67,73,120,121	0
1	CME	A	509	10/11	0.89	0.17	71,77,106,106	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	KE6	D	803	25/25	0.90	0.29	94,94,97,98	0
3	MG	D	802	1/1	0.92	0.23	101,101,101,101	0
4	KE6	B	803	25/25	0.94	0.25	63,65,68,68	0
3	MG	A	802	1/1	0.94	0.14	47,47,47,47	0
4	KE6	A	803	25/25	0.95	0.18	70,71,75,75	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	KE6	C	803	25/25	0.96	0.20	62,63,67,68	0
2	ZN	D	801	1/1	0.98	0.11	75,75,75,75	0
3	MG	B	802	1/1	0.99	0.15	41,41,41,41	0
2	ZN	C	801	1/1	0.99	0.18	57,57,57,57	0
3	MG	C	802	1/1	1.00	0.15	40,40,40,40	0
2	ZN	A	801	1/1	1.00	0.17	59,59,59,59	0
2	ZN	B	801	1/1	1.00	0.14	54,54,54,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









**Electron density around KE6 B 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around KE6 A 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.