



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 01:51 pm BST

PDB ID : 3SMA  
Title : A new N-acetyltransferase fold in the structure and mechanism of the phosphonate biosynthetic enzyme FrbF  
Authors : Bae, B.; Nair, S.K.  
Deposited on : 2011-06-27  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	 % 83% 8% • 8%
1	B	286	 % 84% 8% • 6%
1	C	286	 % 79% 11% • 8%
1	D	286	 % 79% 12% • 8%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FrbF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	Total 2024	C 1266	N 374	O 374	S 10	0	0	0
1	B	268	Total 2067	C 1290	N 383	O 384	S 10	0	0	0
1	C	262	Total 2019	C 1263	N 373	O 373	S 10	0	0	0
1	D	264	Total 2029	C 1269	N 375	O 375	S 10	0	0	0

- Molecule 2 is ACETYL COENZYME \*A (three-letter code: ACO) (formula: C<sub>23</sub>H<sub>38</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	Total 51	C 23	N 7	O 17	P 3	S 1	0	0
2	B	1	Total 51	C 23	N 7	O 17	P 3	S 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	C	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	127	Total	O	0	0
			127	127		
3	B	122	Total	O	0	0
			122	122		
3	C	117	Total	O	0	0
			117	117		
3	D	145	Total	O	0	0
			145	145		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FrbF




- Molecule 1: FrbF



- Molecule 1: FrbF



- Molecule 1: FrbF



E196	P202	G203	R204	R208	E223	I227	D233	E239	S249	R250	C271	M274	T275	R278	R282	ALA	VAL	GLY	ALA
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.83Å 35.69Å 136.09Å 90.00° 117.60° 90.00°	Depositor
Resolution (Å)	45.34 – 2.00 45.35 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (45.34-2.00) 98.9 (45.35-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.32 (at 2.00Å)	Xtrriage
Refinement program	BUSTER 2.8.0	Depositor
R, $R_{free}$	0.193 , 0.246 0.199 , 0.253	Depositor DCC
$R_{free}$ test set	3758 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtrriage
Anisotropy	1.193	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8854	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/2076	0.70	0/2826
1	B	0.53	0/2120	0.72	0/2885
1	C	0.52	0/2071	0.69	0/2819
1	D	0.52	0/2081	0.71	0/2833
All	All	0.52	0/8348	0.70	0/11363

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2024	0	1952	25	0
1	B	2067	0	1989	17	0
1	C	2019	0	1947	27	0
1	D	2029	0	1954	26	0
2	A	51	0	34	0	0
2	B	51	0	34	1	0
2	C	51	0	34	2	0
2	D	51	0	34	2	0
3	A	127	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	122	0	0	1	1
3	C	117	0	0	4	1
3	D	145	0	0	2	0
All	All	8854	0	7978	88	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:ARG:HB2	1:A:93:PRO:HD3	1.50	0.92
1:D:92:ARG:HB3	1:D:93:PRO:HD3	1.52	0.91
1:A:233:ASP:OD2	1:A:278:ARG:HD3	1.76	0.85
1:A:129:ARG:HG2	1:B:129:ARG:HH11	1.45	0.80
1:C:141:ASN:HD21	1:C:162:ARG:HH11	1.31	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:287:HOH:O	3:C:302:HOH:O[2_556]	1.74	0.46

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	261/286 (91%)	254 (97%)	6 (2%)	1 (0%)	34 30
1	B	266/286 (93%)	257 (97%)	6 (2%)	3 (1%)	14 8
1	C	260/286 (91%)	254 (98%)	5 (2%)	1 (0%)	34 30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	D	262/286 (92%)	256 (98%)	5 (2%)	1 (0%)	34 30
All	All	1049/1144 (92%)	1021 (97%)	22 (2%)	6 (1%)	25 19

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	92	ARG
1	B	157	HIS
1	C	92	ARG
1	B	93	PRO
1	B	202	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	207/227 (91%)	196 (95%)	11 (5%)	22 18
1	B	212/227 (93%)	203 (96%)	9 (4%)	30 27
1	C	207/227 (91%)	197 (95%)	10 (5%)	25 22
1	D	207/227 (91%)	195 (94%)	12 (6%)	20 15
All	All	833/908 (92%)	791 (95%)	42 (5%)	24 20

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	224	GLN
1	C	128	ARG
1	D	223	GLU
1	B	278	ARG
1	C	51	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	189	ASN
1	C	138	HIS
1	D	207	HIS
1	C	106	GLN
1	C	141	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACO	A	300	-	45,53,53	1.67	11 (24%)	56,79,79	2.27	17 (30%)
2	ACO	B	300	-	45,53,53	1.84	12 (26%)	56,79,79	2.07	16 (28%)
2	ACO	C	300	-	45,53,53	1.42	6 (13%)	56,79,79	2.01	18 (32%)
2	ACO	D	300	-	45,53,53	1.47	8 (17%)	56,79,79	1.96	16 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACO	A	300	-	-	7/47/67/67	0/3/3/3
2	ACO	B	300	-	-	3/47/67/67	0/3/3/3
2	ACO	C	300	-	-	4/47/67/67	0/3/3/3
2	ACO	D	300	-	-	7/47/67/67	0/3/3/3

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	ACO	P3B-O3B	6.09	1.70	1.59
2	A	300	ACO	P3B-O3B	5.21	1.69	1.59
2	C	300	ACO	P3B-O3B	4.23	1.67	1.59
2	D	300	ACO	P3B-O3B	3.66	1.66	1.59
2	C	300	ACO	C4A-N3A	3.52	1.40	1.35

The worst 5 of 67 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	ACO	N3A-C2A-N1A	-6.70	118.21	128.68
2	B	300	ACO	N3A-C2A-N1A	-6.58	118.40	128.68
2	D	300	ACO	N3A-C2A-N1A	-5.94	119.39	128.68
2	A	300	ACO	C6P-C7P-N8P	-5.37	101.06	111.90
2	C	300	ACO	N3A-C2A-N1A	-5.29	120.42	128.68

There are no chirality outliers.

5 of 21 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	300	ACO	C3B-O3B-P3B-O9A
2	A	300	ACO	C5B-O5B-P1A-O2A
2	A	300	ACO	OAP-CAP-CBP-CEP
2	A	300	ACO	O-C-S1P-C2P
2	A	300	ACO	CH3-C-S1P-C2P


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	300	ACO	1	0
2	C	300	ACO	2	0
2	D	300	ACO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/286 (91%)	-0.29	2 (0%) 86 85	21, 31, 54, 80	0
1	B	268/286 (93%)	-0.26	2 (0%) 87 87	20, 31, 56, 86	0
1	C	262/286 (91%)	-0.25	3 (1%) 80 79	22, 32, 55, 84	0
1	D	264/286 (92%)	-0.29	2 (0%) 86 85	21, 33, 52, 84	0
All	All	1057/1144 (92%)	-0.27	9 (0%) 84 83	20, 32, 55, 86	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	92	ARG	4.7
1	A	92	ARG	4.5
1	C	92	ARG	3.6
1	A	93	PRO	2.9
1	C	37	ARG	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACO	D	300	51/51	0.93	0.10	29,37,61,63	0
2	ACO	C	300	51/51	0.94	0.10	25,35,55,59	0
2	ACO	B	300	51/51	0.95	0.10	28,32,55,57	0
2	ACO	A	300	51/51	0.96	0.09	24,33,60,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









## 6.5 Other polymers [i](#)

There are no such residues in this entry.