



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 3, 2023 – 10:17 PM EDT

PDB ID : 3SWS
Title : Crystal Structure of the Quinone Form of Methylamine Dehydrogenase in Complex with the Diferric Form of MauG
Authors : Jensen, L.M.R.; Wilmot, C.M.
Deposited on : 2011-07-14
Resolution : 1.86 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	83% 10% • 5%
1	B	373	83% 11% • 5%
2	C	137	78% 11% • 8%
2	E	137	81% 9% • 9%
3	D	386	88% 9% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	386	 87% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	EDO	D	388	-	-	X	-

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 15667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylamine utilization protein MauG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	354	2753	1721	494	527	11	0	3	0
1	B	354	2781	1740	495	534	12	0	8	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	368	HIS	-	expression tag	UNP Q51658
A	369	HIS	-	expression tag	UNP Q51658
A	370	HIS	-	expression tag	UNP Q51658
A	371	HIS	-	expression tag	UNP Q51658
A	372	HIS	-	expression tag	UNP Q51658
A	373	HIS	-	expression tag	UNP Q51658
B	368	HIS	-	expression tag	UNP Q51658
B	369	HIS	-	expression tag	UNP Q51658
B	370	HIS	-	expression tag	UNP Q51658
B	371	HIS	-	expression tag	UNP Q51658
B	372	HIS	-	expression tag	UNP Q51658
B	373	HIS	-	expression tag	UNP Q51658

- Molecule 2 is a protein called Methylamine dehydrogenase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	126	979	605	167	194	13	0	2	0
2	E	125	964	596	161	193	14	0	2	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	132	HIS	-	expression tag	UNP P22619
C	133	HIS	-	expression tag	UNP P22619
C	134	HIS	-	expression tag	UNP P22619
C	135	HIS	-	expression tag	UNP P22619
C	136	HIS	-	expression tag	UNP P22619
C	137	HIS	-	expression tag	UNP P22619
E	132	HIS	-	expression tag	UNP P22619
E	133	HIS	-	expression tag	UNP P22619
E	134	HIS	-	expression tag	UNP P22619
E	135	HIS	-	expression tag	UNP P22619
E	136	HIS	-	expression tag	UNP P22619
E	137	HIS	-	expression tag	UNP P22619

- Molecule 3 is a protein called Methylamine dehydrogenase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	376	Total	C	N	O	S	0	12	0
			2994	1904	508	574	8			
3	F	376	Total	C	N	O	S	0	5	0
			2952	1874	504	566	8			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		


- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Na	0	0
			2	2		
5	B	2	Total	Na	0	0
			2	2		

- Molecule 6 is HEME C (three-letter code: HEC) (formula: C₃₄H₃₄FeN₄O₄).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 9 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 10 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	D	1	Total	C	O	0	0
			16	10	6		

- Molecule 11 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	F	1	Total	C O	0	0
			13	8 5		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	359	Total	O	0	0
			359	359		
12	B	415	Total	O	0	0
			415	415		
12	C	151	Total	O	0	0
			151	151		
12	D	475	Total	O	0	0
			475	475		
12	E	131	Total	O	0	0
			131	131		
12	F	478	Total	O	0	0
			478	478		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methylamine utilization protein MauG




- Molecule 1: Methylamine utilization protein MauG




- Molecule 2: Methylamine dehydrogenase light chain




- Molecule 2: Methylamine dehydrogenase light chain





- Molecule 3: Methylamine dehydrogenase heavy chain

Chain D:  88% 9%



- Molecule 3: Methylamine dehydrogenase heavy chain

Chain F:  87% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.22Å 99.42Å 102.83Å 64.77° 74.77° 75.14°	Depositor
Resolution (Å)	49.12 – 1.86 49.11 – 1.86	Depositor EDS
% Data completeness (in resolution range)	97.2 (49.12-1.86) 97.1 (49.11-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.136 , 0.178 0.142 , 0.183	Depositor DCC
R_{free} test set	7734 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for h,h-k,h-l 0.009 for -h,-l,-k 0.004 for -h,-h+l,-h+k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15667	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, 1PE, PG4, HEC, TRQ, CA, MES, NA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	1/2826 (0.0%)	1.01	13/3834 (0.3%)
1	B	1.20	3/2869 (0.1%)	1.07	12/3891 (0.3%)
2	C	1.36	5/994 (0.5%)	1.09	3/1357 (0.2%)
2	E	1.21	2/978 (0.2%)	0.97	1/1335 (0.1%)
3	D	1.29	7/3107 (0.2%)	1.06	7/4231 (0.2%)
3	F	1.26	9/3044 (0.3%)	1.02	7/4147 (0.2%)
All	All	1.23	27/13818 (0.2%)	1.04	43/18795 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	62	TYR	CD2-CE2	7.32	1.50	1.39
3	D	48	TYR	CE1-CZ	6.86	1.47	1.38
3	D	342	GLU	C-O	6.15	1.35	1.23
3	F	287	TYR	CD1-CE1	6.03	1.48	1.39
3	F	242	TRP	CE3-CZ3	5.86	1.48	1.38

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	180	ARG	NE-CZ-NH1	11.31	125.95	120.30
1	B	180	ARG	NE-CZ-NH2	-9.07	115.77	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	118	ASP	CB-CG-OD1	7.79	125.31	118.30
3	D	224	ASP	CB-CG-OD1	7.58	125.12	118.30
3	F	104	ARG	NE-CZ-NH1	7.54	124.07	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	358	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2753	0	2639	26	0
1	B	2781	0	2672	32	0
2	C	979	0	883	10	0
2	E	964	0	868	8	0
3	D	2994	0	2906	20	0
3	F	2952	0	2847	17	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	86	0	60	3	0
6	B	86	0	60	1	0
7	A	4	0	6	0	0
7	B	4	0	6	1	0
7	D	4	0	6	9	0
8	D	12	0	12	1	0
9	D	4	0	3	0	0
10	D	16	0	22	5	0
11	F	13	0	18	6	0
12	A	359	0	0	4	0
12	B	415	0	0	9	0
12	C	151	0	0	3	0
12	D	475	0	0	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	E	131	0	0	2	0
12	F	478	0	0	9	0
All	All	15667	0	13008	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 117 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:372[A]:GLN:OE1	12:F:1498:HOH:O	1.62	1.17
1:B:26[A]:ASN:ND2	12:B:2258:HOH:O	1.82	1.10
1:B:46[B]:GLU:HG2	1:B:51:LEU:HD12	1.40	0.99
3:F:372[B]:GLN:NE2	12:F:1047:HOH:O	2.02	0.93
1:B:48:LYS:H	1:B:62:HIS:HE1	1.03	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/373 (95%)	346 (98%)	8 (2%)	1 (0%)	41	26
1	B	360/373 (96%)	350 (97%)	9 (2%)	1 (0%)	41	26
2	C	125/137 (91%)	120 (96%)	5 (4%)	0	100	100
2	E	124/137 (90%)	121 (98%)	3 (2%)	0	100	100
3	D	386/386 (100%)	372 (96%)	12 (3%)	2 (0%)	29	15
3	F	379/386 (98%)	365 (96%)	13 (3%)	1 (0%)	41	26
All	All	1729/1792 (96%)	1674 (97%)	50 (3%)	5 (0%)	47	26

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	7	ASP
1	A	358	LEU
3	D	102[A]	ILE
3	D	102[B]	ILE
3	F	102	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	279/292 (96%)	267 (96%)	12 (4%)	29 12
1	B	284/292 (97%)	279 (98%)	5 (2%)	59 45
2	C	107/112 (96%)	105 (98%)	2 (2%)	57 43
2	E	106/112 (95%)	103 (97%)	3 (3%)	43 27
3	D	316/311 (102%)	305 (96%)	11 (4%)	36 18
3	F	309/311 (99%)	304 (98%)	5 (2%)	62 49
All	All	1401/1430 (98%)	1363 (97%)	38 (3%)	47 29

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	316	LEU
3	F	293	ARG
3	D	354	LYS
2	E	131	SER
3	F	354	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	54	HIS
1	B	210	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	62	HIS
1	A	210	GLN
1	B	84	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TRQ	C	57	2	13,17,18	3.14	6 (46%)	14,24,26	2.88	6 (42%)
2	TRQ	E	57	2	13,17,18	3.18	3 (23%)	14,24,26	2.80	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRQ	C	57	2	-	0/4/19/21	0/2/2/2
2	TRQ	E	57	2	-	0/4/19/21	0/2/2/2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	57	TRQ	CH2-CZ2	-8.06	1.45	1.54
2	E	57	TRQ	CZ3-CE3	7.16	1.46	1.34
2	C	57	TRQ	CH2-CZ2	-6.93	1.46	1.54
2	C	57	TRQ	CZ3-CE3	6.48	1.45	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	57	TRQ	O7-CZ2	3.91	1.31	1.23

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	57	TRQ	CZ2-CE2-NE1	6.59	130.46	119.94
2	C	57	TRQ	O6-CH2-CZ2	-6.12	114.34	118.51
2	C	57	TRQ	CZ2-CE2-NE1	5.69	129.02	119.94
2	E	57	TRQ	O6-CH2-CZ2	-5.39	114.84	118.51
2	E	57	TRQ	CB-CG-CD1	-4.65	122.22	127.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	57	TRQ	2	0
2	E	57	TRQ	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	HEC	B	600	1	32,50,50	1.41	5 (15%)	24,82,82	2.37	8 (33%)
7	EDO	D	388	-	3,3,3	1.69	0	2,2,2	2.29	1 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	374	-	3,3,3	0.52	0	2,2,2	0.26	0
9	ACT	D	389	-	3,3,3	1.32	0	3,3,3	1.93	1 (33%)
6	HEC	A	600	1	32,50,50	1.52	4 (12%)	24,82,82	3.05	11 (45%)
6	HEC	B	500	12,1	32,50,50	1.77	6 (18%)	24,82,82	2.54	8 (33%)
8	MES	D	387	-	12,12,12	1.70	1 (8%)	14,16,16	3.04	7 (50%)
6	HEC	A	500	12,1	32,50,50	1.54	8 (25%)	24,82,82	2.66	9 (37%)
10	1PE	D	390	-	15,15,15	0.46	0	14,14,14	0.66	0
11	PG4	F	387	-	12,12,12	0.37	0	11,11,11	0.55	0
7	EDO	B	374	-	3,3,3	0.74	0	2,2,2	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	HEC	B	600	1	-	2/10/54/54	-
7	EDO	D	388	-	-	0/1/1/1	-
7	EDO	A	374	-	-	1/1/1/1	-
6	HEC	A	600	1	-	2/10/54/54	-
6	HEC	B	500	12,1	-	2/10/54/54	-
8	MES	D	387	-	-	2/6/14/14	0/1/1/1
6	HEC	A	500	12,1	-	2/10/54/54	-
10	1PE	D	390	-	-	7/13/13/13	-
11	PG4	F	387	-	-	5/10/10/10	-
7	EDO	B	374	-	-	1/1/1/1	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	D	387	MES	C8-S	-5.08	1.70	1.77
6	B	500	HEC	C3A-C4A	4.84	1.53	1.42
6	A	600	HEC	C2B-C3B	-4.75	1.35	1.40
6	B	500	HEC	C3C-C4C	3.76	1.49	1.43
6	B	500	HEC	C2A-C1A	3.62	1.50	1.42

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	600	HEC	CBD-CAD-C3D	-8.79	97.62	112.62
6	A	500	HEC	CBD-CAD-C3D	-7.00	100.67	112.62
6	B	600	HEC	CBD-CAD-C3D	-6.58	101.39	112.62
8	D	387	MES	C5-N4-C3	6.39	123.22	108.83
8	D	387	MES	C6-C5-N4	-6.09	100.87	110.10

There are no chirality outliers.

5 of 24 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	D	387	MES	C8-C7-N4-C3
10	D	390	1PE	OH4-C13-C23-OH3
10	D	390	1PE	OH5-C14-C24-OH4
10	D	390	1PE	OH2-C12-C22-OH3
11	F	387	PG4	O2-C3-C4-O3


There are no ring outliers.


8 monomers are involved in 26 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	600	HEC	1	0
7	D	388	EDO	9	0
6	A	600	HEC	2	0
8	D	387	MES	1	0
6	A	500	HEC	1	0
10	D	390	1PE	5	0
11	F	387	PG4	6	0
7	B	374	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/373 (94%)	-0.58	2 (0%) 89 89	14, 23, 38, 58	0
1	B	354/373 (94%)	-0.82	1 (0%) 94 93	15, 21, 34, 56	0
2	C	125/137 (91%)	-0.36	1 (0%) 86 86	13, 15, 24, 55	0
2	E	124/137 (90%)	-0.65	1 (0%) 86 86	14, 18, 28, 56	0
3	D	376/386 (97%)	-0.62	1 (0%) 94 93	12, 17, 28, 53	0
3	F	376/386 (97%)	-0.65	1 (0%) 94 93	13, 19, 30, 54	0
All	All	1709/1792 (95%)	-0.64	7 (0%) 92 92	12, 19, 34, 58	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	132	HIS	6.2
1	A	6	ALA	5.2
3	F	11	GLN	3.2
1	A	7	ASP	3.0
3	D	11	GLN	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TRQ	C	57	16/17	0.98	0.09	14,16,21,26	0
2	TRQ	E	57	16/17	0.98	0.06	17,18,25,29	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	EDO	D	388	4/4	0.86	0.18	29,29,30,33	0
7	EDO	B	374	4/4	0.87	0.19	38,38,40,41	0
11	PG4	F	387	13/13	0.89	0.11	42,50,62,63	0
7	EDO	A	374	4/4	0.90	0.08	46,47,49,50	0
10	1PE	D	390	16/16	0.91	0.18	38,51,59,62	0
9	ACT	D	389	4/4	0.95	0.11	25,28,31,32	0
8	MES	D	387	12/12	0.96	0.15	32,52,60,62	0
5	NA	A	402	1/1	0.98	0.08	31,31,31,31	0
5	NA	B	402	1/1	0.98	0.05	27,27,27,27	0
5	NA	A	401	1/1	0.98	0.07	24,24,24,24	0
6	HEC	A	600	43/43	0.99	0.06	10,14,17,20	0
6	HEC	B	500	43/43	0.99	0.07	12,15,20,21	0
6	HEC	B	600	43/43	0.99	0.06	10,16,17,18	0
5	NA	B	401	1/1	0.99	0.04	22,22,22,22	0
6	HEC	A	500	43/43	0.99	0.06	15,19,23,24	0
4	CA	B	400	1/1	1.00	0.07	15,15,15,15	0
4	CA	A	400	1/1	1.00	0.04	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers [i](#)

There are no such residues in this entry.