



# Full wwPDB X-ray Structure Validation Report i

Sep 23, 2023 – 07:51 PM EDT

PDB ID : 5SXE  
Title : Crystal Structure of PI3Kalpha in complex with fragments 19 and 28  
Authors : Gabelli, S.B.; Vogelstein, B.; Miller, M.S.; Amzel, L.M.  
Deposited on : 2016-08-09  
Resolution : 3.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1161 (3.60-3.44)
Clashscore	141614	1244 (3.60-3.44)
Ramachandran outliers	138981	1206 (3.60-3.44)
Sidechain outliers	138945	1207 (3.60-3.44)
RSRZ outliers	127900	1080 (3.60-3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	71G	A	1101	-	-	-	X

## 2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 10518 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit alpha isoform.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	1048	8567	5468	1471	1556	2	70	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	initiating methionine	UNP P42336
A	-26	SER	-	expression tag	UNP P42336
A	-25	TYR	-	expression tag	UNP P42336
A	-24	TYR	-	expression tag	UNP P42336
A	-23	HIS	-	expression tag	UNP P42336
A	-22	HIS	-	expression tag	UNP P42336
A	-21	HIS	-	expression tag	UNP P42336
A	-20	HIS	-	expression tag	UNP P42336
A	-19	HIS	-	expression tag	UNP P42336
A	-18	HIS	-	expression tag	UNP P42336
A	-17	ASP	-	expression tag	UNP P42336
A	-16	TYR	-	expression tag	UNP P42336
A	-15	ASP	-	expression tag	UNP P42336
A	-14	ILE	-	expression tag	UNP P42336
A	-13	PRO	-	expression tag	UNP P42336
A	-12	THR	-	expression tag	UNP P42336
A	-11	THR	-	expression tag	UNP P42336
A	-10	GLU	-	expression tag	UNP P42336
A	-9	ASN	-	expression tag	UNP P42336
A	-8	LEU	-	expression tag	UNP P42336
A	-7	TYR	-	expression tag	UNP P42336
A	-6	PHE	-	expression tag	UNP P42336
A	-5	GLN	-	expression tag	UNP P42336
A	-4	GLY	-	expression tag	UNP P42336
A	-3	ALA	-	expression tag	UNP P42336
A	-2	MET	-	expression tag	UNP P42336

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P42336
A	0	SER	-	expression tag	UNP P42336

- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	B	225	Total C N O S 1936 1215 342 373 6	0	0	0

- Molecule 3 is 3-aminobenzonitrile (three-letter code: 71G) (formula: C<sub>7</sub>H<sub>6</sub>N<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N 9 7 2	0	0

- Molecule 4 is 4-bromo-1H-imidazole (three-letter code: ES3) (formula: C<sub>3</sub>H<sub>3</sub>BrN<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	Br	C	N	0	0
			6	1	3	2		


### 3 Residue-property plots [\(i\)](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



- Molecule 2: Phosphatidylinositol 3-kinase regulatory subunit alpha





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.83Å    116.26Å    148.98Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	91.65 – 3.51 48.90 – 3.51	Depositor EDS
% Data completeness (in resolution range)	99.6 (91.65-3.51) 99.7 (48.90-3.51)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.05 (at 3.48Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
$R$ , $R_{free}$	0.191 , 0.271 0.193 , 0.264	Depositor DCC
$R_{free}$ test set	1306 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	118.7	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 101.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.029 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10518	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	139.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, 71G, ES3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/8743	0.78	5/11815 (0.0%)
2	B	0.51	0/1957	0.70	0/2603
All	All	0.54	0/10700	0.76	5/14418 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	870	LEU	CA-CB-CG	6.67	130.65	115.30
1	A	113	LEU	CB-CG-CD1	5.61	120.54	111.00
1	A	513	LEU	CA-CB-CG	5.47	127.87	115.30
1	A	1	MET	CA-CB-CG	5.09	121.95	113.30
1	A	1023	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	113	LEU	Peptide
1	A	323	SER	Peptide
1	A	511	ALA	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8567	0	8543	69	0
2	B	1936	0	1913	8	0
3	A	9	0	0	0	0
4	B	6	0	3	0	0
All	All	10518	0	10459	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:962:ILE:O	1:A:965:SER:O	1.99	0.80
1:A:910:ILE:O	1:A:1025:THR:HG21	1.99	0.62
1:A:190:ILE:HD11	1:A:213:HIS:HA	1.80	0.61
1:A:261:PHE:HA	1:A:270:TYR:CE2	2.37	0.59
1:A:31:ILE:HD11	2:B:531:LEU:HD13	1.86	0.57
1:A:602:LEU:O	1:A:612:ARG:NH2	2.39	0.56
1:A:1022:ILE:HA	1:A:1025:THR:HG22	1.88	0.56
1:A:372:ASN:HB3	1:A:387:LEU:HD21	1.89	0.55
1:A:671:LEU:HB2	1:A:688:LEU:HD21	1.89	0.54
1:A:772:MET:HB2	1:A:778:PRO:HG2	1.90	0.53
1:A:353:LYS:HA	1:A:377:PRO:HB3	1.90	0.53
1:A:717:ASP:O	1:A:721:GLN:HG2	2.09	0.53
1:A:885:ASN:HB3	1:A:889:ILE:HG22	1.91	0.53
1:A:561:ILE:O	1:A:564:ILE:HG22	2.08	0.53
1:A:833:MET:HE1	1:A:904:TYR:HA	1.92	0.52
1:A:749:GLN:O	1:A:762:GLY:HA2	2.11	0.51
1:A:544:THR:HG22	2:B:380:LEU:HB3	1.92	0.50
1:A:58:LEU:HB3	1:A:61:LEU:HD13	1.93	0.50
2:B:517:ASN:HD22	2:B:518:GLU:H	1.60	0.50
1:A:360:ILE:N	1:A:360:ILE:HD12	2.28	0.48
1:A:866:LEU:HD21	1:A:870:LEU:HB2	1.95	0.48
1:A:866:LEU:HD23	1:A:866:LEU:C	2.34	0.48
1:A:866:LEU:HD21	1:A:870:LEU:HD13	1.95	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:446:TRP:CZ2	1:A:465:ASN:HA	2.48	0.48
1:A:1:MET:HB2	1:A:720:LYS:CE	2.43	0.48
1:A:61:LEU:HD11	2:B:508:TYR:CZ	2.49	0.48
1:A:367:LEU:HD12	1:A:368:CYS:O	2.13	0.48
1:A:361:TYR:HA	1:A:366:PRO:HD3	1.95	0.48
1:A:171:VAL:HG12	1:A:172:GLU:O	2.13	0.47
1:A:985:TYR:HH	1:A:1051:TRP:HE3	1.62	0.47
1:A:1023:ARG:O	1:A:1027:ALA:N	2.46	0.47
1:A:612:ARG:NH1	1:A:642:GLU:OE1	2.47	0.46
1:A:0:SER:O	1:A:1:MET:HG2	2.14	0.46
1:A:640:LYS:HE2	1:A:680:VAL:HG11	1.97	0.46
2:B:334:TYR:HH	2:B:368:TYR:HE1	1.60	0.46
1:A:517:LEU:HD21	1:A:554:HIS:HA	1.97	0.46
1:A:181:ILE:HG23	1:A:277:ILE:HG21	1.97	0.46
1:A:639:LEU:HD23	1:A:649:LEU:HD23	1.98	0.45
1:A:403:CYS:O	1:A:456:LEU:HD21	2.16	0.45
1:A:807:LEU:HD12	1:A:838:CYS:SG	2.56	0.45
1:A:735:LEU:O	1:A:739:MET:HG3	2.17	0.45
1:A:193:VAL:HG22	1:A:208:THR:HG22	1.99	0.45
1:A:543:ILE:HD11	1:A:567:LYS:HD3	1.99	0.45
1:A:147:CYS:O	1:A:151:VAL:HG23	2.17	0.44
1:A:28:ASN:ND2	1:A:62:LEU:HD21	2.32	0.44
1:A:61:LEU:HD11	2:B:508:TYR:CE2	2.53	0.44
1:A:830:ASP:OD2	1:A:832:ARG:NH2	2.51	0.43
1:A:83:PHE:HZ	1:A:112:ILE:HD12	1.83	0.43
1:A:728:GLN:NE2	1:A:771:ILE:O	2.51	0.43
1:A:856:THR:O	1:A:857:ILE:C	2.56	0.43
1:A:194:ILE:N	1:A:194:ILE:HD12	2.34	0.43
1:A:0:SER:HB2	1:A:775:ALA:HB3	1.99	0.43
1:A:537:ARG:NH1	1:A:547:GLU:OE1	2.41	0.43
1:A:642:GLU:HG2	1:A:647:ASN:CG	2.40	0.42
1:A:163:ALA:HB2	1:A:297:LEU:HD11	2.00	0.42
1:A:354:ILE:HD11	1:A:381:PRO:CB	2.49	0.42
1:A:424:TRP:CE2	1:A:446:TRP:HB2	2.54	0.42
1:A:763:ASN:HD22	1:A:763:ASN:N	2.18	0.42
1:A:31:ILE:HD12	2:B:531:LEU:HB2	2.01	0.42
1:A:253:LYS:HB3	1:A:286:MET:HB3	2.02	0.42
1:A:92:LEU:HB2	1:A:94:LEU:HD13	2.01	0.42
1:A:374:GLN:O	1:A:377:PRO:HG3	2.20	0.42
1:A:904:TYR:CE2	1:A:930:PHE:HA	2.55	0.42
1:A:897:PHE:O	1:A:901:CYS:HB2	2.20	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:908:THR:HG21	1:A:954:PHE:HB2	2.03	0.41
1:A:863:LYS:O	1:A:865:GLY:N	2.54	0.41
1:A:873:ASN:OD1	1:A:875:HIS:HB2	2.20	0.41
2:B:374:LYS:HD2	2:B:422:VAL:HG21	2.03	0.41
1:A:727:THR:O	1:A:728:GLN:C	2.58	0.41
1:A:908:THR:HG22	1:A:953:PRO:HD2	2.03	0.40
1:A:70:PHE:CD1	1:A:99:LEU:HB3	2.56	0.40
1:A:762:GLY:O	1:A:763:ASN:O	2.39	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1036/1096 (94%)	918 (89%)	101 (10%)	17 (2%)	9 45
2	B	200/279 (72%)	185 (92%)	13 (6%)	2 (1%)	15 54
All	All	1236/1375 (90%)	1103 (89%)	114 (9%)	19 (2%)	10 46

All (19) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	PRO
1	A	378	CYS
1	A	514	SER
1	A	763	ASN
1	A	107	ASN
1	A	508	TYR
1	A	972	THR
2	B	332	GLU
1	A	555	ARG
2	B	513	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	5	PRO
1	A	186	ASP
1	A	264	LYS
1	A	520	ASP
1	A	864	GLY
1	A	1051	TRP
1	A	112	ILE
1	A	471	PRO
1	A	913	ILE

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	956/997 (96%)	904 (95%)	52 (5%)	22 56
2	B	214/259 (83%)	201 (94%)	13 (6%)	18 52
All	All	1170/1256 (93%)	1105 (94%)	65 (6%)	21 55

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-15	ASP
1	A	-14	ILE
1	A	1	MET
1	A	4	ARG
1	A	23	GLU
1	A	38	ARG
1	A	39	GLU
1	A	52	GLU
1	A	99	LEU
1	A	112	ILE
1	A	113	LEU
1	A	114	ASN
1	A	190	ILE
1	A	216	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	236	SER
1	A	239	LEU
1	A	274	ARG
1	A	295	SER
1	A	299	MET
1	A	308	SER
1	A	323	SER
1	A	335	ARG
1	A	350	ASP
1	A	353	LYS
1	A	368	CYS
1	A	373	THR
1	A	380	ASN
1	A	401	ARG
1	A	420	CYS
1	A	441	MET
1	A	452	LEU
1	A	470	THR
1	A	475	LEU
1	A	499	SER
1	A	510	HIS
1	A	513	LEU
1	A	516	ARG
1	A	575	ASN
1	A	595	PRO
1	A	630	GLN
1	A	633	ILE
1	A	642	GLU
1	A	711	LYS
1	A	727	THR
1	A	740	ARG
1	A	791	GLU
1	A	808	ARG
1	A	825	GLN
1	A	870	LEU
1	A	901	CYS
1	A	919	SER
1	A	966	LYS
2	B	372	LEU
2	B	400	SER
2	B	415	GLN
2	B	416	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	441	ASN
2	B	515	GLU
2	B	517	ASN
2	B	529	ASP
2	B	531	LEU
2	B	552	GLN
2	B	585	THR
2	B	594	LEU
2	B	595	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-21	HIS
1	A	107	ASN
1	A	201	ASN
1	A	202	ASN
1	A	331	ASN
1	A	374	GLN
1	A	444	ASN
1	A	526	ASN
1	A	575	ASN
1	A	634	GLN
1	A	643	GLN
1	A	749	GLN
1	A	763	ASN
1	A	815	GLN
1	A	825	GLN
1	A	861	GLN
1	A	878	HIS
1	A	1042	GLN
2	B	517	ASN
2	B	564	ASN
2	B	595	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	SEP	A	7	1	8,9,10	0.79	0	8,12,14	1.28	1 (12%)
1	SEP	A	790	1	8,9,10	0.67	0	8,12,14	2.06	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	7	1	-	5/5/8/10	-
1	SEP	A	790	1	-	4/5/8/10	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	790	SEP	OG-CB-CA	4.22	112.25	108.14
1	A	7	SEP	P-OG-CB	2.94	126.39	118.30
1	A	790	SEP	P-OG-CB	2.34	124.75	118.30
1	A	790	SEP	O2P-P-OG	-2.21	100.84	106.73

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	7	SEP	N-CA-CB-OG
1	A	7	SEP	CA-CB-OG-P
1	A	7	SEP	CB-OG-P-O1P
1	A	7	SEP	CB-OG-P-O2P

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
1	A	7	SEP	CB-OG-P-O3P
1	A	790	SEP	N-CA-CB-OG
1	A	790	SEP	CB-OG-P-O2P
1	A	790	SEP	CB-OG-P-O3P
1	A	790	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ES3	B	701	-	3,6,6	3.60	1 (33%)	2,7,7	1.97	1 (50%)
3	71G	A	1101	-	9,9,9	2.52	2 (22%)	11,11,11	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ES3	B	701	-	-	-	0/1/1/1
3	71G	A	1101	-	-	0/2/2/2	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	71G	C3-C7	-6.65	1.29	1.44
4	B	701	ES3	BR01-C02	6.23	2.00	1.90
3	A	1101	71G	C1-N1	3.08	1.49	1.38

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	701	ES3	C03-N04-C05	2.46	109.61	105.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1046/1096 (95%)	0.10	40 (3%) 40 30	73, 123, 193, 254	0
2	B	225/279 (80%)	0.58	23 (10%) 6 6	111, 189, 231, 275	0
All	All	1271/1375 (92%)	0.19	63 (4%) 28 21	73, 130, 212, 275	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	409	ARG	5.0
1	A	378	CYS	4.8
1	A	346	VAL	4.5
2	B	328	LEU	4.2
2	B	390	TYR	4.2
1	A	423	ALA	3.8
1	A	-26	SER	3.8
1	A	343	TYR	3.7
1	A	872	PHE	3.5
1	A	344	VAL	3.5
1	A	347	ASN	3.5
1	A	869	ALA	3.5
2	B	346	LYS	3.5
1	A	871	GLN	3.4
1	A	351	ILE	3.3
2	B	408	TYR	3.3
1	A	1	MET	3.2
1	A	354	ILE	3.1
1	A	523	LEU	3.0
1	A	341	ALA	3.0
1	A	61	LEU	3.0
2	B	442	ILE	2.9
1	A	-14	ILE	2.9
1	A	810	ASP	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	443	GLU	2.8
2	B	505	SER	2.7
1	A	422	LEU	2.7
1	A	1054	LYS	2.7
1	A	353	LYS	2.6
2	B	399	SER	2.5
1	A	381	PRO	2.5
2	B	401	VAL	2.5
1	A	374	GLN	2.5
1	A	338	ILE	2.5
2	B	329	GLN	2.5
1	A	1009	GLY	2.4
2	B	405	ILE	2.4
1	A	345	ASN	2.4
2	B	397	THR	2.4
2	B	504	TYR	2.4
1	A	870	LEU	2.3
1	A	-13	PRO	2.3
1	A	473	LEU	2.3
1	A	83	PHE	2.3
1	A	358	THR	2.3
2	B	333	TRP	2.3
2	B	508	TYR	2.2
2	B	356	LEU	2.2
2	B	444	ALA	2.2
2	B	510	GLU	2.2
1	A	809	GLN	2.1
1	A	806	ASP	2.1
1	A	350	ASP	2.1
1	A	868	GLY	2.1
1	A	933	ASP	2.1
2	B	331	ALA	2.1
1	A	673	SER	2.1
2	B	368	TYR	2.1
2	B	580	TYR	2.1
1	A	308	SER	2.0
1	A	973	LYS	2.0
2	B	586	GLN	2.0
1	A	364	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	SEP	A	7	10/11	0.64	0.29	163,190,249,258	0
1	SEP	A	790	10/11	0.94	0.13	94,118,199,203	0

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	71G	A	1101	9/9	0.57	1.35	146,183,195,196	0
4	ES3	B	701	6/6	0.65	0.28	191,195,207,240	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.