



## wwPDB EM Validation Summary Report ⓘ

Apr 16, 2026 – 10:42 am BST

PDB ID : 9T0Z / pdb\_00009t0z  
EMDB ID : EMD-55416  
Title : HPFcold Bound Hibernating C. burnetii 30S Ribosome  
Authors : Stuart, W.S.; Isupov, M.N.; Harmer, N.J.  
Deposited on : 2025-10-20  
Resolution : 2.87 Å(reported)  
Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	12062 ( 2.37 - 3.37 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	M	119	 8% 95% . .
2	N	99	 64% 26% 8% . .
3	9	183	 22% 91% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	S	95	
5	A	1535	
6	O	89	
7	C	227	
8	I	139	
9	U	74	
10	F	127	
11	T	90	
12	D	206	
13	E	168	
14	H	130	
15	P	137	
16	L	124	
17	Q	89	
18	4	79	
19	G	191	
20	J	110	
21	K	123	
22	R	73	

## 2 Entry composition

There are 26 unique types of molecules in this entry. The entry contains 52223 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Small ribosomal subunit protein uS13.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	M	115	Total	C	N	O	S	0	0
			907	557	187	160	3		

- Molecule 2 is a protein called Small ribosomal subunit protein uS14.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	N	98	Total	C	N	O	S	0	0
			800	496	167	132	5		

- Molecule 3 is a protein called Ribosome hibernation promoting factor.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	9	175	Total	C	N	O	S	0	0
			1412	892	253	262	5		

- Molecule 4 is a protein called Small ribosomal subunit protein uS19.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	S	84	Total	C	N	O	S	0	0
			668	419	131	114	4		

- Molecule 5 is a RNA chain called 16S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	A	1535	Total	C	N	O	P	0	0
			32928	14693	6016	10684	1535		

- Molecule 6 is a protein called Small ribosomal subunit protein uS15.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	O	88	Total	C	N	O	S	0	0
			716	448	141	125	2		

- Molecule 7 is a protein called Small ribosomal subunit protein uS3.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	C	206	Total	C	N	O	S	0	0
			1641	1037	314	286	4		

- Molecule 8 is a protein called Small ribosomal subunit protein uS9.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	I	125	Total	C	N	O	S	0	0
			981	607	197	175	2		

- Molecule 9 is a protein called Small ribosomal subunit protein bS21.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	U	68	Total	C	N	O	S	0	0
			565	354	111	97	3		

- Molecule 10 is a protein called Small ribosomal subunit protein bS6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	F	106	Total	C	N	O	S	0	0
			867	545	159	158	5		

- Molecule 11 is a protein called Small ribosomal subunit protein bS20.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	T	86	Total	C	N	O		0	0
			669	405	144	120			

- Molecule 12 is a protein called Small ribosomal subunit protein uS4.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	D	205	Total	C	N	O	S	0	0
			1659	1045	309	298	7		

- Molecule 13 is a protein called Small ribosomal subunit protein uS5.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	E	158	Total	C	N	O	S	0	0
			1153	723	213	211	6		

- Molecule 14 is a protein called Small ribosomal subunit protein uS8.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	H	130	Total	C	N	O	S	0	0
			1017	634	182	194	7		

- Molecule 15 is a protein called Small ribosomal subunit protein bS16.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	P	106	Total	C	N	O	S	0	0
			865	542	174	146	3		

- Molecule 16 is a protein called Small ribosomal subunit protein uS12.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	L	123	Total	C	N	O	S	0	0
			958	586	204	166	2		

- Molecule 17 is a protein called Small ribosomal subunit protein uS17.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	83	Total	C	N	O	S	0	0
			669	415	131	119	4		

- Molecule 18 is a protein called Large ribosomal subunit protein bL31.

Mol	Chain	Residues	Atoms				AltConf	Trace
18	4	10	Total	C	N	O	0	0
			93	61	19	13		

- Molecule 19 is a protein called Small ribosomal subunit protein uS7.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	G	162	Total	C	N	O	S	0	0
			1291	808	254	224	5		

- Molecule 20 is a protein called Small ribosomal subunit protein uS10.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	J	102	Total	C	N	O	S	0	0
			828	517	157	153	1		

- Molecule 21 is a protein called Small ribosomal subunit protein uS11.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	K	119	Total	C	N	O	S	0	0
			915	558	190	162	5		

- Molecule 22 is a protein called Small ribosomal subunit protein bS18.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	R	65	Total	C	N	O	S	0	0
			524	331	94	95	4		

- Molecule 23 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
23	N	1	Total	Zn	0
			1	1	
23	R	1	Total	Zn	0
			1	1	

- Molecule 24 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		AltConf
24	A	34	Total	K	0
			34	34	

- Molecule 25 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
25	A	53	Total	Mg	0
			53	53	


- Molecule 26 is water.

Mol	Chain	Residues	Atoms		AltConf
26	A	7	Total	O	0
			7	7	
26	K	1	Total	O	0
			1	1	


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Small ribosomal subunit protein uS13




- Molecule 2: Small ribosomal subunit protein uS14




- Molecule 3: Ribosome hibernation promoting factor




- Molecule 4: Small ribosomal subunit protein uS19



- Molecule 5: 16S ribosomal RNA









- Molecule 6: Small ribosomal subunit protein uS15

Chain O: 97%





- Molecule 7: Small ribosomal subunit protein uS3

Chain C: 90% 7% 9%





- Molecule 8: Small ribosomal subunit protein uS9

Chain I:  88% 10%





- Molecule 9: Small ribosomal subunit protein bS21

Chain U:  18% 89% 8%




- Molecule 10: Small ribosomal subunit protein bS6

Chain F:  5% 82% 17%



- Molecule 11: Small ribosomal subunit protein bS20

Chain T:  94% 5% 1%




- Molecule 12: Small ribosomal subunit protein uS4

Chain D:  97% 2% 1%




- Molecule 13: Small ribosomal subunit protein uS5

Chain E:  93% 6% 1%





- Molecule 14: Small ribosomal subunit protein uS8

Chain H:  99% 1% 0%




- Molecule 15: Small ribosomal subunit protein bS16

Chain P:  76% 23%




- Molecule 16: Small ribosomal subunit protein uS12

Chain L:  99%





- Molecule 17: Small ribosomal subunit protein uS17

Chain Q:  93% 7%






- Molecule 18: Large ribosomal subunit protein bL31

Chain 4:  5% 11% 87%





- Molecule 19: Small ribosomal subunit protein uS7

Chain G:  9% 81% 15%




- Molecule 20: Small ribosomal subunit protein uS10



Chain J:  18% 81% 12% 7%



- Molecule 21: Small ribosomal subunit protein uS11



- Molecule 22: Small ribosomal subunit protein bS18



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	24143	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	45	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.322	Depositor
Minimum map value	-0.074	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.016	Depositor
Recommended contour level	0.07	Depositor
Map size (Å)	423.99997, 423.99997, 423.99997	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K, 5MC, ZN, 4OC, G7M, MA6, UR3, 2MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	M	0.52	0/915	1.19	0/1218
2	N	0.46	0/809	1.09	1/1069 (0.1%)
3	9	0.52	0/1440	1.02	0/1943
4	S	0.53	0/684	0.97	0/918
5	A	0.38	2/36657 (0.0%)	0.81	47/57176 (0.1%)
6	O	0.46	0/724	1.07	0/964
7	C	0.50	0/1666	1.02	1/2239 (0.0%)
8	I	0.51	0/994	1.09	0/1330
9	U	0.48	0/572	1.15	0/757
10	F	0.51	0/882	0.97	0/1185
11	T	0.47	0/673	1.17	0/896
12	D	0.47	0/1684	1.03	0/2258
13	E	0.52	0/1166	1.00	0/1569
14	H	0.51	0/1026	0.97	0/1372
15	P	0.49	0/879	1.03	0/1169
16	L	0.51	0/971	0.97	0/1299
17	Q	0.50	0/677	0.88	0/904
18	4	0.46	0/94	1.24	0/121
19	G	0.50	0/1308	1.11	0/1752
20	J	0.53	0/836	1.08	0/1128
21	K	0.52	0/929	1.05	0/1240
22	R	0.49	0/533	1.08	0/718
All	All	0.43	2/56119 (0.0%)	0.89	49/83225 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1495	UR3	O3'-P	6.02	1.62	1.56
5	A	524	G7M	O3'-P	5.21	1.61	1.56

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	554	G	C2'-C3'-O3'	8.92	127.08	113.70
2	N	15	ARG	CG-CD-NE	8.87	131.51	112.00
5	A	268	C	C2'-C3'-O3'	7.71	125.26	113.70
5	A	1314	C	C2'-C3'-O3'	7.65	125.18	113.70
5	A	1532	C	C4'-C3'-O3'	7.33	123.99	113.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	907	0	966	2	0
2	N	800	0	876	27	0
3	9	1412	0	1404	14	0
4	S	668	0	681	13	0
5	A	32928	0	16586	129	0
6	O	716	0	766	1	0
7	C	1641	0	1721	1	0
8	I	981	0	1004	0	0
9	U	565	0	602	1	0
10	F	867	0	878	0	0
11	T	669	0	715	0	0
12	D	1659	0	1726	5	0
13	E	1153	0	1213	1	0
14	H	1017	0	1073	1	0
15	P	865	0	908	2	0
16	L	958	0	1016	0	0
17	Q	669	0	708	0	0
18	4	93	0	102	1	0
19	G	1291	0	1351	10	0
20	J	828	0	875	16	0
21	K	915	0	935	4	0
22	R	524	0	528	0	0
23	N	1	0	0	0	0
23	R	1	0	0	0	0
24	A	34	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
25	A	53	0	0	0	0
26	A	7	0	0	0	0
26	K	1	0	0	0	0
All	All	52223	0	36634	167	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:9:61:ASN:OD1	5:A:962:2MG:H1'	1.41	1.18
5:A:714:C:H4'	21:K:113:ASN:HD22	1.22	1.02
5:A:714:C:C4'	21:K:113:ASN:HD22	1.85	0.90
5:A:198:G:H21	5:A:466:U:H1'	1.35	0.87
5:A:714:C:H4'	21:K:113:ASN:ND2	1.89	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	113/119 (95%)	109 (96%)	4 (4%)	0	100	100
2	N	96/99 (97%)	89 (93%)	7 (7%)	0	100	100
3	9	173/183 (94%)	167 (96%)	6 (4%)	0	100	100
4	S	82/95 (86%)	80 (98%)	2 (2%)	0	100	100
6	O	86/89 (97%)	84 (98%)	2 (2%)	0	100	100
7	C	204/227 (90%)	199 (98%)	5 (2%)	0	100	100
8	I	121/139 (87%)	118 (98%)	3 (2%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
9	U	66/74 (89%)	66 (100%)	0	0	100	100
10	F	104/127 (82%)	104 (100%)	0	0	100	100
11	T	84/90 (93%)	84 (100%)	0	0	100	100
12	D	203/206 (98%)	199 (98%)	4 (2%)	0	100	100
13	E	156/168 (93%)	153 (98%)	3 (2%)	0	100	100
14	H	128/130 (98%)	126 (98%)	2 (2%)	0	100	100
15	P	104/137 (76%)	102 (98%)	2 (2%)	0	100	100
16	L	121/124 (98%)	118 (98%)	3 (2%)	0	100	100
17	Q	81/89 (91%)	79 (98%)	2 (2%)	0	100	100
18	4	8/79 (10%)	7 (88%)	1 (12%)	0	100	100
19	G	158/191 (83%)	158 (100%)	0	0	100	100
20	J	100/110 (91%)	96 (96%)	4 (4%)	0	100	100
21	K	117/123 (95%)	115 (98%)	2 (2%)	0	100	100
22	R	63/73 (86%)	63 (100%)	0	0	100	100
All	All	2368/2672 (89%)	2316 (98%)	52 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	93/96 (97%)	93 (100%)	0	100	100
2	N	86/87 (99%)	71 (83%)	15 (17%)	2	6
3	9	153/161 (95%)	153 (100%)	0	100	100
4	S	73/82 (89%)	72 (99%)	1 (1%)	59	83
6	O	78/79 (99%)	78 (100%)	0	100	100
7	C	169/186 (91%)	169 (100%)	0	100	100
8	I	101/108 (94%)	99 (98%)	2 (2%)	48	76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
9	U	58/64 (91%)	57 (98%)	1 (2%)	53	80
10	F	95/110 (86%)	93 (98%)	2 (2%)	47	75
11	T	69/73 (94%)	68 (99%)	1 (1%)	59	83
12	D	177/178 (99%)	176 (99%)	1 (1%)	78	92
13	E	119/126 (94%)	119 (100%)	0	100	100
14	H	114/114 (100%)	114 (100%)	0	100	100
15	P	88/110 (80%)	88 (100%)	0	100	100
16	L	100/101 (99%)	100 (100%)	0	100	100
17	Q	76/82 (93%)	76 (100%)	0	100	100
18	4	9/69 (13%)	9 (100%)	0	100	100
19	G	131/150 (87%)	130 (99%)	1 (1%)	73	89
20	J	92/98 (94%)	90 (98%)	2 (2%)	45	75
21	K	94/97 (97%)	94 (100%)	0	100	100
22	R	57/65 (88%)	56 (98%)	1 (2%)	51	78
All	All	2032/2236 (91%)	2005 (99%)	27 (1%)	59	84

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	96	LYS
8	I	125	VAL
20	J	58	ILE
8	I	70	VAL
9	U	33	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
16	L	72	HIS
20	J	105	ASN
20	J	57	ASN
21	K	70	ASN
4	S	57	HIS

### 5.3.3 RNA

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	A	1534/1535 (99%)	228 (14%)	46 (2%)

5 of 228 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	A	4	U
5	A	8	G
5	A	21	G
5	A	31	A
5	A	38	G

5 of 46 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
5	A	904	A
5	A	1314	C
5	A	972	A
5	A	1125	C
5	A	1334	G

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MA6	A	1515	5	23,26,27	1.55	5 (21%)	34,38,41	2.19	12 (35%)
5	UR3	A	1495	5	19,22,23	1.19	1 (5%)	26,32,35	1.78	5 (19%)
5	2MG	A	1513	5	23,26,27	1.24	3 (13%)	32,38,41	2.38	10 (31%)
5	MA6	A	1516	5	23,26,27	1.67	5 (21%)	34,38,41	2.24	12 (35%)
5	5MC	A	963	5	18,22,23	0.99	1 (5%)	26,32,35	1.32	4 (15%)
5	4OC	A	1399	5	20,23,24	0.82	0	26,32,35	1.21	4 (15%)
5	2MG	A	962	5	23,26,27	1.24	4 (17%)	32,38,41	2.32	11 (34%)
5	G7M	A	524	5	23,26,27	2.36	5 (21%)	35,39,42	2.99	10 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MA6	A	1515	5	-	3/11/29/30	0/3/3/3
5	UR3	A	1495	5	-	0/7/25/26	0/2/2/2
5	2MG	A	1513	5	-	2/9/27/28	0/3/3/3
5	MA6	A	1516	5	-	6/11/29/30	0/3/3/3
5	5MC	A	963	5	-	2/7/25/26	0/2/2/2
5	4OC	A	1399	5	-	0/9/29/30	0/2/2/2
5	2MG	A	962	5	-	2/9/27/28	0/3/3/3
5	G7M	A	524	5	-	3/7/25/26	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	524	G7M	C8-N7	7.29	1.46	1.33
5	A	524	G7M	C5-N7	-4.98	1.33	1.39
5	A	1516	MA6	C5-C4	4.84	1.48	1.39
5	A	1515	MA6	C5-C4	4.69	1.47	1.39
5	A	524	G7M	C8-N9	4.00	1.46	1.35

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1513	2MG	C2-N3-C4	7.86	121.78	112.04
5	A	524	G7M	CN7-N7-C8	-7.74	112.89	124.84
5	A	962	2MG	C2-N3-C4	7.42	121.24	112.04
5	A	524	G7M	C8-N7-C5	6.73	116.19	107.78
5	A	524	G7M	N9-C8-N7	-6.56	95.98	112.21

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	962	2MG	N3-C2-N2-CM2
5	A	963	5MC	O4'-C4'-C5'-O5'
5	A	1513	2MG	N1-C2-N2-CM2
5	A	1513	2MG	N3-C2-N2-CM2
5	A	1515	MA6	C5-C6-N6-C10

There are no ring outliers.

7 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1515	MA6	2	0
5	A	1495	UR3	4	0
5	A	1513	2MG	1	0
5	A	1516	MA6	7	0
5	A	1399	4OC	2	0
5	A	962	2MG	2	0
5	A	524	G7M	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 89 ligands modelled in this entry, 89 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-55416. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.




## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


### 6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 212




Y Index: 184




Z Index: 210


### 6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.07. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1013 nm<sup>3</sup>; this corresponds to an approximate mass of 916 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.348 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.348 \text{ \AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.87	-	-
Author-provided FSC curve	2.87	3.24	2.92
Unmasked-calculated*	4.75	9.02	5.13

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.75 differs from the reported value 2.87 by more than 10 %

## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-55416 and PDB model 9T0Z. Per-residue inclusion information can be found in section [3](#) on page [8](#).


### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.07 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.07).















































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.07) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8620	 0.5160
4	 0.3750	 0.3420
9	 0.5850	 0.4590
A	 0.9410	 0.5300
C	 0.7000	 0.4790
D	 0.7950	 0.5110
E	 0.8170	 0.5360
F	 0.7200	 0.4680
G	 0.6360	 0.4630
H	 0.8080	 0.5260
I	 0.7410	 0.4820
J	 0.6190	 0.4340
K	 0.7750	 0.5060
L	 0.7820	 0.5370
M	 0.7020	 0.4550
N	 0.7440	 0.4860
O	 0.8070	 0.5220
P	 0.8480	 0.5300
Q	 0.7570	 0.5270
R	 0.7770	 0.5010
S	 0.6790	 0.4730
T	 0.7920	 0.4990
U	 0.5990	 0.4530

