



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 17, 2024 – 05:56 PM EST

PDB ID : 3T5Q
Title : 3A structure of Lassa virus nucleoprotein in complex with ssRNA
Authors : Hastie, K.M.; Liu, T.; King, L.B.; Ngo, N.; Zandonatti, M.A.; Woods, V.L.;
de la Torre, J.C.; Saphire, E.O.
Deposited on : 2011-07-27
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)
RNA backbone	3102	1173 (3.30-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	71% 14% • 13%
1	B	353	71% 12% • 15%
1	E	353	70% 14% • 15%
1	G	353	2% 71% 13% • 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	353	<p>% 68% 16% 14%</p>
1	K	353	<p>8% 66% 16% 17%</p>
2	C	8	<p>50% 25% 12% 12%</p>
2	F	8	<p>25% 62% 12%</p>
3	D	7	<p>29% 43% 29%</p>
4	H	6	<p>17% 67% 17%</p>
4	L	6	<p>50% 33% 17%</p>
5	J	8	<p>12% 12% 38% 38% 12%</p>

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 14684 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	306	2352	1470	417	453	12	0	0	0
1	B	301	2284	1428	399	446	11	0	0	0
1	E	301	2300	1441	403	445	11	0	0	0
1	G	303	2310	1446	404	447	13	0	0	0
1	I	302	2311	1447	403	449	12	0	0	0
1	K	294	2235	1402	385	436	12	0	0	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	expression tag	UNP Q5S585
A	-11	ALA	-	expression tag	UNP Q5S585
A	-10	HIS	-	expression tag	UNP Q5S585
A	-9	HIS	-	expression tag	UNP Q5S585
A	-8	HIS	-	expression tag	UNP Q5S585
A	-7	HIS	-	expression tag	UNP Q5S585
A	-6	HIS	-	expression tag	UNP Q5S585
A	-5	HIS	-	expression tag	UNP Q5S585
A	-4	ASP	-	expression tag	UNP Q5S585
A	-3	ASP	-	expression tag	UNP Q5S585
A	-2	ASP	-	expression tag	UNP Q5S585
A	-1	LYS	-	expression tag	UNP Q5S585
A	0	MET	-	expression tag	UNP Q5S585
B	-12	MET	-	expression tag	UNP Q5S585
B	-11	ALA	-	expression tag	UNP Q5S585
B	-10	HIS	-	expression tag	UNP Q5S585
B	-9	HIS	-	expression tag	UNP Q5S585

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-8	HIS	-	expression tag	UNP Q5S585
B	-7	HIS	-	expression tag	UNP Q5S585
B	-6	HIS	-	expression tag	UNP Q5S585
B	-5	HIS	-	expression tag	UNP Q5S585
B	-4	ASP	-	expression tag	UNP Q5S585
B	-3	ASP	-	expression tag	UNP Q5S585
B	-2	ASP	-	expression tag	UNP Q5S585
B	-1	LYS	-	expression tag	UNP Q5S585
B	0	MET	-	expression tag	UNP Q5S585
E	-12	MET	-	expression tag	UNP Q5S585
E	-11	ALA	-	expression tag	UNP Q5S585
E	-10	HIS	-	expression tag	UNP Q5S585
E	-9	HIS	-	expression tag	UNP Q5S585
E	-8	HIS	-	expression tag	UNP Q5S585
E	-7	HIS	-	expression tag	UNP Q5S585
E	-6	HIS	-	expression tag	UNP Q5S585
E	-5	HIS	-	expression tag	UNP Q5S585
E	-4	ASP	-	expression tag	UNP Q5S585
E	-3	ASP	-	expression tag	UNP Q5S585
E	-2	ASP	-	expression tag	UNP Q5S585
E	-1	LYS	-	expression tag	UNP Q5S585
E	0	MET	-	expression tag	UNP Q5S585
G	-12	MET	-	expression tag	UNP Q5S585
G	-11	ALA	-	expression tag	UNP Q5S585
G	-10	HIS	-	expression tag	UNP Q5S585
G	-9	HIS	-	expression tag	UNP Q5S585
G	-8	HIS	-	expression tag	UNP Q5S585
G	-7	HIS	-	expression tag	UNP Q5S585
G	-6	HIS	-	expression tag	UNP Q5S585
G	-5	HIS	-	expression tag	UNP Q5S585
G	-4	ASP	-	expression tag	UNP Q5S585
G	-3	ASP	-	expression tag	UNP Q5S585
G	-2	ASP	-	expression tag	UNP Q5S585
G	-1	LYS	-	expression tag	UNP Q5S585
G	0	MET	-	expression tag	UNP Q5S585
I	-12	MET	-	expression tag	UNP Q5S585
I	-11	ALA	-	expression tag	UNP Q5S585
I	-10	HIS	-	expression tag	UNP Q5S585
I	-9	HIS	-	expression tag	UNP Q5S585
I	-8	HIS	-	expression tag	UNP Q5S585
I	-7	HIS	-	expression tag	UNP Q5S585
I	-6	HIS	-	expression tag	UNP Q5S585

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	-5	HIS	-	expression tag	UNP Q5S585
I	-4	ASP	-	expression tag	UNP Q5S585
I	-3	ASP	-	expression tag	UNP Q5S585
I	-2	ASP	-	expression tag	UNP Q5S585
I	-1	LYS	-	expression tag	UNP Q5S585
I	0	MET	-	expression tag	UNP Q5S585
K	-12	MET	-	expression tag	UNP Q5S585
K	-11	ALA	-	expression tag	UNP Q5S585
K	-10	HIS	-	expression tag	UNP Q5S585
K	-9	HIS	-	expression tag	UNP Q5S585
K	-8	HIS	-	expression tag	UNP Q5S585
K	-7	HIS	-	expression tag	UNP Q5S585
K	-6	HIS	-	expression tag	UNP Q5S585
K	-5	HIS	-	expression tag	UNP Q5S585
K	-4	ASP	-	expression tag	UNP Q5S585
K	-3	ASP	-	expression tag	UNP Q5S585
K	-2	ASP	-	expression tag	UNP Q5S585
K	-1	LYS	-	expression tag	UNP Q5S585
K	0	MET	-	expression tag	UNP Q5S585

- Molecule 2 is a RNA chain called RNA (5'-R(P*UP*UP*AP*UP*CP*UP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	P	0	0	0
			164	74	24	58	8			
2	F	8	Total	C	N	O	P	0	0	0
			164	74	24	58	8			

- Molecule 3 is a RNA chain called RNA (5'-R(P*UP*AP*UP*CP*UP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	7	Total	C	N	O	P	0	0	0
			144	65	22	50	7			


- Molecule 4 is a RNA chain called RNA (5'-R(P*UP*AP*UP*CP*UP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	6	Total	C	N	O	P	0	0	0
			122	55	17	44	6			
4	L	6	Total	C	N	O	P	0	0	0
			122	55	17	44	6			

- Molecule 5 is a RNA chain called RNA (5'-R(P*UP*UP*AP*UP*CP*UP*CP*C)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
5	J	8	162	73	22	59	8	0	0	0

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
6	A	1	5	4	1	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total	O	0	0
			3	3		
7	B	3	Total	O	0	0
			3	3		
7	J	1	Total	O	0	0
			1	1		
7	K	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nucleoprotein

Chain A: 




- Molecule 1: Nucleoprotein


Chain B: 





- Molecule 1: Nucleoprotein

Chain E: 










- Molecule 1: Nucleoprotein



- Molecule 1: Nucleoprotein



- Molecule 1: Nucleoprotein




- Molecule 2: RNA (5'-R(P*UP*UP*AP*UP*CP*UP*CP*A)-3')

Chain C:  50% 25% 12% 12%



• Molecule 2: RNA (5'-R(P*UP*UP*AP*UP*CP*UP*CP*A)-3')

Chain F:  25% 62% 12%




• Molecule 3: RNA (5'-R(P*UP*AP*UP*CP*UP*CP*A)-3')

Chain D:  29% 43% 29%




• Molecule 4: RNA (5'-R(P*UP*AP*UP*CP*UP*C)-3')

Chain H:  17% 67% 17%



• Molecule 4: RNA (5'-R(P*UP*AP*UP*CP*UP*C)-3')

Chain L:  50% 33% 17%



• Molecule 5: RNA (5'-R(P*UP*UP*AP*UP*CP*UP*CP*C)-3')

Chain J:  12% 12% 38% 38% 12%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	127.44Å 127.44Å 298.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.77 – 3.00 39.77 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.77-3.00) 99.9 (39.77-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 3.01Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.226 , 0.258 0.222 , 0.254	Depositor DCC
R_{free} test set	2771 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	90.3	Xtrriage
Anisotropy	0.019	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 67.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.021 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14684	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2378	0.69	4/3210 (0.1%)
1	B	0.52	0/2310	0.69	3/3125 (0.1%)
1	E	0.53	2/2326 (0.1%)	0.71	5/3142 (0.2%)
1	G	0.48	0/2336	0.72	5/3161 (0.2%)
1	I	0.47	0/2337	0.69	3/3161 (0.1%)
1	K	0.54	0/2259	0.60	0/3054
2	C	1.27	2/181 (1.1%)	2.44	14/278 (5.0%)
2	F	0.63	0/181	1.21	0/278
3	D	0.68	0/159	1.25	0/244
4	H	0.59	0/134	1.04	1/205 (0.5%)
4	L	0.76	0/134	1.20	0/205
5	J	0.84	0/178	1.37	2/273 (0.7%)
All	All	0.55	4/14913 (0.0%)	0.78	37/20336 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	76	ASN	CG-OD1	-8.88	1.04	1.24
1	E	76	ASN	CG-ND2	-6.14	1.17	1.32
2	C	8	A	C8-N7	5.91	1.35	1.31
2	C	8	A	C6-N1	-5.10	1.31	1.35

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	8	A	N9-C4-C5	16.12	112.25	105.80
1	E	52	ARG	NE-CZ-NH2	-15.58	112.51	120.30
1	B	52	ARG	NE-CZ-NH1	-14.95	112.82	120.30
1	I	52	ARG	NE-CZ-NH2	-14.91	112.85	120.30
1	G	52	ARG	NE-CZ-NH1	-14.53	113.03	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2352	0	2383	56	1
1	B	2284	0	2272	35	0
1	E	2300	0	2323	52	0
1	G	2310	0	2319	34	1
1	I	2311	0	2329	50	0
1	K	2235	0	2265	59	0
2	C	164	0	85	3	0
2	F	164	0	85	7	0
3	D	144	0	75	10	0
4	H	122	0	64	3	0
4	L	122	0	64	8	0
5	J	162	0	85	11	0
6	A	5	0	0	0	0
7	A	3	0	0	0	0
7	B	3	0	0	1	0
7	J	1	0	0	0	0
7	K	2	0	0	0	0
All	All	14684	0	14349	281	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 281 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:66:ARG:HH11	1:K:66:ARG:HG2	1.09	1.14
1:I:135:ASP:HB3	1:I:138:ARG:HH12	1.14	1.12
1:A:66:ARG:HH11	1:A:66:ARG:HG2	1.10	1.12
1:B:66:ARG:HG2	1:B:66:ARG:HH11	1.14	1.11
1:I:66:ARG:HG2	1:I:66:ARG:HH11	1.13	1.10

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:340:SER:OG	1:G:62:ASN:OD1[6_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	300/353 (85%)	289 (96%)	10 (3%)	1 (0%)	41 76
1	B	295/353 (84%)	285 (97%)	9 (3%)	1 (0%)	41 76
1	E	295/353 (84%)	283 (96%)	12 (4%)	0	100 100
1	G	297/353 (84%)	282 (95%)	15 (5%)	0	100 100
1	I	296/353 (84%)	283 (96%)	11 (4%)	2 (1%)	22 60
1	K	288/353 (82%)	276 (96%)	10 (4%)	2 (1%)	22 60
All	All	1771/2118 (84%)	1698 (96%)	67 (4%)	6 (0%)	41 76

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	130	SER
1	K	131	SER
1	B	135	ASP
1	K	138	ARG
1	I	163	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/305 (85%)	250 (96%)	9 (4%)	36	71
1	B	247/305 (81%)	236 (96%)	11 (4%)	27	64
1	E	252/305 (83%)	246 (98%)	6 (2%)	49	79
1	G	253/305 (83%)	245 (97%)	8 (3%)	39	74
1	I	255/305 (84%)	244 (96%)	11 (4%)	29	66
1	K	248/305 (81%)	241 (97%)	7 (3%)	43	77
All	All	1514/1830 (83%)	1462 (97%)	52 (3%)	37	72

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	95	THR
1	I	66	ARG
1	K	132	GLN
1	G	134	LEU
1	G	289	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	240	ASN
1	K	196	ASN
1	I	49	ASN
1	K	240	ASN
1	K	49	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	7/8 (87%)	2 (28%)	0
2	F	7/8 (87%)	1 (14%)	0
3	D	6/7 (85%)	2 (33%)	0
4	H	5/6 (83%)	1 (20%)	0
4	L	5/6 (83%)	1 (20%)	0
5	J	7/8 (87%)	4 (57%)	0
All	All	37/43 (86%)	11 (29%)	0

5 of 11 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	U
2	C	8	A
3	D	4	U
3	D	5	C
2	F	7	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PO4	A	341	-	4,4,4	0.94	0	6,6,6	0.50	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/353 (86%)	-0.12	1 (0%) 94 84	54, 75, 108, 140	0
1	B	301/353 (85%)	-0.18	0 100 100	58, 78, 121, 154	0
1	E	301/353 (85%)	-0.20	1 (0%) 94 84	53, 77, 107, 140	0
1	G	303/353 (85%)	-0.05	8 (2%) 56 27	58, 82, 113, 142	0
1	I	302/353 (85%)	-0.10	5 (1%) 70 41	52, 81, 114, 144	0
1	K	294/353 (83%)	0.31	29 (9%) 7 2	61, 85, 115, 143	0
2	C	8/8 (100%)	0.38	0 100 100	51, 74, 120, 124	0
2	F	8/8 (100%)	-0.09	0 100 100	72, 95, 126, 127	0
3	D	7/7 (100%)	0.11	0 100 100	68, 85, 113, 121	0
4	H	6/6 (100%)	0.12	0 100 100	80, 86, 130, 148	0
4	L	6/6 (100%)	0.15	0 100 100	107, 115, 131, 146	0
5	J	8/8 (100%)	0.43	1 (12%) 3 1	75, 98, 133, 140	0
All	All	1850/2161 (85%)	-0.05	45 (2%) 59 30	51, 80, 116, 154	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	292	PHE	3.8
1	K	334	THR	3.6
1	K	298	GLY	3.5
1	K	56	LYS	3.4
1	I	297	PRO	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	PO4	A	341	5/5	0.98	0.25	73,91,99,105	0

6.5 Other polymers [i](#)

There are no such residues in this entry.