



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2023 – 05:51 PM EDT

PDB ID : 5T5W  
Title : Structure of an affinity matured lambda-IFN/IFN-lambdaR1/IL-10Rbeta receptor complex  
Authors : Mendoza, J.L.; Jude, K.M.; Garcia, K.C.  
Deposited on : 2016-08-31  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 4% 83% 9% 7%
2	B	218	 11% 73% 13% 14%
3	C	176	 2% 81% 5% 14%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-10 receptor subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	198	1608	1028	263	306	11	0	0	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	GLY	-	expression tag	UNP Q08334
A	19	SER	-	engineered mutation	UNP Q08334
A	49	GLN	ASN	engineered mutation	UNP Q08334
A	68	GLN	ASN	engineered mutation	UNP Q08334
A	102	GLN	ASN	engineered mutation	UNP Q08334
A	161	GLN	ASN	engineered mutation	UNP Q08334
A	221	ALA	-	expression tag	UNP Q08334
A	222	ALA	-	expression tag	UNP Q08334
A	223	ALA	-	expression tag	UNP Q08334
A	224	HIS	-	expression tag	UNP Q08334
A	225	HIS	-	expression tag	UNP Q08334
A	226	HIS	-	expression tag	UNP Q08334
A	227	HIS	-	expression tag	UNP Q08334
A	228	HIS	-	expression tag	UNP Q08334
A	229	HIS	-	expression tag	UNP Q08334
A	230	HIS	-	expression tag	UNP Q08334
A	231	HIS	-	expression tag	UNP Q08334

- Molecule 2 is a protein called Interferon lambda receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	188	1437	938	226	264	9	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	THR	-	expression tag	UNP Q8IU57
B	-1	SER	-	expression tag	UNP Q8IU57
B	0	ARG	-	expression tag	UNP Q8IU57
B	207	ALA	-	expression tag	UNP Q8IU57
B	208	ALA	-	expression tag	UNP Q8IU57
B	209	ALA	-	expression tag	UNP Q8IU57
B	210	LEU	-	expression tag	UNP Q8IU57
B	211	GLU	-	expression tag	UNP Q8IU57
B	212	VAL	-	expression tag	UNP Q8IU57
B	213	LEU	-	expression tag	UNP Q8IU57
B	214	PHE	-	expression tag	UNP Q8IU57
B	215	GLN	-	expression tag	UNP Q8IU57

- Molecule 3 is a protein called Interferon lambda-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	151	1161	729	212	213	7	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	expression tag	UNP Q8IZI9
C	0	SER	-	expression tag	UNP Q8IZI9
C	15	ARG	GLN	engineered mutation	UNP Q8IZI9
C	73	ASP	GLU	engineered mutation	UNP Q8IZI9
C	120	ARG	HIS	engineered mutation	UNP Q8IZI9
C	150	ALA	THR	engineered mutation	UNP Q8IZI9
C	163	GLU	-	expression tag	UNP Q8IZI9
C	164	ALA	-	expression tag	UNP Q8IZI9
C	165	ALA	-	expression tag	UNP Q8IZI9
C	166	ALA	-	expression tag	UNP Q8IZI9
C	167	HIS	-	expression tag	UNP Q8IZI9
C	168	HIS	-	expression tag	UNP Q8IZI9
C	169	HIS	-	expression tag	UNP Q8IZI9
C	170	HIS	-	expression tag	UNP Q8IZI9
C	171	HIS	-	expression tag	UNP Q8IZI9
C	172	HIS	-	expression tag	UNP Q8IZI9
C	173	HIS	-	expression tag	UNP Q8IZI9
C	174	HIS	-	expression tag	UNP Q8IZI9

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	B	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	7	7	7	0	0
5	B	5	5	5	0	0
5	C	6	6	6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-10 receptor subunit beta



- Molecule 2: Interferon lambda receptor 1



- Molecule 3: Interferon lambda-3



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.79Å 106.79Å 129.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.24 – 2.85 46.24 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.24-2.85) 93.4 (46.24-2.85)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.00 (at 2.86Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.198 , 0.253 0.197 , 0.252	Depositor DCC
$R_{free}$ test set	1036 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	99.2	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 94.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4266	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	127.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/1656	0.51	0/2262
2	B	0.29	0/1484	0.48	0/2042
3	C	0.29	0/1181	0.46	0/1604
All	All	0.30	0/4321	0.49	0/5908

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1608	0	1494	11	0
2	B	1437	0	1349	18	0
3	C	1161	0	1132	5	0
4	B	42	0	39	1	0
5	A	7	0	0	0	0
5	B	5	0	0	0	0
5	C	6	0	0	0	0
All	All	4266	0	4014	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:51:VAL:HG12	2:B:54:CYS:H	1.56	0.70
1:A:156:VAL:HB	1:A:169:ILE:HG13	1.77	0.65
2:B:37:VAL:HG23	2:B:51:VAL:HB	1.83	0.61
1:A:81:LYS:HA	1:A:107:PRO:HG2	1.83	0.60
2:B:174:HIS:CE1	2:B:200:LEU:HD13	2.37	0.60
2:B:39:TYR:CE1	2:B:49:ARG:HB3	2.40	0.57
2:B:67:LEU:HB2	2:B:70:GLN:HB2	1.86	0.56
3:C:27:LEU:HD21	3:C:86:LEU:HD11	1.88	0.54
2:B:159:GLY:HA2	4:B:301:NAG:H5	1.90	0.53
2:B:105:PRO:HG2	2:B:190:SER:HB3	1.93	0.50
2:B:34:THR:OG1	2:B:83:VAL:O	2.29	0.50
1:A:155:ASN:HB3	1:A:192:ARG:HG2	1.95	0.49
2:B:172:GLU:HB3	2:B:174:HIS:CE1	2.48	0.49
1:A:142:THR:HG22	2:B:155:VAL:HG11	1.95	0.48
1:A:68:GLN:NE2	1:A:92:GLU:OE1	2.46	0.48
1:A:80:SER:O	1:A:81:LYS:HG2	2.15	0.47
1:A:58:SER:HB2	3:C:87:ASP:OD2	2.15	0.47
2:B:108:PRO:HB3	2:B:125:TYR:CZ	2.50	0.46
1:A:32:VAL:HG22	1:A:112:ILE:HB	1.98	0.46
3:C:69:LEU:HD12	3:C:90:LEU:HD13	1.98	0.46
2:B:143:TRP:HB3	2:B:150:LYS:HA	1.99	0.45
1:A:45:PHE:CE2	1:A:47:LYS:HB2	2.52	0.43
1:A:146:LYS:HG3	1:A:152:TRP:CD1	2.54	0.43
2:B:83:VAL:HG12	2:B:88:LYS:HG2	2.01	0.42
3:C:129:PRO:HA	3:C:137:LEU:HD11	2.01	0.41
2:B:119:LEU:HB2	2:B:166:LEU:HD11	2.02	0.41
3:C:42:LEU:HD12	3:C:43:PHE:CG	2.54	0.41
2:B:128:PRO:HA	2:B:129:PRO:HD3	1.95	0.41
1:A:158:TYR:HB3	1:A:189:VAL:HG12	2.02	0.41
2:B:37:VAL:CG2	2:B:54:CYS:HB2	2.51	0.41
2:B:50:GLU:OE1	2:B:50:GLU:N	2.54	0.40
2:B:119:LEU:HD21	2:B:176:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/214 (92%)	184 (94%)	12 (6%)	0	100	100
2	B	184/218 (84%)	172 (94%)	12 (6%)	0	100	100
3	C	147/176 (84%)	143 (97%)	4 (3%)	0	100	100
All	All	527/608 (87%)	499 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/195 (90%)	175 (99%)	1 (1%)	86	93
2	B	155/196 (79%)	153 (99%)	2 (1%)	69	84
3	C	122/149 (82%)	122 (100%)	0	100	100
All	All	453/540 (84%)	450 (99%)	3 (1%)	84	91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	TRP
2	B	60	LEU
2	B	201	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
2	B	174	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	B	302	2	14,14,15	0.20	0	17,19,21	0.46	0
4	NAG	B	303	2	14,14,15	0.36	0	17,19,21	0.50	0
4	NAG	B	301	2	14,14,15	0.15	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	302	2	-	2/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	303	2	-	4/6/23/26	0/1/1/1
4	NAG	B	301	2	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	301	NAG	O5-C5-C6-O6
4	B	302	NAG	C8-C7-N2-C2
4	B	302	NAG	O7-C7-N2-C2
4	B	303	NAG	C8-C7-N2-C2
4	B	303	NAG	O7-C7-N2-C2
4	B	303	NAG	O5-C5-C6-O6
4	B	301	NAG	C4-C5-C6-O6
4	B	303	NAG	C4-C5-C6-O6
4	B	301	NAG	C1-C2-N2-C7
4	B	301	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/214 (92%)	0.31	8 (4%) 38 30	71, 114, 183, 228	0
2	B	188/218 (86%)	0.68	24 (12%) 3 2	79, 138, 202, 226	0
3	C	151/176 (85%)	0.29	4 (2%) 56 51	63, 114, 189, 260	0
All	All	537/608 (88%)	0.43	36 (6%) 17 12	63, 123, 200, 260	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	82	THR	6.5
2	B	8	GLN	4.9
2	B	92	VAL	4.9
2	B	23	TRP	4.6
2	B	90	PRO	3.8
2	B	7	PRO	3.8
2	B	34	THR	3.7
2	B	83	VAL	3.5
1	A	217	THR	3.4
2	B	80	VAL	3.4
1	A	44	ALA	3.3
2	B	88	LYS	3.2
2	B	35	TYR	3.2
2	B	11	THR	3.2
2	B	89	SER	3.1
2	B	25	PRO	3.1
2	B	10	VAL	3.0
2	B	143	TRP	2.9
1	A	93	PHE	2.9
2	B	176	LEU	2.7
1	A	50	LEU	2.7
2	B	85	PRO	2.6
2	B	200	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	C	5	HIS	2.5
2	B	36	PHE	2.4
3	C	129	PRO	2.4
3	C	40	SER	2.4
1	A	42	SER	2.3
2	B	12	LEU	2.3
3	C	4	CYS	2.3
2	B	78	GLY	2.3
2	B	84	SER	2.3
1	A	24	PRO	2.2
2	B	147	ALA	2.1
1	A	164	ASP	2.0
1	A	52	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	B	302	14/15	0.73	0.36	158,183,203,208	0
4	NAG	B	301	14/15	0.85	0.29	102,146,176,179	0
4	NAG	B	303	14/15	0.90	0.26	151,168,176,177	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.