



wwPDB EM Validation Summary Report ⓘ

Sep 17, 2024 – 02:25 pm BST

PDB ID : 6TBM
EMDB ID : EMD-10446
Title : Structure of SAGA bound to TBP, including Spt8 and DUB
Authors : Papai, G.; Frechard, A.; Kolesnikova, O.; Crucifix, C.; Schultz, P.; Ben-Shem, A.
Deposited on : 2019-11-01
Resolution : 20.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev112
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 20.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|--------------------------|--------------------------|
| Clashscore | 210492 | 15764 |
| Ramachandran outliers | 207382 | 16835 |
| Sidechain outliers | 206894 | 16415 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | M | 240 | |
| 2 | A | 448 | |
| 3 | C | 698 | |
| 4 | F | 517 | |
| 5 | D | 341 | |
| 6 | E | 1191 | |
| 7 | J | 217 | |
| 8 | K | 609 | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 9 | G | 722 | |
| 10 | H | 485 | |
| 11 | I | 153 | |
| 12 | L | 3825 | |
| 13 | B | 722 | |
| 14 | N | 400 | |
| 15 | R | 76 | |
| 16 | Q | 502 | |
| 17 | O | 123 | |
| 18 | P | 96 | |

2 Entry composition

There are 19 unique types of molecules in this entry. The entry contains 48800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TATA-box Binding Protein (TBP).

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 1 | M | 180 | 1415 | 921 | 242 | 246 | 6 | 0 | 0 |

- Molecule 2 is a protein called Transcriptional coactivator HFI1/ADA1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 2 | A | 173 | 1300 | 816 | 228 | 250 | 6 | 0 | 0 |

- Molecule 3 is a protein called Subunit of SAGA histone acetyltransferase complex.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 3 | C | 95 | 761 | 480 | 142 | 133 | 6 | 0 | 0 |

There are 54 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| C | 645 | LEU | - | expression tag | UNP C4QZ39 |
| C | 646 | GLU | - | expression tag | UNP C4QZ39 |
| C | 647 | GLY | - | expression tag | UNP C4QZ39 |
| C | 648 | GLY | - | expression tag | UNP C4QZ39 |
| C | 649 | GLY | - | expression tag | UNP C4QZ39 |
| C | 650 | GLY | - | expression tag | UNP C4QZ39 |
| C | 651 | SER | - | expression tag | UNP C4QZ39 |
| C | 652 | MET | - | expression tag | UNP C4QZ39 |
| C | 653 | ASP | - | expression tag | UNP C4QZ39 |
| C | 654 | GLU | - | expression tag | UNP C4QZ39 |
| C | 655 | LYS | - | expression tag | UNP C4QZ39 |
| C | 656 | THR | - | expression tag | UNP C4QZ39 |
| C | 657 | THR | - | expression tag | UNP C4QZ39 |
| C | 658 | GLY | - | expression tag | UNP C4QZ39 |
| C | 659 | TRP | - | expression tag | UNP C4QZ39 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| C | 660 | ARG | - | expression tag | UNP C4QZ39 |
| C | 661 | GLY | - | expression tag | UNP C4QZ39 |
| C | 662 | GLY | - | expression tag | UNP C4QZ39 |
| C | 663 | HIS | - | expression tag | UNP C4QZ39 |
| C | 664 | VAL | - | expression tag | UNP C4QZ39 |
| C | 665 | VAL | - | expression tag | UNP C4QZ39 |
| C | 666 | GLU | - | expression tag | UNP C4QZ39 |
| C | 667 | GLY | - | expression tag | UNP C4QZ39 |
| C | 668 | LEU | - | expression tag | UNP C4QZ39 |
| C | 669 | ALA | - | expression tag | UNP C4QZ39 |
| C | 670 | GLY | - | expression tag | UNP C4QZ39 |
| C | 671 | GLU | - | expression tag | UNP C4QZ39 |
| C | 672 | LEU | - | expression tag | UNP C4QZ39 |
| C | 673 | GLU | - | expression tag | UNP C4QZ39 |
| C | 674 | GLN | - | expression tag | UNP C4QZ39 |
| C | 675 | LEU | - | expression tag | UNP C4QZ39 |
| C | 676 | ARG | - | expression tag | UNP C4QZ39 |
| C | 677 | ALA | - | expression tag | UNP C4QZ39 |
| C | 678 | ARG | - | expression tag | UNP C4QZ39 |
| C | 679 | LEU | - | expression tag | UNP C4QZ39 |
| C | 680 | GLU | - | expression tag | UNP C4QZ39 |
| C | 681 | HIS | - | expression tag | UNP C4QZ39 |
| C | 682 | HIS | - | expression tag | UNP C4QZ39 |
| C | 683 | PRO | - | expression tag | UNP C4QZ39 |
| C | 684 | GLN | - | expression tag | UNP C4QZ39 |
| C | 685 | GLY | - | expression tag | UNP C4QZ39 |
| C | 686 | GLN | - | expression tag | UNP C4QZ39 |
| C | 687 | ARG | - | expression tag | UNP C4QZ39 |
| C | 688 | GLU | - | expression tag | UNP C4QZ39 |
| C | 689 | PRO | - | expression tag | UNP C4QZ39 |
| C | 690 | GLY | - | expression tag | UNP C4QZ39 |
| C | 691 | GLY | - | expression tag | UNP C4QZ39 |
| C | 692 | SER | - | expression tag | UNP C4QZ39 |
| C | 693 | HIS | - | expression tag | UNP C4QZ39 |
| C | 694 | HIS | - | expression tag | UNP C4QZ39 |
| C | 695 | HIS | - | expression tag | UNP C4QZ39 |
| C | 696 | HIS | - | expression tag | UNP C4QZ39 |
| C | 697 | HIS | - | expression tag | UNP C4QZ39 |
| C | 698 | HIS | - | expression tag | UNP C4QZ39 |

- Molecule 4 is a protein called Spt20.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 4 | F | 210 | 1682 | 1071 | 292 | 315 | 4 | 0 | 0 |

- Molecule 5 is a protein called Subunit of the SAGA and SAGA-like transcriptional regulatory complexes, interacts with Spt15p to act.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 5 | D | 209 | 1616 | 1016 | 298 | 295 | 7 | 0 | 0 |

- Molecule 6 is a protein called Subunit of the SAGA transcriptional regulatory complex, involved in proper assembly of the complex.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 6 | E | 154 | 1232 | 784 | 208 | 233 | 7 | 0 | 0 |

- Molecule 7 is a protein called Transcription initiation factor TFIID subunit 10.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 7 | J | 96 | 768 | 489 | 120 | 156 | 3 | 0 | 0 |

- Molecule 8 is a protein called Subunit (61/68 kDa) of TFIID and SAGA complexes.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 8 | K | 154 | 1192 | 747 | 216 | 226 | 3 | 0 | 0 |

- Molecule 9 is a protein called Subunit (90 kDa) of TFIID and SAGA complexes.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 9 | G | 522 | 4075 | 2581 | 719 | 756 | 19 | 0 | 0 |

- Molecule 10 is a protein called Subunit (60 kDa) of TFIID and SAGA complexes.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 10 | H | 421 | 3263 | 2084 | 556 | 617 | 6 | 0 | 0 |

- Molecule 11 is a protein called Subunit (17 kDa) of TFIID and SAGA complexes, involved

in RNA polymerase II transcription initiation.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 11 | I | 123 | 981 | 632 | 169 | 178 | 2 | 0 | 0 |

- Molecule 12 is a protein called Transcription-associated protein.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-------|------|------|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 12 | L | 2968 | 22318 | 14296 | 3864 | 4071 | 87 | 0 | 0 |

- Molecule 13 is a protein called Transcriptional regulator involved in glucose repression of Gal4p-regulated genes.

| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---------|-------|
| | | | Total | C | N | O | | |
| 13 | B | 76 | 373 | 221 | 76 | 76 | 0 | 0 |

- Molecule 14 is a protein called Spt8.

| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| | | | Total | C | N | O | | |
| 14 | N | 335 | 1648 | 978 | 335 | 335 | 0 | 0 |

- Molecule 15 is a protein called Polyubiquitin-B.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 15 | R | 76 | 611 | 384 | 105 | 121 | 1 | 2 | 0 |

There is a discrepancy between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| R | 76 | GLZ | - | expression tag | UNP J3QS39 |

- Molecule 16 is a protein called Ubiquitin carboxyl-terminal hydrolase.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| | | | Total | C | N | O | S | | |
| 16 | Q | 482 | 3817 | 2410 | 657 | 719 | 31 | 0 | 0 |

- Molecule 17 is a protein called SAGA-associated factor 11.


| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 17 | O | 117 | 922 | 563 | 169 | 184 | 6 | 0 | 0 |

- Molecule 18 is a protein called Transcription and mRNA export factor SUS1.


| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| | | | Total | C | N | O | S | | |
| 18 | P | 90 | 735 | 470 | 122 | 142 | 1 | 0 | 0 |


- Molecule 19 is water.


| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|-------|----|---------|
| 19 | R | 83 | Total | O | 0 |
| | | | 83 | 83 | |
| 19 | Q | 8 | Total | O | 0 |
| | | | 8 | 8 | |



● Molecule 10: Subunit (60 kDa) of TFIID and SAGA complexes









● Molecule 13: Transcriptional regulator involved in glucose repression of Gal4p-regulated genes

Chain B:  10%  89%






• Molecule 16: Ubiquitin carboxyl-terminal hydrolase




• Molecule 17: SAGA-associated factor 11



• Molecule 18: Transcription and mRNA export factor SUS1





4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, C1 | Depositor |
| Number of particles used | 354104 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 52.8 | Depositor |
| Minimum defocus (nm) | 800 | Depositor |
| Maximum defocus (nm) | 4500 | Depositor |
| Magnification | 105000 | Depositor |
| Image detector | GATAN K2 QUANTUM (4k x 4k) | Depositor |
| Maximum map value | 0.032 | Depositor |
| Minimum map value | -0.024 | Depositor |
| Average map value | 0.000 | Depositor |
| Map value standard deviation | 0.001 | Depositor |
| Recommended contour level | 0.004 | Depositor |
| Map size (Å) | 558.08, 558.08, 558.08 | wwPDB |
| Map dimensions | 256, 256, 256 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 2.18, 2.18, 2.18 | Depositor |

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | M | 0.88 | 2/1442 (0.1%) | 0.78 | 2/1942 (0.1%) |
| 2 | A | 0.44 | 0/1319 | 0.60 | 0/1794 |
| 3 | C | 0.53 | 0/774 | 0.75 | 0/1039 |
| 4 | F | 0.34 | 0/1718 | 0.58 | 0/2335 |
| 5 | D | 0.51 | 0/1641 | 0.65 | 0/2213 |
| 6 | E | 0.45 | 0/1246 | 0.62 | 0/1667 |
| 7 | J | 0.47 | 0/779 | 0.60 | 0/1051 |
| 8 | K | 0.42 | 0/1213 | 0.66 | 0/1647 |
| 9 | G | 0.52 | 0/4177 | 0.60 | 0/5661 |
| 10 | H | 0.38 | 0/3315 | 0.60 | 0/4500 |
| 11 | I | 0.44 | 0/1006 | 0.63 | 0/1374 |
| 12 | L | 0.33 | 0/22712 | 0.54 | 0/30825 |
| 13 | B | 0.64 | 0/370 | 0.70 | 1/509 (0.2%) |
| 15 | R | 0.83 | 1/619 (0.2%) | 0.75 | 0/833 |
| 16 | Q | 0.71 | 0/3897 | 1.09 | 15/5265 (0.3%) |
| 17 | O | 0.79 | 0/931 | 1.13 | 2/1250 (0.2%) |
| 18 | P | 0.75 | 0/747 | 1.01 | 0/1011 |
| All | All | 0.47 | 3/47906 (0.0%) | 0.67 | 20/64916 (0.0%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | M | 0 | 1 |
| 16 | Q | 2 | 5 |
| All | All | 2 | 6 |

All (3) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 1 | M | 194 | ILE | C-N | 27.38 | 1.97 | 1.34 |
| 1 | M | 183 | SER | C-N | 6.38 | 1.48 | 1.34 |
| 15 | R | 24 | GLU | CB-CG | -5.25 | 1.42 | 1.52 |

The worst 5 of 20 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|-------|-------------|----------|
| 16 | Q | 61 | TYR | CB-CG-CD1 | 9.83 | 126.90 | 121.00 |
| 16 | Q | 61 | TYR | CB-CG-CD2 | -9.38 | 115.37 | 121.00 |
| 1 | M | 62 | GLY | N-CA-C | 6.94 | 130.46 | 113.10 |
| 16 | Q | 78 | ARG | NE-CZ-NH1 | -6.80 | 116.90 | 120.30 |
| 13 | B | 527 | PRO | N-CA-CB | 6.57 | 111.18 | 103.30 |

All (2) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 16 | Q | 31 | SER | CA |
| 16 | Q | 222 | THR | CA |

5 of 6 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 1 | M | 183 | SER | Mainchain |
| 16 | Q | 149 | ARG | Sidechain |
| 16 | Q | 212 | TYR | Sidechain |
| 16 | Q | 38 | TYR | Sidechain |
| 16 | Q | 61 | TYR | Sidechain |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | M | 1415 | 0 | 1491 | 102 | 0 |
| 2 | A | 1300 | 0 | 1254 | 30 | 0 |
| 3 | C | 761 | 0 | 779 | 194 | 0 |
| 4 | F | 1682 | 0 | 1622 | 30 | 0 |
| 5 | D | 1616 | 0 | 1558 | 122 | 0 |
| 6 | E | 1232 | 0 | 1276 | 40 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 7 | J | 768 | 0 | 754 | 11 | 0 |
| 8 | K | 1192 | 0 | 1214 | 89 | 0 |
| 9 | G | 4075 | 0 | 3934 | 102 | 0 |
| 10 | H | 3263 | 0 | 3258 | 69 | 0 |
| 11 | I | 981 | 0 | 982 | 18 | 0 |
| 12 | L | 22318 | 0 | 20960 | 376 | 0 |
| 13 | B | 373 | 0 | 190 | 2 | 0 |
| 14 | N | 1648 | 0 | 382 | 61 | 0 |
| 15 | R | 611 | 0 | 636 | 20 | 0 |
| 16 | Q | 3817 | 0 | 3772 | 521 | 0 |
| 17 | O | 922 | 0 | 897 | 411 | 0 |
| 18 | P | 735 | 0 | 743 | 191 | 0 |
| 19 | Q | 8 | 0 | 0 | 10 | 0 |
| 19 | R | 83 | 0 | 0 | 7 | 0 |
| All | All | 48800 | 0 | 45702 | 1602 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 1602 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-------------------|------------------|--------------------------|-------------------|
| 16:Q:472:PHE:CZ | 17:O:45:GLN:HB3 | 1.29 | 1.65 |
| 17:O:11:TRP:CD1 | 18:P:83:LEU:HD23 | 1.14 | 1.61 |
| 16:Q:479:ILE:HG23 | 17:O:47:SER:CA | 1.19 | 1.61 |
| 16:Q:479:ILE:CG2 | 17:O:47:SER:HA | 1.14 | 1.60 |
| 17:O:34:ARG:HH11 | 18:P:57:PRO:CB | 1.11 | 1.58 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|------------------|------------|----------|----------|-------------|-----|
| 1 | M | 178/240 (74%) | 147 (83%) | 24 (14%) | 7 (4%) | 2 | 19 |
| 2 | A | 169/448 (38%) | 151 (89%) | 18 (11%) | 0 | 100 | 100 |
| 3 | C | 89/698 (13%) | 83 (93%) | 6 (7%) | 0 | 100 | 100 |
| 4 | F | 202/517 (39%) | 177 (88%) | 24 (12%) | 1 (0%) | 25 | 64 |
| 5 | D | 203/341 (60%) | 177 (87%) | 19 (9%) | 7 (3%) | 3 | 21 |
| 6 | E | 150/1191 (13%) | 135 (90%) | 13 (9%) | 2 (1%) | 10 | 43 |
| 7 | J | 92/217 (42%) | 82 (89%) | 9 (10%) | 1 (1%) | 12 | 47 |
| 8 | K | 150/609 (25%) | 125 (83%) | 23 (15%) | 2 (1%) | 10 | 43 |
| 9 | G | 512/722 (71%) | 448 (88%) | 63 (12%) | 1 (0%) | 44 | 78 |
| 10 | H | 413/485 (85%) | 365 (88%) | 48 (12%) | 0 | 100 | 100 |
| 11 | I | 119/153 (78%) | 98 (82%) | 21 (18%) | 0 | 100 | 100 |
| 12 | L | 2874/3825 (75%) | 2623 (91%) | 233 (8%) | 18 (1%) | 22 | 60 |
| 13 | B | 70/722 (10%) | 63 (90%) | 7 (10%) | 0 | 100 | 100 |
| 15 | R | 76/76 (100%) | 75 (99%) | 1 (1%) | 0 | 100 | 100 |
| 16 | Q | 480/502 (96%) | 444 (92%) | 30 (6%) | 6 (1%) | 10 | 43 |
| 17 | O | 115/123 (94%) | 103 (90%) | 6 (5%) | 6 (5%) | 1 | 15 |
| 18 | P | 88/96 (92%) | 85 (97%) | 3 (3%) | 0 | 100 | 100 |
| All | All | 5980/10965 (54%) | 5381 (90%) | 548 (9%) | 51 (1%) | 17 | 52 |

5 of 51 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | M | 77 | GLY |
| 1 | M | 99 | PHE |
| 1 | M | 110 | LYS |
| 5 | D | 200 | THR |
| 5 | D | 327 | ARG |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|-----|
| 1 | M | 152/205 (74%) | 138 (91%) | 14 (9%) | 7 | 23 |
| 2 | A | 134/394 (34%) | 132 (98%) | 2 (2%) | 60 | 75 |
| 3 | C | 85/627 (14%) | 85 (100%) | 0 | 100 | 100 |
| 4 | F | 179/471 (38%) | 179 (100%) | 0 | 100 | 100 |
| 5 | D | 166/306 (54%) | 154 (93%) | 12 (7%) | 12 | 32 |
| 6 | E | 142/1101 (13%) | 136 (96%) | 6 (4%) | 25 | 46 |
| 7 | J | 85/183 (46%) | 84 (99%) | 1 (1%) | 67 | 78 |
| 8 | K | 133/524 (25%) | 130 (98%) | 3 (2%) | 45 | 64 |
| 9 | G | 439/635 (69%) | 430 (98%) | 9 (2%) | 48 | 66 |
| 10 | H | 352/438 (80%) | 348 (99%) | 4 (1%) | 70 | 80 |
| 11 | I | 104/130 (80%) | 103 (99%) | 1 (1%) | 73 | 82 |
| 12 | L | 2156/3450 (62%) | 2114 (98%) | 42 (2%) | 52 | 69 |
| 15 | R | 70/68 (103%) | 69 (99%) | 1 (1%) | 62 | 75 |
| 16 | Q | 431/449 (96%) | 425 (99%) | 6 (1%) | 62 | 75 |
| 17 | O | 103/109 (94%) | 102 (99%) | 1 (1%) | 73 | 82 |
| 18 | P | 85/91 (93%) | 85 (100%) | 0 | 100 | 100 |
| All | All | 4816/9181 (52%) | 4714 (98%) | 102 (2%) | 49 | 66 |

5 of 102 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 12 | L | 1122 | ASN |
| 12 | L | 2489 | ARG |
| 16 | Q | 254 | LEU |
| 12 | L | 1314 | LEU |
| 12 | L | 1419 | LYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 12 | L | 2884 | GLN |
| 17 | O | 76 | ASN |
| 15 | R | 60 | ASN |
| 18 | P | 36 | GLN |
| 17 | O | 26 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|-------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 15 | GLZ | R | 76 | 15,16 | 3,3,3 | 0.70 | 0 | 0,2,2 | - | - |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|-------|---------|----------|-------|
| 15 | GLZ | R | 76 | 15,16 | - | 0/0/1/1 | - |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 14 | N | 6 |
| 13 | B | 2 |
| 1 | M | 1 |

The worst 5 of 9 chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1 | B | 414:ASP | C | 462:ALA | N | 28.86 |
| 1 | B | 471:ASP | C | 509:ASP | N | 23.98 |
| 1 | N | 117:UNK | C | 172:UNK | N | 15.59 |
| 1 | N | 287:UNK | C | 332:UNK | N | 5.97 |
| 1 | N | 213:UNK | C | 214:UNK | N | 4.21 |


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10446. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 121



Y Index: 93




Z Index: 133


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)

6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.004. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1341 nm³; this corresponds to an approximate mass of 1211 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.050\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-10446 and PDB model 6TBM. Per-residue inclusion information can be found in section [3](#) on page [9](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.004 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.004).







































9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.004) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|--|--|
| All |  0.7880 |  0.0800 |
| A |  0.7340 |  0.0550 |
| B |  1.0000 |  0.1030 |
| C |  0.5900 |  0.0520 |
| D |  0.8780 |  0.0850 |
| E |  0.9840 |  0.1110 |
| F |  0.8990 |  0.0920 |
| G |  0.8960 |  0.0740 |
| H |  0.9010 |  0.0800 |
| I |  0.8680 |  0.0710 |
| J |  0.9480 |  0.0730 |
| K |  0.8180 |  0.0500 |
| L |  0.9280 |  0.1010 |
| M |  0.8060 |  0.0760 |
| N |  0.0380 |  0.0210 |
| O |  0.1700 |  0.0130 |
| P |  0.2190 |  0.0410 |
| Q |  0.2220 |  0.0270 |
| R |  0.3800 |  0.0620 |

