



wwPDB EM Validation Summary Report ⓘ

Dec 11, 2022 – 11:39 pm GMT

PDB ID : 6TDY
EMDB ID : EMD-10471
Title : Cryo-EM structure of *Euglena gracilis* mitochondrial ATP synthase, OSCP/F1/c-ring in rotational state 1
Authors : Muhleip, A.; Amunts, A.
Deposited on : 2019-11-10
Resolution : 3.04 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 158937 | 4297 |
| Ramachandran outliers | 154571 | 4023 |
| Sidechain outliers | 154315 | 3826 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 561 | |
| 1 | B | 561 | |
| 1 | C | 561 | |
| 2 | D | 494 | |
| 2 | E | 494 | |
| 2 | F | 494 | |
| 3 | G | 306 | |
| 4 | H | 176 | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 5 | I | 76 | |
| 6 | J | 192 | |
| 6 | K | 192 | |
| 6 | L | 192 | |
| 7 | M | 267 | |
| 8 | N | 103 | |
| 9 | O | 104 | |
| 9 | P | 104 | |
| 9 | Q | 104 | |
| 9 | R | 104 | |
| 9 | S | 104 | |
| 9 | T | 104 | |
| 9 | U | 104 | |
| 9 | V | 104 | |
| 9 | W | 104 | |
| 9 | X | 104 | |
| 10 | h | 476 | |
| 11 | c | 169 | |

2 Entry composition [i](#)

There are 15 unique types of molecules in this entry. The entry contains 81811 atoms, of which 41207 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP synthase subunit alpha.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|------|------|-----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 1 | A | 526 | Total | C | H | N | O | S | 0 | 0 |
| | | | 8292 | 2611 | 4197 | 700 | 766 | 18 | | |
| 1 | B | 526 | Total | C | H | N | O | S | 0 | 0 |
| | | | 8292 | 2611 | 4197 | 700 | 766 | 18 | | |
| 1 | C | 529 | Total | C | H | N | O | S | 0 | 0 |
| | | | 8336 | 2625 | 4219 | 703 | 771 | 18 | | |

- Molecule 2 is a protein called ATP synthase subunit beta.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|------|------|-----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 2 | D | 487 | Total | C | H | N | O | S | 0 | 0 |
| | | | 7407 | 2318 | 3729 | 620 | 713 | 27 | | |
| 2 | E | 487 | Total | C | H | N | O | S | 0 | 0 |
| | | | 7408 | 2318 | 3730 | 620 | 713 | 27 | | |
| 2 | F | 487 | Total | C | H | N | O | S | 0 | 0 |
| | | | 7407 | 2318 | 3729 | 620 | 713 | 27 | | |

- Molecule 3 is a protein called ATP synthase subunit gamma.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|------|------|-----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 3 | G | 303 | Total | C | H | N | O | S | 0 | 0 |
| | | | 4898 | 1543 | 2462 | 420 | 459 | 14 | | |

- Molecule 4 is a protein called ATP synthase subunit delta.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|-----|------|-----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 4 | H | 160 | Total | C | H | N | O | S | 0 | 0 |
| | | | 2448 | 787 | 1202 | 207 | 251 | 1 | | |

- Molecule 5 is a protein called ATP synthase subunit epsilon.

| Mol | Chain | Residues | Atoms | | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|----|----|---|---------|-------|
| 5 | I | 66 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1077 | 346 | 541 | 91 | 98 | 1 | | |

- Molecule 6 is a protein called p18.

| Mol | Chain | Residues | Atoms | | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|------|-----|-----|---|---------|-------|
| 6 | J | 170 | Total | C | H | N | O | S | 0 | 0 |
| | | | 2596 | 829 | 1294 | 217 | 249 | 7 | | |
| 6 | K | 170 | Total | C | H | N | O | S | 0 | 0 |
| | | | 2596 | 829 | 1294 | 217 | 249 | 7 | | |
| 6 | L | 170 | Total | C | H | N | O | S | 0 | 0 |
| | | | 2596 | 829 | 1294 | 217 | 249 | 7 | | |

- Molecule 7 is a protein called oligomycin sensitivity conferring protein (OSCP).

| Mol | Chain | Residues | Atoms | | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|-----|-----|---|---------|-------|
| 7 | M | 243 | Total | C | H | N | O | S | 0 | 0 |
| | | | 3778 | 1212 | 1885 | 310 | 370 | 1 | | |

- Molecule 8 is a protein called inhibitor of F1 (IF1).

| Mol | Chain | Residues | Atoms | | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|----|----|---|---------|-------|
| 8 | N | 49 | Total | C | H | N | O | S | 0 | 0 |
| | | | 802 | 247 | 399 | 72 | 82 | 2 | | |

- Molecule 9 is a protein called ATP synthase subunit c.

| Mol | Chain | Residues | Atoms | | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|----|-----|---|---------|-------|
| 9 | O | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | P | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | Q | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | R | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | S | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | T | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | U | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|-----|-----|----|-----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 9 | V | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | W | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |
| 9 | X | 81 | Total | C | H | N | O | S | 0 | 0 |
| | | | 1185 | 383 | 605 | 89 | 102 | 6 | | |


- Molecule 10 is a protein called ATP synthase subunit d.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|-----|-----|----|----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 10 | h | 49 | Total | C | H | N | O | S | 0 | 0 |
| | | | 807 | 255 | 408 | 67 | 76 | 1 | | |

- Molecule 11 is a protein called ATPTB4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace | |
|-----|-------|----------|-------|-----|-----|----|----|---------|-------|---|
| | | | Total | C | H | N | O | | | S |
| 11 | c | 57 | Total | C | H | N | O | S | 0 | 0 |
| | | | 944 | 300 | 481 | 78 | 83 | 2 | | |

- Molecule 12 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | | | AltConf | |
|-----|-------|----------|-------|----|----|---|----|---------|---|
| | | | Total | C | H | N | O | | P |
| 12 | A | 1 | Total | C | H | N | O | P | 0 |
| | | | 43 | 10 | 12 | 5 | 13 | 3 | |

Continued on next page...


Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | AltConf | |
|-----|-------|----------|-------------|---------|---------|--------|---------|---------|---|
| | | | Total | C | H | N | O | | P |
| 12 | B | 1 | Total 43 | C 10 | H 12 | N 5 | O 13 | P 3 | 0 |
| 12 | C | 1 | Total 43 | C 10 | H 12 | N 5 | O 13 | P 3 | 0 |
| 12 | F | 1 | Total 43 | C 10 | H 12 | N 5 | O 13 | P 3 | 0 |

- Molecule 13 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


| Mol | Chain | Residues | Atoms | | AltConf |
|-----|-------|----------|------------|---------|---------|
| | | | Total | Mg | |
| 13 | A | 1 | Total 1 | Mg 1 | 0 |
| 13 | B | 1 | Total 1 | Mg 1 | 0 |
| 13 | C | 1 | Total 1 | Mg 1 | 0 |
| 13 | D | 1 | Total 1 | Mg 1 | 0 |
| 13 | F | 1 | Total 1 | Mg 1 | 0 |

- Molecule 14 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | | | AltConf | |
|-----|-------|----------|-------------|---------|---------|--------|---------|---------|---|
| | | | Total | C | H | N | O | | P |
| 14 | D | 1 | Total 39 | C 10 | H 12 | N 5 | O 10 | P 2 | 0 |

- Molecule 15 is FRAGMENT OF TRITON X-100 (three-letter code: TRT) (formula: $C_{21}H_{36}O_4$).





| Mol | Chain | Residues | Atoms | | | | AltConf |
|-----|-------|----------|-------|----|----|---|---------|
| | | | Total | C | H | O | |
| 15 | E | 1 | 61 | 21 | 36 | 4 | 0 |

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP synthase subunit alpha

Chain A: 





- Molecule 1: ATP synthase subunit alpha


Chain B: 




- Molecule 1: ATP synthase subunit alpha

Chain C: 








• Molecule 2: ATP synthase subunit beta




• Molecule 2: ATP synthase subunit beta



• Molecule 2: ATP synthase subunit beta





• Molecule 3: ATP synthase subunit gamma









• Molecule 4: ATP synthase subunit delta





• Molecule 5: ATP synthase subunit epsilon





• Molecule 6: p18




• Molecule 6: p18




• Molecule 6: p18




• Molecule 7: oligomycin sensitivity conferring protein (OSCP)




• Molecule 9: ATP synthase subunit c




• Molecule 9: ATP synthase subunit c



• Molecule 9: ATP synthase subunit c




• Molecule 9: ATP synthase subunit c





• Molecule 9: ATP synthase subunit c







• Molecule 9: ATP synthase subunit c



• Molecule 9: ATP synthase subunit c



• Molecule 10: ATP synthase subunit d



GLN
ALA
LEU
THR
ARG
LEU
HIS
LEU
GLU
ARG
ARG
LEU
LEU
ARG
GLY
GLN
ASP
VAL
ILE
ASP
LEU
GLU
TRP
ALA
TRP
GLN
ASP
VAL
ASN
PRO
ILE
ASP
LEU
ASN
GLU
ALA
TRP
ILE
LEU
LEU
ALA
LEU
GLN
ASP
THR
HIS
TYR
HIS
HIS
PRO
ALA
LEU
VAL
ARG
ASP
VAL
ASP
HIS
THR
TYR
LEU
LEU
ALA
ILE
PHE
LEU
GLY
ASP
PRO
THR
GLU
VAL
PHE
ARG
GLY
ALA
THR
ALA
TYR
ILE
TRP
LYS
LEU
PHE
PRO
HIS
THR
ASP
GLY
GLU
GLU
MET
PHE

LYS
ASP
THR
ARG
LEU
PHE
ASP
VAL
LEU
LEU
PRO
ARG
GLN
VAL
ASN
PRO
ILE
ASP
LEU
ASN
GLU
ALA
TRP
ILE
LEU
LEU
ALA
LEU
GLN
ASP
THR
HIS
TYR
HIS
HIS
PRO
ALA
LEU
VAL
ARG
ASP
VAL
ASP
HIS
THR
TYR
LEU
LEU
ALA
ILE
PHE
LEU
GLY
ASP
PRO
THR
GLU
VAL
PHE
ARG
GLY
ALA
THR
ALA
TYR
ILE
TRP
LYS
LEU
PHE
PRO
HIS
THR
ASP
GLY
GLU
GLU
MET
PHE

• Molecule 11: ATPTB4



MET
PHE
ARG
GLY
PHE
LEU
ASP
PRO
VAL
LEU
ALA
ALA
ASP
ALA
V14
K15
F16
L19
Q27
K30
D31
Q32
D37
C38
N39
L40
T41
A42
I43
F44
GLY
ALA
SER
TRP
LYS
ALA
ASP
LEU
ASN
LYS
TRP
PHE
ASP
SER
GLU
TYR
ALA
PRO
LYS
LEU
PRO
ALA
GLU
ARG
ASP
SER

ALA
LYS
SER
LEU
ASP
L78
Y79
L80
K81
R82
V83
D84
R87
E91
P103
GLY
LYS
VAL
ASP
ALA
LEU
THR
GLU
LYS
HIS
LEU
LEU
GLU
THR
GLY
LYS
ALA
ARG
LEU
GLU
LEU
THR
ALA
GLY
LEU
GLY
ASN
LYS
ASP
GLU
GLY
VAL
ASN
ALA
PHE
ARG
LYS
GLU
VAL
GLU

GLN
GLU
GLY
LYS
TYR
ALA
ASN
TRP
PRO
ALA
GLU
LYS
SER
LYS
ALA
LEU
ALA
ASP
LYS
VAL
ILE
ALA
ALA
SER
PRO

4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, C1 | Depositor |
| Number of particles used | 150744 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 36.3 | Depositor |
| Minimum defocus (nm) | Not provided | |
| Maximum defocus (nm) | Not provided | |
| Magnification | 130000 | Depositor |
| Image detector | GATAN K2 QUANTUM (4k x 4k) | Depositor |
| Maximum map value | 0.267 | Depositor |
| Minimum map value | -0.151 | Depositor |
| Average map value | 0.001 | Depositor |
| Map value standard deviation | 0.005 | Depositor |
| Recommended contour level | 0.025 | Depositor |
| Map size (Å) | 461.99997, 461.99997, 461.99997 | wwPDB |
| Map dimensions | 440, 440, 440 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 1.05, 1.05, 1.05 | Depositor |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, TRT, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|----------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 0.50 | 1/4164 (0.0%) | 0.57 | 1/5635 (0.0%) |
| 1 | B | 0.48 | 0/4164 | 0.56 | 0/5635 |
| 1 | C | 0.52 | 0/4186 | 0.60 | 1/5665 (0.0%) |
| 2 | D | 0.47 | 0/3732 | 0.58 | 1/5056 (0.0%) |
| 2 | E | 0.46 | 0/3732 | 0.55 | 0/5056 |
| 2 | F | 0.51 | 0/3732 | 0.58 | 0/5056 |
| 3 | G | 0.41 | 0/2476 | 0.54 | 0/3337 |
| 4 | H | 0.31 | 0/1270 | 0.51 | 0/1720 |
| 5 | I | 0.31 | 0/549 | 0.50 | 0/744 |
| 6 | J | 0.39 | 0/1328 | 0.49 | 0/1792 |
| 6 | K | 0.39 | 0/1328 | 0.49 | 0/1792 |
| 6 | L | 0.41 | 0/1328 | 0.51 | 0/1792 |
| 7 | M | 0.37 | 0/1933 | 0.55 | 0/2623 |
| 8 | N | 0.36 | 0/408 | 0.51 | 0/547 |
| 9 | O | 0.37 | 0/590 | 0.53 | 0/802 |
| 9 | P | 0.32 | 0/590 | 0.50 | 0/802 |
| 9 | Q | 0.31 | 0/590 | 0.50 | 0/802 |
| 9 | R | 0.31 | 0/590 | 0.49 | 0/802 |
| 9 | S | 0.30 | 0/590 | 0.46 | 0/802 |
| 9 | T | 0.36 | 0/590 | 0.52 | 0/802 |
| 9 | U | 0.35 | 0/590 | 0.47 | 0/802 |
| 9 | V | 0.31 | 0/590 | 0.47 | 0/802 |
| 9 | W | 0.33 | 0/590 | 0.52 | 0/802 |
| 9 | X | 0.31 | 0/590 | 0.53 | 0/802 |
| 10 | h | 0.33 | 0/405 | 0.56 | 0/547 |
| 11 | c | 0.38 | 0/470 | 0.55 | 0/635 |
| All | All | 0.44 | 1/41105 (0.0%) | 0.55 | 3/55652 (0.0%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a

sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | C | 0 | 1 |
| 2 | E | 0 | 1 |
| 2 | F | 0 | 1 |
| All | All | 0 | 3 |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 1 | A | 395 | ASN | CB-CG | -5.01 | 1.39 | 1.51 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|------|-------------|----------|
| 1 | A | 117 | ILE | C-N-CA | 7.29 | 139.93 | 121.70 |
| 2 | D | 239 | ARG | NE-CZ-NH1 | 5.62 | 123.11 | 120.30 |
| 1 | C | 435 | ARG | NE-CZ-NH1 | 5.01 | 122.80 | 120.30 |

There are no chirality outliers.

All (3) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 1 | C | 284 | ASP | Peptide |
| 2 | E | 266 | ASP | Peptide |
| 2 | F | 288 | ALA | Peptide |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 4095 | 4197 | 4197 | 27 | 0 |
| 1 | B | 4095 | 4197 | 4197 | 35 | 0 |
| 1 | C | 4117 | 4219 | 4219 | 43 | 0 |
| 2 | D | 3678 | 3729 | 3729 | 32 | 0 |
| 2 | E | 3678 | 3730 | 3730 | 23 | 0 |
| 2 | F | 3678 | 3729 | 3729 | 33 | 0 |
| 3 | G | 2436 | 2462 | 2462 | 19 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 4 | H | 1246 | 1202 | 1202 | 8 | 0 |
| 5 | I | 536 | 541 | 541 | 2 | 0 |
| 6 | J | 1302 | 1294 | 1294 | 6 | 0 |
| 6 | K | 1302 | 1294 | 1294 | 11 | 0 |
| 6 | L | 1302 | 1294 | 1294 | 9 | 0 |
| 7 | M | 1893 | 1885 | 1885 | 14 | 0 |
| 8 | N | 403 | 399 | 399 | 4 | 0 |
| 9 | O | 580 | 605 | 605 | 9 | 0 |
| 9 | P | 580 | 605 | 605 | 7 | 0 |
| 9 | Q | 580 | 605 | 605 | 6 | 0 |
| 9 | R | 580 | 605 | 605 | 10 | 0 |
| 9 | S | 580 | 605 | 605 | 9 | 0 |
| 9 | T | 580 | 605 | 605 | 7 | 0 |
| 9 | U | 580 | 605 | 605 | 8 | 0 |
| 9 | V | 580 | 605 | 605 | 10 | 0 |
| 9 | W | 580 | 605 | 605 | 8 | 0 |
| 9 | X | 580 | 605 | 605 | 11 | 0 |
| 10 | h | 399 | 408 | 408 | 0 | 0 |
| 11 | c | 463 | 481 | 481 | 0 | 0 |
| 12 | A | 31 | 12 | 12 | 0 | 0 |
| 12 | B | 31 | 12 | 12 | 0 | 0 |
| 12 | C | 31 | 12 | 12 | 0 | 0 |
| 12 | F | 31 | 12 | 12 | 1 | 0 |
| 13 | A | 1 | 0 | 0 | 0 | 0 |
| 13 | B | 1 | 0 | 0 | 0 | 0 |
| 13 | C | 1 | 0 | 0 | 0 | 0 |
| 13 | D | 1 | 0 | 0 | 0 | 0 |
| 13 | F | 1 | 0 | 0 | 0 | 0 |
| 14 | D | 27 | 12 | 12 | 0 | 0 |
| 15 | E | 25 | 36 | 36 | 2 | 0 |
| All | All | 40604 | 41207 | 41207 | 278 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 7:M:147:VAL:HG12 | 7:M:191:TYR:HB3 | 1.58 | 0.84 |
| 9:W:84:LEU:HD22 | 9:X:53:VAL:HG11 | 1.60 | 0.81 |
| 1:C:452:ILE:HG21 | 1:C:511:LEU:HD11 | 1.65 | 0.79 |

Continued on next page...

Continued from previous page...

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|-----------------|--------------------------|-------------------|
| 1:C:227:ASN:OD1 | 1:C:230:ARG:NH2 | 2.17 | 0.78 |
| 1:A:224:ARG:NH2 | 2:D:340:ASP:OD1 | 2.17 | 0.77 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1 | A | 522/561 (93%) | 499 (96%) | 23 (4%) | 0 | 100 | 100 |
| 1 | B | 522/561 (93%) | 493 (94%) | 29 (6%) | 0 | 100 | 100 |
| 1 | C | 525/561 (94%) | 497 (95%) | 28 (5%) | 0 | 100 | 100 |
| 2 | D | 485/494 (98%) | 460 (95%) | 23 (5%) | 2 (0%) | 34 | 69 |
| 2 | E | 485/494 (98%) | 458 (94%) | 26 (5%) | 1 (0%) | 47 | 80 |
| 2 | F | 485/494 (98%) | 462 (95%) | 21 (4%) | 2 (0%) | 34 | 69 |
| 3 | G | 301/306 (98%) | 294 (98%) | 7 (2%) | 0 | 100 | 100 |
| 4 | H | 158/176 (90%) | 148 (94%) | 10 (6%) | 0 | 100 | 100 |
| 5 | I | 64/76 (84%) | 59 (92%) | 5 (8%) | 0 | 100 | 100 |
| 6 | J | 168/192 (88%) | 164 (98%) | 4 (2%) | 0 | 100 | 100 |
| 6 | K | 168/192 (88%) | 167 (99%) | 1 (1%) | 0 | 100 | 100 |
| 6 | L | 168/192 (88%) | 165 (98%) | 3 (2%) | 0 | 100 | 100 |
| 7 | M | 241/267 (90%) | 231 (96%) | 9 (4%) | 1 (0%) | 34 | 69 |
| 8 | N | 47/103 (46%) | 46 (98%) | 1 (2%) | 0 | 100 | 100 |
| 9 | O | 79/104 (76%) | 78 (99%) | 1 (1%) | 0 | 100 | 100 |
| 9 | P | 79/104 (76%) | 78 (99%) | 1 (1%) | 0 | 100 | 100 |
| 9 | Q | 79/104 (76%) | 78 (99%) | 1 (1%) | 0 | 100 | 100 |
| 9 | R | 79/104 (76%) | 77 (98%) | 2 (2%) | 0 | 100 | 100 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 9 | S | 79/104 (76%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 9 | T | 79/104 (76%) | 78 (99%) | 1 (1%) | 0 | 100 | 100 |
| 9 | U | 79/104 (76%) | 78 (99%) | 1 (1%) | 0 | 100 | 100 |
| 9 | V | 79/104 (76%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 9 | W | 79/104 (76%) | 77 (98%) | 2 (2%) | 0 | 100 | 100 |
| 9 | X | 79/104 (76%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 10 | h | 47/476 (10%) | 46 (98%) | 1 (2%) | 0 | 100 | 100 |
| 11 | c | 53/169 (31%) | 53 (100%) | 0 | 0 | 100 | 100 |
| All | All | 5229/6354 (82%) | 5014 (96%) | 209 (4%) | 6 (0%) | 54 | 84 |

5 of 6 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | D | 289 | VAL |
| 2 | F | 289 | VAL |
| 2 | D | 288 | ALA |
| 2 | F | 288 | ALA |
| 7 | M | 148 | ALA |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|------------|----------|-------------|-----|
| 1 | A | 450/474 (95%) | 450 (100%) | 0 | 100 | 100 |
| 1 | B | 450/474 (95%) | 450 (100%) | 0 | 100 | 100 |
| 1 | C | 452/474 (95%) | 452 (100%) | 0 | 100 | 100 |
| 2 | D | 397/400 (99%) | 397 (100%) | 0 | 100 | 100 |
| 2 | E | 397/400 (99%) | 397 (100%) | 0 | 100 | 100 |
| 2 | F | 397/400 (99%) | 397 (100%) | 0 | 100 | 100 |
| 3 | G | 261/264 (99%) | 260 (100%) | 1 (0%) | 91 | 96 |
| 4 | H | 131/146 (90%) | 131 (100%) | 0 | 100 | 100 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|-------------|----------|-------------|-----|
| 5 | I | 58/66 (88%) | 58 (100%) | 0 | 100 | 100 |
| 6 | J | 132/151 (87%) | 132 (100%) | 0 | 100 | 100 |
| 6 | K | 132/151 (87%) | 132 (100%) | 0 | 100 | 100 |
| 6 | L | 132/151 (87%) | 132 (100%) | 0 | 100 | 100 |
| 7 | M | 202/221 (91%) | 202 (100%) | 0 | 100 | 100 |
| 8 | N | 44/87 (51%) | 44 (100%) | 0 | 100 | 100 |
| 9 | O | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | P | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | Q | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | R | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | S | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | T | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | U | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | V | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | W | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 9 | X | 58/76 (76%) | 58 (100%) | 0 | 100 | 100 |
| 10 | h | 44/414 (11%) | 44 (100%) | 0 | 100 | 100 |
| 11 | c | 52/137 (38%) | 52 (100%) | 0 | 100 | 100 |
| All | All | 4311/5170 (83%) | 4310 (100%) | 1 (0%) | 100 | 100 |

All (1) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | G | 19 | LYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 160 | ASN |
| 3 | G | 239 | ASN |

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 14 | ADP | D | 601 | 13 | 24,29,29 | 0.97 | 1 (4%) | 29,45,45 | 1.46 | 4 (13%) |
| 15 | TRT | E | 600 | - | 25,25,25 | 0.61 | 0 | 33,33,33 | 0.71 | 0 |
| 12 | ATP | F | 601 | 13 | 26,33,33 | 1.01 | 2 (7%) | 31,52,52 | 1.65 | 7 (22%) |
| 12 | ATP | C | 601 | 13 | 26,33,33 | 0.88 | 0 | 31,52,52 | 1.62 | 5 (16%) |
| 12 | ATP | B | 601 | 13 | 26,33,33 | 0.86 | 0 | 31,52,52 | 1.49 | 4 (12%) |
| 12 | ATP | A | 601 | 13 | 26,33,33 | 0.82 | 0 | 31,52,52 | 1.55 | 5 (16%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|------------|---------|
| 14 | ADP | D | 601 | 13 | - | 0/12/32/32 | 0/3/3/3 |
| 15 | TRT | E | 600 | - | - | 8/23/23/23 | 0/1/1/1 |
| 12 | ATP | F | 601 | 13 | - | 4/18/38/38 | 0/3/3/3 |
| 12 | ATP | C | 601 | 13 | - | 3/18/38/38 | 0/3/3/3 |
| 12 | ATP | B | 601 | 13 | - | 2/18/38/38 | 0/3/3/3 |
| 12 | ATP | A | 601 | 13 | - | 0/18/38/38 | 0/3/3/3 |

All (3) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 12 | F | 601 | ATP | C2'-C1' | -2.55 | 1.49 | 1.53 |
| 14 | D | 601 | ADP | C5-C4 | 2.18 | 1.46 | 1.40 |
| 12 | F | 601 | ATP | C5-C4 | 2.04 | 1.46 | 1.40 |

The worst 5 of 25 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|-------|-------------|----------|
| 12 | C | 601 | ATP | PB-O3B-PG | -4.38 | 117.81 | 132.83 |
| 12 | F | 601 | ATP | PB-O3B-PG | -4.00 | 119.09 | 132.83 |
| 12 | B | 601 | ATP | PB-O3B-PG | -3.94 | 119.31 | 132.83 |
| 12 | A | 601 | ATP | PB-O3B-PG | -3.73 | 120.02 | 132.83 |
| 12 | A | 601 | ATP | N3-C2-N1 | -3.58 | 123.08 | 128.68 |

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-----------------|
| 12 | F | 601 | ATP | C5'-O5'-PA-O1A |
| 12 | F | 601 | ATP | C5'-O5'-PA-O3A |
| 15 | E | 600 | TRT | O15-C16-C17-O18 |
| 12 | C | 601 | ATP | PG-O3B-PB-O1B |
| 15 | E | 600 | TRT | C5-C6-C9-C10 |


There are no ring outliers.


2 monomers are involved in 3 short contacts:


| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 15 | E | 600 | TRT | 2 | 0 |
| 12 | F | 601 | ATP | 1 | 0 |


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10471. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 220




Y Index: 220




Z Index: 220


6.2.2 Raw map



X Index: 220



Y Index: 220




Z Index: 220


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 295




Y Index: 198




Z Index: 294


6.3.2 Raw map



X Index: 291



Y Index: 207



Z Index: 281

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.025. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_10471_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 291 nm³; this corresponds to an approximate mass of 263 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.329 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.329 Å⁻¹

8.2 Resolution estimates [i](#)


| Resolution estimate (Å) | Estimation criterion (FSC cut-off) | | |
|---------------------------|------------------------------------|------|----------|
| | 0.143 | 0.5 | Half-bit |
| Reported by author | 3.04 | - | - |
| Author-provided FSC curve | - | - | - |
| Unmasked-calculated* | 3.62 | 5.93 | 3.69 |

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.62 differs from the reported value 3.04 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-10471 and PDB model 6TDY. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.025 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.025).























































9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.025) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|--|--|
| All |  0.7332 |  0.4000 |
| A |  0.8311 |  0.4500 |
| B |  0.8402 |  0.4650 |
| C |  0.8192 |  0.4360 |
| D |  0.8053 |  0.4380 |
| E |  0.8263 |  0.4640 |
| F |  0.8236 |  0.4560 |
| G |  0.7821 |  0.4250 |
| H |  0.6710 |  0.3690 |
| I |  0.7203 |  0.3760 |
| J |  0.7880 |  0.3810 |
| K |  0.7841 |  0.3840 |
| L |  0.7381 |  0.3300 |
| M |  0.6466 |  0.3380 |
| N |  0.6768 |  0.3870 |
| O |  0.3513 |  0.2240 |
| P |  0.4052 |  0.2470 |
| Q |  0.4452 |  0.2730 |
| R |  0.4626 |  0.2910 |
| S |  0.4539 |  0.2760 |
| T |  0.4626 |  0.2790 |
| U |  0.4974 |  0.2690 |
| V |  0.4504 |  0.2230 |
| W |  0.4157 |  0.2310 |
| X |  0.3496 |  0.2100 |
| c |  0.4702 |  0.2570 |
| h |  0.3427 |  0.2660 |

