



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2024 – 04:28 pm GMT

PDB ID : 6TG4
Title : Tankyrase 2 in complex with an inhibitor (OM-1700)
Authors : Sowa, S.T.; Lehtio, L.
Deposited on : 2019-11-15
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	240	
1	BBB	240	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6532 atoms, of which 3177 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tankyrase-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	AAA	206	3215	1032	1576	300	297	10	104	0	0
1	BBB	200	3166	1015	1555	295	291	10	98	0	0

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	923	MET	-	initiating methionine	UNP Q9H2K2
AAA	924	HIS	-	expression tag	UNP Q9H2K2
AAA	925	HIS	-	expression tag	UNP Q9H2K2
AAA	926	HIS	-	expression tag	UNP Q9H2K2
AAA	927	HIS	-	expression tag	UNP Q9H2K2
AAA	928	HIS	-	expression tag	UNP Q9H2K2
AAA	929	HIS	-	expression tag	UNP Q9H2K2
AAA	930	SER	-	expression tag	UNP Q9H2K2
AAA	931	SER	-	expression tag	UNP Q9H2K2
AAA	932	GLY	-	expression tag	UNP Q9H2K2
AAA	933	VAL	-	expression tag	UNP Q9H2K2
AAA	934	ASP	-	expression tag	UNP Q9H2K2
AAA	935	LEU	-	expression tag	UNP Q9H2K2
AAA	936	GLY	-	expression tag	UNP Q9H2K2
AAA	937	THR	-	expression tag	UNP Q9H2K2
AAA	938	GLU	-	expression tag	UNP Q9H2K2
AAA	939	ASN	-	expression tag	UNP Q9H2K2
AAA	940	LEU	-	expression tag	UNP Q9H2K2
AAA	941	TYR	-	expression tag	UNP Q9H2K2
AAA	942	PHE	-	expression tag	UNP Q9H2K2
AAA	943	GLN	-	expression tag	UNP Q9H2K2
AAA	944	SER	-	expression tag	UNP Q9H2K2
AAA	945	MET	-	expression tag	UNP Q9H2K2
BBB	923	MET	-	initiating methionine	UNP Q9H2K2
BBB	924	HIS	-	expression tag	UNP Q9H2K2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
BBB	925	HIS	-	expression tag	UNP Q9H2K2
BBB	926	HIS	-	expression tag	UNP Q9H2K2
BBB	927	HIS	-	expression tag	UNP Q9H2K2
BBB	928	HIS	-	expression tag	UNP Q9H2K2
BBB	929	HIS	-	expression tag	UNP Q9H2K2
BBB	930	SER	-	expression tag	UNP Q9H2K2
BBB	931	SER	-	expression tag	UNP Q9H2K2
BBB	932	GLY	-	expression tag	UNP Q9H2K2
BBB	933	VAL	-	expression tag	UNP Q9H2K2
BBB	934	ASP	-	expression tag	UNP Q9H2K2
BBB	935	LEU	-	expression tag	UNP Q9H2K2
BBB	936	GLY	-	expression tag	UNP Q9H2K2
BBB	937	THR	-	expression tag	UNP Q9H2K2
BBB	938	GLU	-	expression tag	UNP Q9H2K2
BBB	939	ASN	-	expression tag	UNP Q9H2K2
BBB	940	LEU	-	expression tag	UNP Q9H2K2
BBB	941	TYR	-	expression tag	UNP Q9H2K2
BBB	942	PHE	-	expression tag	UNP Q9H2K2
BBB	943	GLN	-	expression tag	UNP Q9H2K2
BBB	944	SER	-	expression tag	UNP Q9H2K2
BBB	945	MET	-	expression tag	UNP Q9H2K2

- Molecule 2 is N-[3-[5-(5-ethoxypyridin-2-yl)-4-(2-fluorophenyl)-1,2,4-triazol-3-yl]cyclobutyl]pyridine-2-carboxamide (three-letter code: N8K) (formula: C₂₅H₂₃FN₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	AAA	1	Total	C	F	H	N	O	0	0
			57	25	1	23	6	2		
2	BBB	1	Total	C	F	H	N	O	0	0
			57	25	1	23	6	2		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	1	Total	Zn	0	0
			1	1		
3	BBB	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	20	Total	O	0	0
			20	20		
4	BBB	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tankyrase-2



- Molecule 1: Tankyrase-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.68Å 77.90Å 149.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.46 – 2.76 37.43 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.46-2.76) 99.6 (37.43-2.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.230 , 0.281 0.234 , 0.281	Depositor DCC
R_{free} test set	670 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtrriage
Anisotropy	0.449	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 30.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6532	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.0017e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: N8K, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.66	0/1682	0.77	0/2268
1	BBB	0.64	0/1652	0.77	0/2221
All	All	0.65	0/3334	0.77	0/4489

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1639	1576	1550	4	0
1	BBB	1611	1555	1534	9	0
2	AAA	34	23	0	0	0
2	BBB	34	23	0	1	0
3	AAA	1	0	0	0	0
3	BBB	1	0	0	0	0
4	AAA	20	0	0	0	0
4	BBB	15	0	0	1	0
All	All	3355	3177	3084	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:979:HIS:NE2	1:BBB:1066:SER:OG	2.31	0.58
1:BBB:976:VAL:O	1:BBB:977:ARG:NH1	2.45	0.49
1:AAA:975:THR:HB	1:AAA:1064:ASN:HA	1.96	0.48
1:BBB:975:THR:HB	1:BBB:1064:ASN:HA	1.96	0.47
1:BBB:1037:ASN:H	1:BBB:1037:ASN:ND2	2.12	0.47
1:BBB:1143:ARG:HB3	1:BBB:1145:GLU:OE1	2.17	0.45
1:BBB:979:HIS:CD2	1:BBB:1066:SER:HG	2.31	0.45
1:BBB:1069:ASN:O	1:BBB:1072:VAL:HG23	2.19	0.42
1:AAA:976:VAL:O	1:AAA:977:ARG:NH1	2.52	0.42
1:BBB:1079:THR:N	4:BBB:1302:HOH:O	2.53	0.41
1:BBB:1095:GLN:HA	1:BBB:1153:ILE:O	2.20	0.41
1:AAA:1095:GLN:HA	1:AAA:1153:ILE:O	2.21	0.41
1:AAA:956:ILE:HG23	1:AAA:999:LYS:HB3	2.01	0.41
2:BBB:1201:N8K:C20	2:BBB:1201:N8K:C10	3.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	202/240 (84%)	196 (97%)	6 (3%)	0	100	100
1	BBB	194/240 (81%)	189 (97%)	5 (3%)	0	100	100
All	All	396/480 (82%)	385 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	171/204 (84%)	168 (98%)	3 (2%)	59	75
1	BBB	169/204 (83%)	160 (95%)	9 (5%)	22	38
All	All	340/408 (83%)	328 (96%)	12 (4%)	36	56

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	956	ILE
1	AAA	990	ASN
1	AAA	1079	THR
1	BBB	967	SER
1	BBB	990	ASN
1	BBB	1004	LYS
1	BBB	1037	ASN
1	BBB	1051	ILE
1	BBB	1067	LYS
1	BBB	1085	LYS
1	BBB	1110	PHE
1	BBB	1145	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	N8K	AAA	1201	-	36,38,38	2.99	8 (22%)	39,53,53	1.52	8 (20%)
2	N8K	BBB	1201	-	36,38,38	2.77	9 (25%)	39,53,53	1.69	8 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	N8K	AAA	1201	-	-	8/19/31/31	0/5/5/5
2	N8K	BBB	1201	-	-	12/19/31/31	0/5/5/5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	1201	N8K	N2-N1	-10.42	1.17	1.37
2	AAA	1201	N8K	C19-N5	-9.77	1.34	1.45
2	BBB	1201	N8K	C19-N5	-9.33	1.35	1.45
2	BBB	1201	N8K	N2-N1	-8.08	1.21	1.37
2	AAA	1201	N8K	C6-C7	-7.44	1.33	1.47
2	BBB	1201	N8K	C6-C7	-7.11	1.34	1.47
2	BBB	1201	N8K	C13-C12	-4.31	1.40	1.50
2	AAA	1201	N8K	C13-C12	-4.12	1.40	1.50
2	BBB	1201	N8K	C17-N4	3.60	1.42	1.34
2	AAA	1201	N8K	C17-N4	3.45	1.41	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	1201	N8K	C10-C9	-3.08	1.50	1.56
2	AAA	1201	N8K	C25-N6	3.02	1.40	1.34
2	BBB	1201	N8K	C8-N5	-2.99	1.32	1.37
2	BBB	1201	N8K	C18-C9	-2.94	1.51	1.56
2	BBB	1201	N8K	C25-N6	2.90	1.40	1.34
2	AAA	1201	N8K	C8-N5	-2.23	1.33	1.37
2	BBB	1201	N8K	C8-C9	2.14	1.53	1.50

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	BBB	1201	N8K	C19-N5-C7	4.44	130.54	124.57
2	AAA	1201	N8K	F1-C24-C19	3.80	123.97	121.04
2	BBB	1201	N8K	C14-C13-N4	-3.73	117.63	122.72
2	BBB	1201	N8K	C17-N4-C13	3.71	121.74	116.93
2	AAA	1201	N8K	C17-N4-C13	3.34	121.27	116.93
2	BBB	1201	N8K	C14-C13-C12	2.92	124.32	119.57
2	BBB	1201	N8K	C25-N6-C6	2.85	121.66	117.90
2	AAA	1201	N8K	C14-C13-N4	-2.76	118.97	122.72
2	AAA	1201	N8K	C10-C9-C8	-2.68	109.62	119.20
2	BBB	1201	N8K	C15-C14-C13	2.68	121.87	118.63
2	AAA	1201	N8K	C14-C13-C12	2.51	123.65	119.57
2	BBB	1201	N8K	C10-C9-C8	-2.41	110.58	119.20
2	AAA	1201	N8K	C25-N6-C6	2.32	120.95	117.90
2	AAA	1201	N8K	C16-C17-N4	-2.12	119.97	123.43
2	BBB	1201	N8K	F1-C24-C19	2.09	122.65	121.04
2	AAA	1201	N8K	C7-C6-N6	2.03	120.11	116.35

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	1201	N8K	C5-C6-C7-N1
2	AAA	1201	N8K	N6-C6-C7-N1
2	AAA	1201	N8K	N6-C6-C7-N5
2	BBB	1201	N8K	C5-C6-C7-N1
2	BBB	1201	N8K	N6-C6-C7-N1
2	BBB	1201	N8K	N6-C6-C7-N5
2	AAA	1201	N8K	N3-C12-C13-C14
2	AAA	1201	N8K	O2-C12-C13-N4
2	AAA	1201	N8K	O2-C12-C13-C14
2	AAA	1201	N8K	N3-C12-C13-N4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	BBB	1201	N8K	O2-C12-C13-C14
2	BBB	1201	N8K	N3-C12-C13-C14
2	BBB	1201	N8K	O2-C12-C13-N4
2	BBB	1201	N8K	N3-C12-C13-N4
2	BBB	1201	N8K	C4-C3-O1-C2
2	BBB	1201	N8K	C25-C3-O1-C2
2	BBB	1201	N8K	C1-C2-O1-C3
2	AAA	1201	N8K	C5-C6-C7-N5
2	BBB	1201	N8K	C5-C6-C7-N5
2	BBB	1201	N8K	C18-C11-N3-C12

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	1201	N8K	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	206/240 (85%)	0.27	5 (2%) 59 68	30, 45, 68, 86	0
1	BBB	200/240 (83%)	0.05	3 (1%) 73 81	35, 48, 76, 95	0
All	All	406/480 (84%)	0.16	8 (1%) 65 73	30, 46, 74, 95	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	1131	VAL	3.2
1	BBB	1051	ILE	3.1
1	AAA	1115	MET	2.9
1	BBB	1110	PHE	2.8
1	AAA	980	ARG	2.3
1	AAA	1132	ASN	2.1
1	BBB	1052	GLY	2.1
1	AAA	1042	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	N8K	AAA	1201	34/34	0.94	0.19	34,37,47,48	0
2	N8K	BBB	1201	34/34	0.95	0.20	37,44,52,53	0
3	ZN	AAA	1202	1/1	0.99	0.10	41,41,41,41	0
3	ZN	BBB	1202	1/1	0.99	0.12	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.