

Full wwPDB X-ray Structure Validation Report (i)

Feb 7, 2024 – 02:57 PM EST

PDB ID : 8TJ4 Title : CRYSTAL STRUCTURE OF THE A/Bangkok/1/1979(H3N2) IN-FLUENZA VIRUS HEMAGGLUTININ WITH HUMAN RECEPTOR ANA-LOG 6'-SLNLN Authors : Wu, N.C.; Zhu, X.; Wilson, I.A. Deposited on : 2023-07-20

Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution
IVIETIC	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	323	92%	6% •
1	С	323	89%	8% •
1	Е	323	% 93%	
1	G	323	92%	6% ·



Chain Length Quality of chain Mol .% 2В 174. . 95% 2% 2D 6% • 17493% 2% 2F 174. . 96% 2% Η 25%• 17494% Ι 3 2100% 23 \mathbf{L} 50% 50% 2Ο 3 100% \mathbf{S} 3 2100% Т 23 50% 50% U $\mathbf{2}$ 3 100% V 23 50% 50% 2W 3 50% 50% 3 Х $\mathbf{2}$ 50% 50% J 3 4 67% 33% 3 М 4 33% 67% 4 Ν 3 100% Κ 5540% 60% Q 5540% 20% 40% R 5560% 40% Р 6 4 50% 50% Υ 6 4 25% 50% 25% Ζ 7 633% 50% 17% 3 8 \mathbf{a} 33% 67% 8 3 \mathbf{c} 100% 9 b 250% 50% Continued on next page...



	Mol	Chain	Chain 🛛	Length	Quality	of chain
9 d 2	Q	d	d	2	5.0%	E 0%/



2 Entry composition (i)

There are 14 unique types of molecules in this entry. The entry contains 18202 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	217	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	517	2460	1543	434	471	12	0	0	U
1	C	317	Total	С	Ν	0	S	0	0	0
	U		2460	1543	434	471	12			0
1	F	217	Total	С	Ν	0	S	0	0	0
		317	2460	1543	434	471	12		0	0
1	1 G	G 317	Total	С	Ν	0	S	0	0	0
			2460	1543	434	471	12		0	0

• Molecule 1 is a protein called Hemagglutinin HA1 chain.

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	7	ALA	-	expression tag	UNP P03441
А	8	ASP	-	expression tag	UNP P03441
А	9	PRO	-	expression tag	UNP P03441
А	10	GLY	-	expression tag	UNP P03441
А	294	PHE	UNK	conflict	UNP P03441
С	7	ALA	-	expression tag	UNP P03441
С	8	ASP	-	expression tag	UNP P03441
С	9	PRO	-	expression tag	UNP P03441
С	10	GLY	-	expression tag	UNP P03441
С	294	PHE	UNK	conflict	UNP P03441
Е	7	ALA	-	expression tag	UNP P03441
Е	8	ASP	-	expression tag	UNP P03441
Е	9	PRO	-	expression tag	UNP P03441
Е	10	GLY	-	expression tag	UNP P03441
E	294	PHE	UNK	conflict	UNP P03441
G	7	ALA	-	expression tag	UNP P03441
G	8	ASP	-	expression tag	UNP P03441
G	9	PRO	-	expression tag	UNP P03441
G	10	GLY	-	expression tag	UNP P03441
G	294	PHE	UNK	conflict	UNP P03441



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
0	D	179	Total	С	Ν	0	\mathbf{S}	0	0	0
	Б 172	172	1386	860	243	277	6	0	0	0
2	Л	179	Total	С	Ν	0	S	0	0	0
	D	172	1386	860	243	277	6	0	0	0
9	Б	179	Total	С	Ν	0	S	0	0	0
	2 F	172	1392	863	246	277	6		0	0
9	2 H	179	Total	С	Ν	0	S	0	0	0
		172	1392	863	246	277	6			0

• Molecule 2 is a protein called Hemagglutinin HA2 chain.

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	18	VAL	UNK	conflict	UNP P03441
В	19	ASP	UNK	conflict	UNP P03441
D	18	VAL	UNK	conflict	UNP P03441
D	19	ASP	UNK	conflict	UNP P03441
F	18	VAL	UNK	conflict	UNP P03441
F	19	ASP	UNK	conflict	UNP P03441
Н	18	VAL	UNK	conflict	UNP P03441
Н	19	ASP	UNK	conflict	UNP P03441

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	Ι	2	Total C N O 28 16 2 10	0	0	0
3	L	2	Total C N O 28 16 2 10	0	0	0
3	О	2	Total C N O 28 16 2 10	0	0	0
3	S	2	Total C N O 28 16 2 10	0	0	0
3	Т	2	Total C N O 28 16 2 10	0	0	0
3	U	2	Total C N O 28 16 2 10	0	0	0



0 0									
Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace			
3	V	2	Total C N O 28 16 2 10	0	0	0			
3	W	2	Total C N O 28 16 2 10	0	0	0			
3	Х	2	Total C N O 28 16 2 10	0	0	0			

• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	J	3	Total C N O 39 22 2 15	0	0	0
4	М	3	Total C N O 39 22 2 15	0	0	0
4	Ν	3	Total C N O 39 22 2 15	0	0	0

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	K	5	Total C N O 61 34 2 25	0	0	0
5	Q	5	Total C N O 61 34 2 25	0	0	0
5	R	5	Total C N O 61 34 2 25	0	0	0

• Molecule 6 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-be ta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyr anose.





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
6	Р	4	Total 48	C 28	N 2	O 18	0	0	0
6	Y	4	Total 48	C 28	N 2	0 18	0	0	0

• Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyran ose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
7	Z	6	Total 72	C 40	N 2	O 30	0	0	0

• Molecule 8 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galacto pyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
8	a	3	Total 46	C 25	N 2	O 19	0	0	0
8	с	3	Total 46	$\begin{array}{c} \mathrm{C} \\ \mathrm{25} \end{array}$	N 2	O 19	0	0	0

• Molecule 9 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galacto pyranose.





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
9	b	2	Total 32	C N 17 1	O 14	0	0	0
9	d	2	Total 32	C N 17 1	0 14	0	0	0

• Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	1	Total C N O 14 8 1 5	0	0
10	В	1	Total C N O 14 8 1 5	0	0
10	D	1	Total C N O 14 8 1 5	0	0
10	Е	1	Total C N O 14 8 1 5	0	0
10	F	1	Total C N O 14 8 1 5	0	0
10	G	1	Total C N O 14 8 1 5	0	0
10	Н	1	Total C N O 14 8 1 5	0	0

• Molecule 11 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	В	1	Total C O	0	0
	D	1	$13 \ 8 \ 5$	0	0
11	Б	1	Total C O	0	0
11	Г	1	$13 \ 8 \ 5$	0	

• Molecule 12 is 2-(2-METHOXYETHOXY)ETHANOL (three-letter code: PG0) (formula: $C_5H_{12}O_3$).



Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
12	В	1	Total 8	C 5	O 3	0	0



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	Н	1	$\begin{array}{cc} {\rm Total} & {\rm C} \\ 8 & 5 \end{array}$	O 3	0	0

• Molecule 13 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	Е	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0
13	F	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0
13	F	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0

• Molecule 14 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	А	233	Total O 233 233	0	0
14	В	207	Total O 207 207	0	0
14	С	234	Total O 234 234	0	0
14	D	216	Total O 216 216	0	0
14	Е	227	Total O 227 227	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	F	214	Total O 214 214	0	0
14	G	237	Total O 237 237	0	0
14	Н	201	Total O 201 201	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 6% • 92% • Molecule 1: Hemagglutinin HA1 chain Chain C: 89% 8% LYS GLN THR • Molecule 1: Hemagglutinin HA1 chain Chain E: 93% AL/ ASF • Molecule 1: Hemagglutinin HA1 chain Chain G: 92% 6% • Molecule 2: Hemagglutinin HA2 chain Chain B: 95%
- Molecule 1: Hemagglutinin HA1 chain



• Molecule 2:	Hemagglutinin HA2 chain	
Chain D:	93%	6% ·
G1 K51 V55 156 E57 K58 T59 N60	K62 K161 F111 Q172 LVS LVS	
• Molecule 2:	Hemagglutinin HA2 chain	
Chain F:	96%	
61 D19 E57 K58 K58 K58 K58 K58 K58 K58 K58 K58 K58	LVS	
• Molecule 2:	Hemagglutinin HA2 chain	
Chain H:	94%	5%•
61 64 751 754 755 755	LYS	
• Molecule 3: opyranose	2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-ace	etamido-2-deoxy-beta-D-gluc
Chain I:	100%	
NAG1 NAG2		
• Molecule 3: opyranose	2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-ace	etamido-2-deoxy-beta-D-gluc
Chain L:	50% 50%	
NAG2 NAG2		
• Molecule 3: opyranose	2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-ace	etamido-2-deoxy-beta-D-gluc
Chain O:	100%	
NAG1 NAG2		
• Molecule 3: opyranose	2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-ace	etamido-2-deoxy-beta-D-gluc

Chain S:

100%



NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain T:	50%	50%	
NAG1 NAG2			
• Molecule : opyranose	3: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain U:	10	00%	
NAG1 NAG2			
• Molecule 3 opyranose	3: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain V:	50%	50%	
NAG1 NAG2			
• Molecule : opyranose	3: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain W:	50%	50%	•
NAG1 NAG2			
• Molecule : opyranose	3: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain X:	50%	50%	
NAG2 NAG2			

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:	67%	33%
NAG1 NAG2 BMA3		



• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

33%

Chain M:

NAG1 NAG2 BMA3

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:	100%	
NAG1 BMA3 BMA3		

67%

 \bullet Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:	40%	60%
NAG1 BMA3 MAN4 MAN5 MAN5		

 \bullet Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose nose

α · α			
Chain Q:	40%	20%	40%
•			

NAG1 NAG2 BMA3 MAN4 MAN5

 \bullet Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose nose

Chain R:	60%	40%
NAG1 NAG2 MAN4 MAN5 MAN5		

 $\bullet \ Molecule \ 6: \ alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)] \\ 2-acetamido-2-deoxy-beta-D-glucopyranose \\ (1-6)] \\ 2-acetamido-2-deoxy-beta-D-glucopyranose \\ (1-6)] \\ (1-6)$

50%

Chain P: 50%

NAG1 FUC2 NAG3 FUC4



 $\bullet \ Molecule \ 6: \ alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)] \\ 2-acetamido-2-deoxy-beta-D-glucopyranose \\ (1-6)] \\ (1$

Chain Y:	25%	50%	25%
NAG1 FUC2 NAG3 FUC4			

• Molecule 7: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Z:	33%	50%	17%
NAG1 NAG2 BMA3 MAN4 MAN5 MAN6 MAN6			

• Molecule 8: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-de
oxy-beta-D-glucopyranose

Chain a:	33%	67%
NAG1 GAL2 SIA3		

• Molecule 8: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-de
oxy-beta-D-glucopyranose

Chain c:	1	100%
NAG1 GAL2 SIA3		
• Molecule 9: 1	N-acetyl-alpha-neuraminic	acid-(2-6)-beta-D-galactopyranose
Chain b:	50%	50%
GAL1 SIA2		
• Molecule 9: 2	N-acetyl-alpha-neuraminic	acid-(2-6)-beta-D-galactopyranose
Chain d:	50%	50%
GAL1 SIA2		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	Н 3	Depositor
Cell constants	100.66Å 100.66Å 687.90Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	45.23 - 1.95	Depositor
Resolution (A)	45.23 - 1.95	EDS
% Data completeness	99.2 (45.23-1.95)	Depositor
(in resolution range)	99.5~(45.23-1.95)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.13	Depositor
$< I/\sigma(I) > 1$	$1.80 (at 1.95 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
D D.	0.189 , 0.223	Depositor
Π, Π_{free}	0.190 , 0.222	DCC
R_{free} test set	9453 reflections (5.01%)	wwPDB-VP
Wilson B-factor $(Å^2)$	23.9	Xtriage
Anisotropy	0.370	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 39.4	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.477 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18202	wwPDB-VP
Average B, all atoms $(Å^2)$	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 41.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5092e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, PG4, MAN, PG0, FUC, NAG, SIA, BMA, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Chain		Bond lengths		Bond angles	
			# Z > 5	RMSZ	# Z > 5
1	А	0.36	0/2518	0.61	0/3423
1	С	0.37	0/2518	0.62	0/3423
1	Е	0.36	0/2518	0.60	0/3423
1	G	0.38	0/2518	0.62	0/3423
2	В	0.42	0/1410	0.63	0/1896
2	D	0.47	0/1410	0.73	0/1896
2	F	0.43	0/1416	0.66	0/1903
2	Н	0.50	0/1416	0.74	0/1903
All	All	0.40	0/15724	0.64	0/21290

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2460	0	2405	9	0
1	С	2460	0	2405	12	0
1	Е	2460	0	2405	8	0
1	G	2460	0	2404	9	0
2	В	1386	0	1300	6	0



	Chain	Non H	puye	II(addad)	Clashag	Summe Clashes
	D	1900	H(model)	H(added)	Clasnes	Symm-Clasnes
2		1380	0	1300	5	0
2		1392	0	1311	3 C	0
2	Н	1392	0	1311	0	0
<u>3</u>	l T	28	0	25	0	0
3	L	28	0	25	0	0
3	0	28	0	25	0	0
3	S	28	0	25	1	0
3	Т	28	0	25	0	0
3	U	28	0	25	0	0
3	V	28	0	25	0	0
3	W	28	0	25	1	0
3	X	28	0	25	0	0
4	J	39	0	34	0	0
4	М	39	0	34	0	0
4	N	39	0	34	0	0
5	K	61	0	52	0	0
5	Q	61	0	52	1	0
5	R	61	0	52	0	0
6	Р	48	0	43	0	0
6	Y	48	0	43	2	0
7	Z	72	0	61	1	0
8	a	46	0	40	0	0
8	с	46	0	40	0	0
9	b	32	0	28	0	0
9	d	32	0	28	0	0
10	А	14	0	13	0	0
10	В	14	0	13	0	0
10	D	14	0	13	0	0
10	Е	14	0	13	0	0
10	F	14	0	13	0	0
10	G	14	0	13	0	0
10	Н	14	0	13	0	0
11	В	13	0	18	2	0
11	F	13	0	18	1	0
12	В	8	0	12	1	0
12	Н	8	0	12	0	0
13	Е	7	0	10	2	0
13	F	14	0	20	3	0
14	А	233	0	0	1	0
14	В	207	0	0	2	0
14	С	234	0	0	1	0
14	D	216	0	0	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	Е	227	0	0	1	0
14	F	214	0	0	1	0
14	G	237	0	0	0	0
14	Н	201	0	0	0	0
All	All	18202	0	15788	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:G:177:LEU:HB2	1:G:260:ILE:HD11	1.74	0.69
2:F:71:SER:HB3	13:F:204:PEG:H31	1.73	0.69
1:A:177:LEU:HB2	1:A:260:ILE:HD11	1.76	0.66
1:C:177:LEU:HB2	1:C:260:ILE:HD11	1.79	0.65
1:G:222:TRP:CZ2	1:G:225:GLY:HA2	2.33	0.63
1:A:222:TRP:CZ2	1:A:225:GLY:HA2	2.36	0.61
1:C:222:TRP:CZ2	1:C:225:GLY:HA2	2.38	0.59
1:A:102:VAL:HG22	1:A:232:ILE:HB	1.86	0.58
1:G:126:ASN:OD1	1:G:128:THR:HG23	2.02	0.58
14:E:615:HOH:O	13:F:203:PEG:H41	2.06	0.55
13:E:402:PEG:H42	13:F:203:PEG:H11	1.88	0.55
2:B:39:LYS:HE3	14:B:482:HOH:O	2.08	0.54
1:E:102:VAL:HG22	1:E:232:ILE:HB	1.89	0.54
1:G:77:ASP:OD2	1:G:141:ARG:NH1	2.42	0.53
2:D:51:LYS:O	2:D:55:VAL:HG23	2.09	0.52
1:C:77:ASP:OD2	1:C:141:ARG:NH1	2.43	0.51
2:H:51:LYS:O	2:H:55:VAL:HG23	2.11	0.50
13:E:402:PEG:H11	3:S:1:NAG:H83	1.94	0.50
1:E:195:TYR:O	1:E:196:VAL:HG22	2.11	0.49
2:D:1:GLY:HA2	14:D:310:HOH:O	2.12	0.49
2:H:54:ARG:HA	2:H:57:GLU:HG3	1.95	0.48
1:E:174:PHE:C	1:E:174:PHE:CD1	2.87	0.48
1:A:83:LYS:HE2	1:A:83:LYS:HB2	1.70	0.48
1:E:320:MET:HG3	1:E:321:ARG:O	2.14	0.48
2:F:19:ASP:OD2	14:F:301:HOH:O	2.20	0.47
1:C:126:ASN:OD1	1:C:128:THR:HG23	2.15	0.47
2:B:58:LYS:NZ	2:B:60:ASN:HA	2.30	0.47
1:G:167:THR:OG1	1:G:242:ILE:HD11	2.15	0.47
1:C:132:GLN:NE2	1:C:152:ASN:HD21	2.14	0.46



		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
11:B:202:PG4:H41	11:B:202:PG4:H22	1.76	0.46	
7:Z:3:BMA:H62	7:Z:4:MAN:H2	1.70	0.46	
2:B:58:LYS:HE3	2:B:59:THR:C	2.35	0.46	
1:A:234:TRP:NE1	14:A:502:HOH:O	2.23	0.46	
1:E:32:ASP:OD1	1:E:33:GLN:HG2	2.17	0.45	
5:Q:3:BMA:H62	5:Q:5:MAN:H2	1.75	0.45	
11:F:202:PG4:H21	11:F:202:PG4:H41	1.57	0.45	
1:A:143:SER:HB2	2:D:161:VAL:CG1	2.47	0.45	
2:B:39:LYS:HE2	14:B:458:HOH:O	2.17	0.45	
1:C:27:LYS:HG2	1:C:32:ASP:O	2.17	0.45	
12:B:203:PG0:H31	12:B:203:PG0:H11	1.42	0.45	
2:H:171:PHE:O	2:H:172:GLN:HB2	2.18	0.44	
1:E:22:ASN:OD1	1:E:22:ASN:N	2.50	0.44	
1:C:237:VAL:HG21	1:C:243:LEU:HB2	1.99	0.43	
1:G:322:ASN:O	1:G:324:PRO:HD3	2.19	0.43	
1:C:324:PRO:O	1:C:325:GLU:HB2	2.19	0.42	
2:D:171:PHE:O	2:D:172:GLN:HB2	2.19	0.42	
2:B:171:PHE:O	2:B:172:GLN:HB2	2.19	0.42	
1:C:207:LYS:HD2	1:C:241:ASP:HA	2.02	0.42	
2:H:54:ARG:HG2	2:H:57:GLU:OE2	2.19	0.42	
1:E:207:LYS:HG3	1:E:241:ASP:OD1	2.19	0.42	
2:F:171:PHE:O	2:F:172:GLN:HB2	2.19	0.42	
1:A:189:LYS:HB2	1:A:189:LYS:HE3	1.77	0.42	
1:G:177:LEU:HB2	1:G:260:ILE:CD1	2.46	0.42	
3:W:1:NAG:H3	6:Y:2:FUC:O4	2.20	0.42	
1:C:166:VAL:HG12	1:C:245:ILE:HB	2.02	0.41	
2:H:4:GLY:O	2:H:8:GLY:HA3	2.21	0.41	
2:D:60:ASN:HB2	2:D:62:LYS:HE2	2.02	0.41	
1:E:29:ILE:HD13	1:E:29:ILE:HA	1.93	0.41	
1:C:197:ARG:NH1	1:C:248:ASN:O	2.54	0.41	
1:G:186:SER:HA	1:G:218:GLY:O	2.20	0.41	
1:C:321:ARG:HD2	14:C:553:HOH:O	2.20	0.41	
6:Y:2:FUC:H5	6:Y:3:NAG:O5	2.21	0.41	
1:G:259:LYS:HE3	1:G:259:LYS:HB3	1.84	0.41	
2:H:54:ARG:HB2	2:H:54:ARG:NH1	2.36	0.41	
1:A:84:TRP:CE2	1:A:116:GLY:HA2	2.56	0.40	
1:A:316:LEU:HD23	2:B:52:LEU:HD13	2.03	0.40	
11:B:202:PG4:H52	11:B:202:PG4:H72	1.69	0.40	

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	315/323~(98%)	304 (96%)	11 (4%)	0	100 100
1	С	315/323~(98%)	304 (96%)	11 (4%)	0	100 100
1	Е	315/323~(98%)	305~(97%)	9~(3%)	1 (0%)	41 30
1	G	315/323~(98%)	304~(96%)	11 (4%)	0	100 100
2	В	170/174~(98%)	160 (94%)	10 (6%)	0	100 100
2	D	170/174~(98%)	161~(95%)	9~(5%)	0	100 100
2	F	170/174~(98%)	161 (95%)	9~(5%)	0	100 100
2	Н	170/174~(98%)	160 (94%)	10 (6%)	0	100 100
All	All	1940/1988~(98%)	1859 (96%)	80 (4%)	1 (0%)	51 43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Ε	196	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	278/283~(98%)	272~(98%)	6 (2%)	52	44
1	С	278/283~(98%)	272 (98%)	6 (2%)	52	44
1	Ε	278/283~(98%)	272 (98%)	6 (2%)	52	44
1	G	278/283~(98%)	274 (99%)	4 (1%)	67	62



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
2	В	145/148~(98%)	145 (100%)	0	100	100
2	D	145/148~(98%)	143~(99%)	2(1%)	67	62
2	F	146/148~(99%)	145~(99%)	1 (1%)	84	82
2	Н	146/148~(99%)	145~(99%)	1 (1%)	84	82
All	All	1694/1724~(98%)	1668~(98%)	26~(2%)	65	60

Continued from previous page...

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	18	HIS
1	А	137	TYR
1	А	140	LYS
1	А	160	LYS
1	А	321	ARG
1	А	325	GLU
1	С	18	HIS
1	С	24	THR
1	С	83	LYS
1	С	137	TYR
1	С	160	LYS
1	С	321	ARG
2	D	58	LYS
2	D	59	THR
1	Е	18	HIS
1	Е	137	TYR
1	Е	174	PHE
1	Е	189	LYS
1	Е	197	ARG
1	Е	321	ARG
2	F	59	THR
1	G	18	HIS
1	G	137	TYR
1	G	264	LYS
1	G	321	ARG
2	Н	58	LYS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such side chains are listed below:

Mol	Chain	Res	Type
1	А	33	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	А	132	GLN
1	С	132	GLN
1	Е	33	GLN
1	G	33	GLN
2	Н	27	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

66 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Bos	Link	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
IVIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	Ι	1	1,3	14,14,15	0.18	0	17,19,21	0.71	0
3	NAG	Ι	2	3	$14,\!14,\!15$	0.47	0	17,19,21	0.45	0
4	NAG	J	1	1,4	14,14,15	0.92	1 (7%)	17,19,21	0.45	0
4	NAG	J	2	4	14,14,15	0.54	0	17,19,21	0.60	0
4	BMA	J	3	4	11,11,12	0.88	0	$15,\!15,\!17$	0.64	0
5	NAG	K	1	1,5	14,14,15	0.32	0	17,19,21	0.58	0
5	NAG	K	2	5	$14,\!14,\!15$	0.16	0	17,19,21	0.47	0
5	BMA	К	3	5	11,11,12	0.83	0	$15,\!15,\!17$	1.05	1 (6%)
5	MAN	K	4	5	11,11,12	0.82	0	$15,\!15,\!17$	0.99	1 (6%)
5	MAN	K	5	5	11,11,12	0.91	0	15,15,17	1.09	2 (13%)
3	NAG	L	1	1,3	14,14,15	0.43	0	17,19,21	0.54	0
3	NAG	L	2	3	14,14,15	1.38	2 (14%)	17,19,21	1.08	1 (5%)



Mal	Turne	Chain	Dec	T in le	Bond lengths		Bond angles			
WIOI	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	NAG	М	1	1,4	$14,\!14,\!15$	0.24	0	$17,\!19,\!21$	0.78	1 (5%)
4	NAG	М	2	4	14,14,15	0.49	0	17,19,21	0.58	0
4	BMA	М	3	4	11,11,12	0.98	0	$15,\!15,\!17$	0.82	0
4	NAG	N	1	1,4	14,14,15	0.58	0	17,19,21	0.45	0
4	NAG	N	2	4	14,14,15	0.40	0	17,19,21	0.50	0
4	BMA	N	3	4	11,11,12	0.92	0	15, 15, 17	0.84	0
3	NAG	0	1	1,3	14,14,15	0.32	0	17,19,21	0.56	0
3	NAG	0	2	3	14,14,15	0.20	0	17,19,21	0.80	0
6	NAG	Р	1	1,6	14,14,15	0.40	0	17,19,21	0.73	1 (5%)
6	FUC	Р	2	6	10,10,11	0.83	0	14,14,16	0.89	1 (7%)
6	NAG	P	3	6	14,14,15	0.34	0	17,19,21	0.50	0
6	FUC	P	4	6	10,10,11	0.89	0	14,14,16	0.74	0
5	NAG	Q	1	1,5	14,14,15	0.29	0	17,19,21	0.46	0
5	NAG	Q	2	5	14,14,15	0.20	0	17,19,21	0.45	0
5	BMA	Q	3	5	11,11,12	0.53	0	15, 15, 17	0.93	1 (6%)
5	MAN	Q	4	5	11,11,12	0.85	0	15, 15, 17	0.98	2 (13%)
5	MAN	Q	5	5	11,11,12	0.99	0	$15,\!15,\!17$	0.87	1 (6%)
5	NAG	R	1	1,5	14,14,15	0.29	0	17,19,21	0.60	0
5	NAG	R	2	5	14,14,15	0.27	0	17,19,21	0.46	0
5	BMA	R	3	5	11,11,12	0.82	0	$15,\!15,\!17$	0.78	0
5	MAN	R	4	5	11,11,12	1.44	2 (18%)	15, 15, 17	1.15	1 (6%)
5	MAN	R	5	5	11,11,12	0.82	0	$15,\!15,\!17$	1.05	2 (13%)
3	NAG	S	1	1,3	14,14,15	0.22	0	17,19,21	0.68	0
3	NAG	S	2	3	14,14,15	0.99	1 (7%)	$17,\!19,\!21$	1.30	1 (5%)
3	NAG	Т	1	1,3	$14,\!14,\!15$	0.73	1 (7%)	$17,\!19,\!21$	0.45	0
3	NAG	Т	2	3	14,14,15	0.44	0	17,19,21	0.38	0
3	NAG	U	1	1,3	14,14,15	0.40	0	17,19,21	0.44	0
3	NAG	U	2	3	14,14,15	0.52	0	17,19,21	0.51	0
3	NAG	V	1	1,3	$14,\!14,\!15$	0.29	0	$17,\!19,\!21$	0.75	1 (5%)
3	NAG	V	2	3	14,14,15	0.31	0	17,19,21	0.56	0
3	NAG	W	1	1,3	14,14,15	0.21	0	17,19,21	0.72	1(5%)
3	NAG	W	2	3	14,14,15	0.43	0	17,19,21	0.48	0
3	NAG	X	1	1,3	14,14,15	0.60	0	17,19,21	0.43	0
3	NAG	X	2	3	14,14,15	0.67	1 (7%)	17,19,21	0.68	0
6	NAG	Y	1	1,6	14,14,15	0.36	0	17,19,21	0.65	0
6	FUC	Y	2	6	10,10,11	0.91	0	14,14,16	0.91	1 (7%)
6	NAG	Y	3	6	14,14,15	0.45	0	17,19,21	0.51	0
6	FUC	Y	4	6	10,10,11	1.01	1 (10%)	14, 14, 16	1.05	1 (7%)
7	NAG	Ζ	1	1,7	14,14,15	0.39	0	17,19,21	0.43	0



Mal	Tuno	Chain	Bog	Link	Bo	Bond lengths			Bond angles		
	туре	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
7	NAG	Z	2	7	14,14,15	0.16	0	$17,\!19,\!21$	0.55	0	
7	BMA	Ζ	3	7	11,11,12	0.60	0	$15,\!15,\!17$	0.82	0	
7	MAN	Z	4	7	11,11,12	1.25	1 (9%)	$15,\!15,\!17$	1.17	2 (13%)	
7	MAN	Z	5	7	11,11,12	0.68	0	$15,\!15,\!17$	1.07	2 (13%)	
7	MAN	Z	6	7	11,11,12	0.84	0	$15,\!15,\!17$	1.01	2 (13%)	
8	NAG	a	1	8	15,15,15	1.48	3 (20%)	21,21,21	1.49	3 (14%)	
8	GAL	a	2	8	11,11,12	0.82	0	$15,\!15,\!17$	1.10	0	
8	SIA	a	3	8	20,20,21	2.10	3 (15%)	24,28,31	1.42	3 (12%)	
9	GAL	b	1	9	12,12,12	0.54	0	$17,\!17,\!17$	0.82	0	
9	SIA	b	2	9	20,20,21	2.21	3 (15%)	24,28,31	1.48	4 (16%)	
8	NAG	С	1	8	15,15,15	1.35	1 (6%)	$21,\!21,\!21$	1.25	3 (14%)	
8	GAL	С	2	8	11,11,12	0.84	0	$15,\!15,\!17$	1.16	1 (6%)	
8	SIA	С	3	8	20,20,21	2.02	3 (15%)	24,28,31	1.30	4 (16%)	
9	GAL	d	1	9	12,12,12	0.65	0	$17,\!17,\!17$	0.83	0	
9	SIA	d	2	9	20,20,21	2.07	3 (15%)	24,28,31	1.41	5 (20%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	Ι	1	1,3	-	4/6/23/26	0/1/1/1
3	NAG	Ι	2	3	-	0/6/23/26	0/1/1/1
4	NAG	J	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	2/6/23/26	0/1/1/1
4	BMA	J	3	4	-	0/2/19/22	0/1/1/1
5	NAG	Κ	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	Κ	2	5	-	2/6/23/26	0/1/1/1
5	BMA	Κ	3	5	-	2/2/19/22	0/1/1/1
5	MAN	К	4	5	-	2/2/19/22	0/1/1/1
5	MAN	K	5	5	-	0/2/19/22	0/1/1/1
3	NAG	L	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	2/6/23/26	0/1/1/1
4	NAG	М	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	М	2	4	-	2/6/23/26	0/1/1/1
4	BMA	М	3	4	-	0/2/19/22	0/1/1/1
4	NAG	N	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Ν	2	4	-	0/6/23/26	0/1/1/1



8T	J4
----	----

Conti	nuea fro	m previoi	is page	• • •			
Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BMA	N	3	4	-	0/2/19/22	0/1/1/1
3	NAG	0	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	Ο	2	3	-	3/6/23/26	0/1/1/1
6	NAG	Р	1	1,6	-	0/6/23/26	0/1/1/1
6	FUC	Р	2	6	-	-	0/1/1/1
6	NAG	Р	3	6	-	0/6/23/26	0/1/1/1
6	FUC	P	4	6	-	-	0/1/1/1
5	NAG	Q	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	Q	2	5	-	0/6/23/26	0/1/1/1
5	BMA	Q	3	5	-	2/2/19/22	0/1/1/1
5	MAN	Q	4	5	-	1/2/19/22	0/1/1/1
5	MAN	Q	5	5	-	0/2/19/22	0/1/1/1
5	NAG	R	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	R	2	5	-	0/6/23/26	0/1/1/1
5	BMA	R	3	5	-	2/2/19/22	0/1/1/1
5	MAN	R	4	5	-	0/2/19/22	0/1/1/1
5	MAN	R	5	5	-	$\frac{2/2}{19/22}$	0/1/1/1
3	NAG	S	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	S	2	3	-	2/6/23/26	0/1/1/1
3	NAG	Т	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	Т	2	3	-	2/6/23/26	0/1/1/1
3	NAG	U	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	U	2	3	-	2/6/23/26	0/1/1/1
3	NAG	V	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	V	2	3	-	1/6/23/26	0/1/1/1
3	NAG	W	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	W	2	3	-	0/6/23/26	0/1/1/1
3	NAG	Х	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	Х	2	3	-	2/6/23/26	0/1/1/1
6	NAG	Y	1	1,6	-	0/6/23/26	0/1/1/1
6	FUC	Y	2	6	-	-	0/1/1/1
6	NAG	Y	3	6	-	0/6/23/26	0/1/1/1
6	FUC	Y	4	6	-	-	0/1/1/1
	NAG	Z	1	1,7	-	0/6/23/26	0/1/1/1
(NAG	Z	2	(-	0/6/23/26	0/1/1/1
- 1	BMA	Z	3		-	2/2/19/22	0/1/1/1
7	MAN	Z	4	7	-	0/2/19/22	0/1/1/1
7	MAN	Z	5		-	0/2/19/22	$\frac{0/1/1/1}{1/1}$
(MAN		0	(-	0/2/19/22	0/1/1/1

 α tin d fa .



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	a	1	8	-	2/6/26/26	0/1/1/1
8	GAL	a	2	8	-	0/2/19/22	0/1/1/1
8	SIA	a	3	8	-	0/18/34/38	0/1/1/1
9	GAL	b	1	9	-	0/2/22/22	0/1/1/1
9	SIA	b	2	9	-	0/18/34/38	0/1/1/1
8	NAG	с	1	8	-	1/6/26/26	0/1/1/1
8	GAL	с	2	8	-	1/2/19/22	0/1/1/1
8	SIA	с	3	8	-	0/18/34/38	0/1/1/1
9	GAL	d	1	9	-	0/2/22/22	0/1/1/1
9	SIA	d	2	9	-	1/18/34/38	0/1/1/1

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	b	2	SIA	C2-C1	8.58	1.60	1.52
9	d	2	SIA	C2-C1	7.97	1.59	1.52
8	a	3	SIA	C2-C1	7.82	1.59	1.52
8	с	3	SIA	C2-C1	7.63	1.59	1.52
8	с	1	NAG	C1-C2	4.83	1.58	1.52
8	а	1	NAG	C1-C2	4.64	1.58	1.52
3	L	2	NAG	O5-C1	-3.66	1.37	1.43
3	S	2	NAG	O5-C1	3.49	1.49	1.43
5	R	4	MAN	O5-C1	-3.16	1.38	1.43
4	J	1	NAG	O5-C1	-3.10	1.38	1.43
3	L	2	NAG	C1-C2	3.08	1.56	1.52
7	Ζ	4	MAN	C2-C3	2.94	1.56	1.52
9	d	2	SIA	O6-C2	2.85	1.47	1.43
9	b	2	SIA	O6-C2	2.79	1.47	1.43
8	с	3	SIA	O6-C2	2.74	1.47	1.43
8	a	3	SIA	O6-C2	2.67	1.47	1.43
8	a	3	SIA	C7-C6	2.54	1.56	1.53
8	с	3	SIA	C7-C6	2.49	1.56	1.53
8	a	1	NAG	O5-C1	-2.32	1.37	1.42
5	R	4	MAN	C4-C5	2.22	1.57	1.53
3	Х	2	NAG	C1-C2	2.22	1.55	1.52
9	d	2	SIA	C7-C6	2.21	1.55	1.53
3	Т	1	NAG	O5-C1	-2.20	1.40	1.43
6	Y	4	FUC	O5-C1	-2.14	1.40	1.43
9	b	2	SIA	C7-C6	2.14	1.55	1.53
8	a	1	NAG	C3-C2	2.12	1.57	1.53

All (49) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	S	2	NAG	C1-O5-C5	5.05	119.03	112.19
8	a	1	NAG	C4-C3-C2	4.48	116.90	110.34
8	a	1	NAG	C1-C2-C3	3.75	115.66	110.54
8	a	3	SIA	O1A-C1-C2	-3.57	114.14	122.57
8	с	1	NAG	C1-C2-C3	3.48	115.29	110.54
3	L	2	NAG	C4-C3-C2	3.39	115.98	111.02
9	b	2	SIA	O1A-C1-C2	-3.27	114.84	122.57
8	с	1	NAG	C4-C3-C2	3.23	115.07	110.34
7	Ζ	4	MAN	O3-C3-C2	3.00	115.75	109.99
9	d	2	SIA	O1A-C1-C2	-2.95	115.60	122.57
8	a	3	SIA	C6-O6-C2	2.92	117.59	111.34
8	с	3	SIA	O1A-C1-C2	-2.91	115.70	122.57
8	с	3	SIA	C6-O6-C2	2.88	117.49	111.34
8	с	3	SIA	O1B-C1-O1A	2.78	130.39	124.09
9	b	2	SIA	C6-O6-C2	2.76	117.24	111.34
8	a	3	SIA	O1B-C1-O1A	2.75	130.34	124.09
5	К	5	MAN	C1-O5-C5	2.75	115.91	112.19
7	Ζ	5	MAN	C1-O5-C5	2.64	115.77	112.19
8	a	1	NAG	O5-C5-C4	-2.62	104.93	109.69
5	R	5	MAN	C1-O5-C5	2.61	115.73	112.19
3	V	1	NAG	C1-O5-C5	2.53	115.62	112.19
9	d	2	SIA	C6-O6-C2	2.45	116.58	111.34
9	d	2	SIA	O1B-C1-O1A	2.41	129.57	124.09
4	М	1	NAG	C1-O5-C5	2.41	115.46	112.19
5	R	4	MAN	C3-C4-C5	2.39	114.50	110.24
7	Ζ	6	MAN	C1-O5-C5	2.38	115.42	112.19
9	b	2	SIA	O6-C2-C3	-2.38	107.18	110.46
7	Ζ	6	MAN	O2-C2-C3	-2.37	105.38	110.14
7	Ζ	4	MAN	C1-O5-C5	2.37	115.40	112.19
7	Ζ	5	MAN	O2-C2-C3	-2.25	105.62	110.14
8	с	1	NAG	O5-C5-C4	-2.20	105.69	109.69
5	Q	5	MAN	O2-C2-C3	-2.20	105.73	110.14
6	Р	1	NAG	C1-O5-C5	2.20	115.17	112.19
8	с	2	GAL	C1-O5-C5	2.20	115.17	112.19
9	b	2	SIA	O1B-C1-O1A	2.18	129.03	124.09
5	Κ	4	MAN	C1-O5-C5	2.18	115.14	112.19
6	Y	2	FUC	O2-C2-C1	2.17	113.59	109.15
5	Q	4	MAN	O2-C2-C3	-2.17	105.80	110.14
5	Q	3	BMA	C1-O5-C5	2.16	115.12	112.19
9	d	2	SIA	O6-C2-C3	-2.16	107.49	110.46
8	с	3	SIA	C3-C4-C5	-2.12	108.89	111.46
6	Y	4	FUC	O2-C2-C1	2.12	113.48	109.15
5	R	5	MAN	O2-C2-C3	-2.11	105.92	110.14



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	Κ	5	MAN	O2-C2-C3	-2.06	106.00	110.14
9	d	2	SIA	C4-C3-C2	2.03	113.44	109.81
6	Р	2	FUC	O2-C2-C1	2.02	113.29	109.15
3	W	1	NAG	C1-O5-C5	2.02	114.93	112.19
5	Q	4	MAN	C1-O5-C5	2.01	114.92	112.19
5	Κ	3	BMA	C2-C3-C4	2.01	114.37	110.89

There are no chirality outliers.

Mol	Mol Chain		Type	Atoms	
3	Х	2	NAG	O5-C5-C6-O6	
3	0	2	NAG	C4-C5-C6-O6	
5	Q	3	BMA	C4-C5-C6-O6	
7	Ζ	3	BMA	C4-C5-C6-O6	
3	S	2	NAG	O5-C5-C6-O6	
3	U	1	NAG	O5-C5-C6-O6	
3	L	2	NAG	O5-C5-C6-O6	
3	Т	2	NAG	O5-C5-C6-O6	
5	R	3	BMA	O5-C5-C6-O6	
7	Z	3	BMA	O5-C5-C6-O6	
4	J	2	NAG	O5-C5-C6-O6	
3	0	2	NAG	O5-C5-C6-O6	
5	Q	3	BMA	O5-C5-C6-O6	
3	Х	2	NAG	C4-C5-C6-O6	
3	U	1	NAG	C4-C5-C6-O6	
3	L	2	NAG	C4-C5-C6-O6	
5	R	3	BMA	C4-C5-C6-O6	
3	Т	2	NAG	C4-C5-C6-O6	
4	J	2	NAG	C4-C5-C6-O6	
5	K	4	MAN	C4-C5-C6-O6	
3	Ι	1	NAG	O5-C5-C6-O6	
5	K	2	NAG	C4-C5-C6-O6	
8	a	1	NAG	C4-C5-C6-O6	
5	R	5	MAN	O5-C5-C6-O6	
3	Ι	1	NAG	C4-C5-C6-O6	
5	К	3	BMA	C4-C5-C6-O6	
5	K	4	MAN	O5-C5-C6-O6	
5	K	2	NAG	O5-C5-C6-O6	
3	S	2	NAG	C4-C5-C6-O6	
8	с	2	GAL	O5-C5-C6-O6	
3	U	2	NAG	C4-C5-C6-O6	

All (47) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	Q	4	MAN	O5-C5-C6-O6
3	S	1	NAG	C1-C2-N2-C7
5	Κ	3	BMA	O5-C5-C6-O6
8	с	1	NAG	C4-C5-C6-O6
3	Ι	1	NAG	C1-C2-N2-C7
8	a	1	NAG	O5-C5-C6-O6
3	S	1	NAG	C4-C5-C6-O6
3	0	2	NAG	C3-C2-N2-C7
3	V	2	NAG	C3-C2-N2-C7
5	R	5	MAN	C4-C5-C6-O6
3	U	2	NAG	O5-C5-C6-O6
9	d	2	SIA	O1B-C1-C2-O6
4	М	2	NAG	O5-C5-C6-O6
4	М	2	NAG	C4-C5-C6-O6
3	Ι	1	NAG	C3-C2-N2-C7
3	S	1	NAG	O5-C5-C6-O6

Continued from previous page...

There are no ring outliers.

8 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	S	1	NAG	1	0
7	Ζ	4	MAN	1	0
5	Q	5	MAN	1	0
6	Y	3	NAG	1	0
3	W	1	NAG	1	0
5	Q	3	BMA	1	0
7	Ζ	3	BMA	1	0
6	Y	2	FUC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


















































































5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Bos	Link	Bo	ond leng	ths	Bond angles		
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
10	NAG	А	401	1	14,14,15	0.52	0	17,19,21	0.71	0
10	NAG	Н	201	2	14,14,15	0.75	1 (7%)	17,19,21	0.60	0
12	PG0	В	203	-	7,7,7	0.47	0	6,6,6	0.23	0
10	NAG	D	201	2	14,14,15	0.41	0	17,19,21	1.02	2 (11%)
12	PG0	Н	202	-	7,7,7	0.47	0	6,6,6	0.34	0
10	NAG	F	201	2	14,14,15	0.61	1 (7%)	17,19,21	0.51	0
13	PEG	F	203	-	6,6,6	0.18	0	$5,\!5,\!5$	0.16	0
13	PEG	F	204	-	6,6,6	0.19	0	$5,\!5,\!5$	0.07	0
13	PEG	Е	402	-	6,6,6	0.21	0	$5,\!5,\!5$	0.17	0
10	NAG	В	201	2	14,14,15	0.61	1 (7%)	17,19,21	0.57	0
11	PG4	В	202	-	12,12,12	0.25	0	11,11,11	0.65	0
11	PG4	F	202	-	12,12,12	0.21	0	11,11,11	0.68	0
10	NAG	Е	401	1	14,14,15	0.61	0	17,19,21	0.50	0
10	NAG	G	401	1	14,14,15	0.69	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	А	401	1	-	1/6/23/26	0/1/1/1
10	NAG	Н	201	2	-	0/6/23/26	0/1/1/1
12	PG0	В	203	-	-	<mark>5/5/5/5</mark>	-
10	NAG	D	201	2	-	1/6/23/26	0/1/1/1
12	PG0	Н	202	-	-	2/5/5/5	-
10	NAG	F	201	2	-	2/6/23/26	0/1/1/1
13	PEG	F	203	-	-	2/4/4/4	-
13	PEG	F	204	-	-	1/4/4/4	-
13	PEG	Е	402	-	-	2/4/4/4	-

Continued on next page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	В	201	2	-	0/6/23/26	0/1/1/1
11	PG4	В	202	-	-	7/10/10/10	-
11	PG4	F	202	-	-	7/10/10/10	-
10	NAG	Е	401	1	-	0/6/23/26	0/1/1/1
10	NAG	G	401	1	-	0/6/23/26	0/1/1/1

Continued from previous page...

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
10	Н	201	NAG	C1-C2	2.38	1.55	1.52
10	F	201	NAG	C1-C2	2.19	1.55	1.52
10	В	201	NAG	C1-C2	2.07	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
10	D	201	NAG	C1-O5-C5	3.32	116.69	112.19
10	D	201	NAG	C2-N2-C7	2.01	125.77	122.90

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
13	F	204	PEG	C1-C2-O2-C3
13	F	203	PEG	C1-C2-O2-C3
12	В	203	PG0	C1-C2-O1-C3
11	F	202	PG4	C4-C3-O2-C2
13	Е	402	PEG	C1-C2-O2-C3
11	В	202	PG4	C4-C3-O2-C2
12	Н	202	PG0	O1-C3-C4-O2
11	F	202	PG4	O1-C1-C2-O2
11	В	202	PG4	C5-C6-O4-C7
12	В	203	PG0	O1-C3-C4-O2
11	В	202	PG4	O1-C1-C2-O2
13	Е	402	PEG	O2-C3-C4-O4
11	F	202	PG4	O3-C5-C6-O4
12	Н	202	PG0	OTT-C1-C2-O1
10	D	201	NAG	C3-C2-N2-C7
12	В	203	PG0	C3-C4-O2-C5
11	В	202	PG4	O4-C7-C8-O5
13	F	203	PEG	C4-C3-O2-C2

All (30) torsion outliers are listed below:

Continued on next page...



Mol	Chain	Res	Type	Atoms
11	В	202	PG4	C1-C2-O2-C3
11	F	202	PG4	C5-C6-O4-C7
12	В	203	PG0	OTT-C1-C2-O1
11	F	202	PG4	C8-C7-O4-C6
11	F	202	PG4	C1-C2-O2-C3
10	F	201	NAG	C4-C5-C6-O6
10	F	201	NAG	O5-C5-C6-O6
11	В	202	PG4	O3-C5-C6-O4
11	В	202	PG4	O2-C3-C4-O3
11	F	202	PG4	O2-C3-C4-O3
10	А	401	NAG	C3-C2-N2-C7
12	В	203	PG0	C4-C3-O1-C2

Continued from previous page...

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	В	203	PG0	1	0
13	F	203	PEG	2	0
13	F	204	PEG	1	0
13	Е	402	PEG	2	0
11	В	202	PG4	2	0
11	F	202	PG4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.





6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	317/323~(98%)	-0.15	0 100 100	18, 36, 58, 76	0
1	С	317/323~(98%)	-0.11	1 (0%) 94 96	17, 34, 61, 82	0
1	Е	317/323~(98%)	-0.13	2 (0%) 89 93	17, 36, 60, 75	0
1	G	317/323~(98%)	-0.17	0 100 100	17, 35, 61, 83	0
2	В	172/174~(98%)	-0.23	1 (0%) 89 93	17, 26, 42, 69	0
2	D	172/174~(98%)	-0.01	3 (1%) 70 77	16, 26, 46, 80	0
2	F	172/174~(98%)	-0.16	3 (1%) 70 77	18, 26, 43, 68	0
2	Н	172/174~(98%)	-0.22	3 (1%) 70 77	16, 26, 45, 78	0
All	All	1956/1988~(98%)	-0.15	13 (0%) 87 92	16, 31, 58, 83	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Н	58	LYS	5.6
2	Н	57	GLU	5.3
2	D	58	LYS	4.5
2	F	59	THR	4.0
2	D	57	GLU	3.8
2	Н	59	THR	3.2
2	F	57	GLU	2.8
1	С	172	GLY	2.6
1	Е	155	TYR	2.5
2	В	59	THR	2.5
2	D	59	THR	2.5
1	Ε	242	ILE	2.3
2	F	58	LYS	2.1



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	K	5	11/12	0.40	0.27	98,103,107,108	0
5	MAN	Q	4	11/12	0.51	0.19	84,88,92,94	0
7	MAN	Z	6	11/12	0.51	0.18	82,87,89,90	0
8	NAG	с	1	15/15	0.57	0.31	71,97,103,103	0
5	MAN	R	4	11/12	0.60	0.27	97,102,104,106	0
5	MAN	Q	5	11/12	0.62	0.27	76,82,93,94	0
8	NAG	a	1	15/15	0.62	0.37	73,96,107,111	0
3	NAG	U	2	14/15	0.62	0.21	85,97,103,105	0
4	BMA	М	3	11/12	0.65	0.22	92,102,104,105	0
5	MAN	K	4	11/12	0.65	0.17	83,95,99,99	0
3	NAG	V	2	14/15	0.65	0.39	87,94,99,100	0
4	BMA	J	3	11/12	0.67	0.18	92,100,108,108	0
4	BMA	Ν	3	11/12	0.69	0.18	92,102,110,110	0
5	MAN	R	5	11/12	0.71	0.20	91,100,104,106	0
6	NAG	Y	3	14/15	0.72	0.29	75,83,87,87	0
3	NAG	0	2	14/15	0.72	0.28	87,96,102,104	0
7	MAN	Ζ	5	11/12	0.75	0.20	82,93,98,99	0
7	MAN	Ζ	4	11/12	0.75	0.14	82,85,94,97	0
3	NAG	Ι	1	14/15	0.78	0.18	46,62,69,72	0
5	BMA	Q	3	11/12	0.78	0.14	$63,\!69,\!77,\!87$	0
3	NAG	S	1	14/15	0.79	0.18	46,62,67,71	0
6	NAG	Y	1	14/15	0.79	0.17	45,62,77,84	0
6	FUC	Y	4	10/11	0.80	0.38	88,93,99,100	0
3	NAG	Ι	2	14/15	0.80	0.31	$66,\!80,\!92,\!92$	0
5	BMA	R	3	11/12	0.81	0.14	$77,\!82,\!91,\!101$	0
3	NAG	S	2	14/15	0.81	0.19	54,76,86,90	0
3	NAG	Х	2	14/15	0.81	0.19	$59,\!77,\!83,\!87$	0
6	NAG	Р	3	14/15	0.81	0.18	$74,\!81,\!86,\!87$	0
4	NAG	Ν	2	14/15	0.83	0.17	$58,\!80,\!92,\!101$	0
9	GAL	b	1	12/12	0.83	0.24	49,72,79,87	0
6	FUC	Р	4	10/11	0.84	0.19	$7\overline{7,85,90,90}$	0
3	NAG	W	2	14/15	0.84	0.24	62,72,86,90	0

Continued on next page...



8TJ4

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
8	GAL	с	2	11/12	0.85	0.23	48,72,89,95	0
6	FUC	Y	2	10/11	0.86	0.25	58,68,71,71	0
6	FUC	Р	2	10/11	0.86	0.17	59,64,67,68	0
6	NAG	Р	1	14/15	0.87	0.10	43,61,74,80	0
4	NAG	М	2	14/15	0.87	0.29	61,78,94,102	0
3	NAG	Т	2	14/15	0.87	0.16	61,71,79,82	0
4	NAG	J	2	14/15	0.87	0.14	58,72,93,98	0
3	NAG	L	2	14/15	0.87	0.16	89,98,107,107	0
5	BMA	K	3	11/12	0.88	0.08	71,78,88,90	0
5	NAG	R	2	14/15	0.88	0.11	41,60,75,79	0
3	NAG	L	1	14/15	0.89	0.13	55,66,81,89	0
3	NAG	V	1	14/15	0.89	0.17	55,62,72,84	0
3	NAG	U	1	14/15	0.89	0.18	$53,\!65,\!80,\!89$	0
9	GAL	d	1	12/12	0.89	0.16	43,65,79,79	0
5	NAG	Q	1	14/15	0.90	0.16	47,54,65,66	0
5	NAG	Q	2	14/15	0.90	0.12	44,55,60,62	0
3	NAG	W	1	14/15	0.90	0.13	$45,\!50,\!60,\!61$	0
8	GAL	a	2	11/12	0.90	0.15	45,72,85,90	0
4	NAG	М	1	14/15	0.91	0.11	44,52,60,60	0
3	NAG	Х	1	14/15	0.91	0.09	30,42,54,60	0
5	NAG	K	1	14/15	0.91	0.13	$45,\!51,\!57,\!59$	0
3	NAG	Т	1	14/15	0.91	0.11	$30,\!41,\!56,\!58$	0
5	NAG	K	2	14/15	0.92	0.11	41,55,72,76	0
5	NAG	R	1	14/15	0.92	0.12	45,51,58,59	0
7	BMA	Z	3	11/12	0.92	0.10	62,70,79,85	0
4	NAG	N	1	14/15	0.92	0.09	32,39,52,65	0
4	NAG	J	1	14/15	0.93	0.09	27,39,52,56	0
3	NAG	0	1	14/15	0.93	0.10	54,61,71,86	0
8	SIA	с	3	20/21	0.93	0.10	39,46,53,53	0
7	NAG	Z	1	14/15	0.93	0.11	51,55,62,62	0
9	SIA	b	2	20/21	0.93	0.13	36,45,51,59	0
8	SIA	a	3	20/21	0.93	0.10	37,45,52,55	0
9	SIA	d	2	20/21	0.93	0.10	36,44,50,55	0
7	NAG	Ζ	2	14/15	0.94	0.10	$47,\!58,\!63,\!63$	0

d fa α tia

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.












































































6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
10	NAG	А	401	14/15	0.58	0.23	67, 79, 87, 92	0
10	NAG	Н	201	14/15	0.58	0.31	45,74,81,82	0
10	NAG	Е	401	14/15	0.67	0.14	71,75,80,82	0
10	NAG	G	401	14/15	0.70	0.14	70,76,79,80	0
13	PEG	F	203	7/7	0.76	0.16	47,52,58,59	0
11	PG4	В	202	13/13	0.77	0.22	44,50,60,61	0
11	PG4	F	202	13/13	0.78	0.21	41,51,59,61	0
12	PG0	Н	202	8/8	0.79	0.17	42,48,50,59	0
10	NAG	D	201	14/15	0.79	0.18	48,70,74,77	0
13	PEG	E	402	7/7	0.81	0.19	$37,\!42,\!50,\!53$	0
10	NAG	В	201	14/15	0.81	0.16	46,57,70,73	0
13	PEG	F	204	7/7	0.82	0.14	$46,\!51,\!55,\!63$	0
10	NAG	F	201	14/15	0.83	0.15	47,58,70,71	0
12	PG0	В	203	8/8	0.88	0.24	42,54,61,66	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.5 Other polymers (i)

There are no such residues in this entry.

