



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 15, 2023 – 05:32 AM EDT

PDB ID : 1TWC
Title : RNA polymerase II complexed with GTP
Authors : Westover, K.D.; Bushnell, D.A.; Kornberg, R.D.
Deposited on : 2004-06-30
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	
2	B	1224	
3	C	318	
4	E	215	
5	F	155	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	H	146	
7	I	122	
8	J	70	
9	K	120	
10	L	70	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	GTP	B	3008	X	-	-	-

2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 27772 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II largest subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1351	10625	6704	1844	2019	58	0	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase II 140 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	1091	8690	5511	1516	1610	53	0	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase II 45 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	266	2095	1317	348	417	13	0	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerases I, II, and III 27 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	E	215	1760	1116	310	322	12	0	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III 23 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	F	83	670	428	114	125	3	0	0	0

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III 14.5 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	H	133	1068	673	180	211	4	0	0	0

- Molecule 7 is a protein called DNA-DIRECTED RNA POLYMERASE II 14.2KD POLYPEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	I	121	990	610	181	188	11	0	0	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III 8.3 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	J	64	525	334	92	93	6	0	0	0

- Molecule 9 is a protein called DNA-directed RNA polymerase II 13.6 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	K	114	919	590	156	171	2	0	0	0

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III 7.7 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	L	46	364	224	72	64	4	0	0	0

- Molecule 11 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	2	Total	Zn	0	0
			2	2		
11	B	1	Total	Zn	0	0
			1	1		
11	C	1	Total	Zn	0	0
			1	1		
11	I	2	Total	Zn	0	0
			2	2		
11	J	1	Total	Zn	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	L	1	Total	Zn	0	0
			1	1		

- Molecule 12 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	2	Total	Mn	0	0
			2	2		

- Molecule 13 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
13	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	11	Total	O	0	0
			11	11		
14	B	11	Total	O	0	0
			11	11		
14	F	1	Total	O	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	L	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase II largest subunit





• Molecule 2: DNA-directed RNA polymerase II 140 kDa polypeptide





MET	SER	ASP	LEU	ALA	ASN	GLY	F18	F19	E19	D20	E21	S22	A23	P24	I25	T26	A27	E28	D29	S30	W31	A32	V33	P34	S35	A36	F37	F38	R39	E40	K41	G42	G43	Q46	D49	S50	F51	D56	Y57	L58	T59	Q60	D61	I62	E65	D66																											
S67	T68	L69	I70	GLU	LEU	GLN	LEU	K134	K135	ALA	GLN	HIS	THR	THR	GLU	ASP	PRO	ASP	VAL	PRO	GLY	ILE	SER	ARG	ARG	LEU	LYS	TYR	E89	V211	L212	GLU	P224	S91	F92	G93	K94	I95	Y96	V97	T98	K99	P100	M101	V225	A102	N103	E104	S105	D106	G107	V108	T109	H110	A111	L112	Y113	P114	Q115	E116	A117	R118	L119	C179	Y180	R120	L244	E245	N246	L122	T123	Y124	S126
G127	L128	F129	L130	V131	V132	K133	K134	K135	D198	T136	Y137	E138	ALA	ILE	ASP	ASP	VAL	PRO	GLY	ARG	GLU	GLU	LEU	LYS	TYR	LYS	T272	V211	L212	GLU	P224	S91	F92	I167	G107	V108	T109	H110	A111	L112	M173	L174	R175	S176	K177	R178	C179	Y180	R120	L244	E245	N246	L122	T123	A184	T185	R186	R249	F250														
D188	L189	Y190	K191	L192	K193	E194	F197	D198	M199	G200	G201	Y202	F203	I204	A266	R267	N206	G207	S208	E209	K210	V211	L212	I213	A214	Q215	E216	R217	I280	S218	V31	A219	V223	Q224	V225	F226	K227	K228	A229	A230	P231	S232	P233	I234	S235	H236	V237	A238	E239	I240	R241	S242	A243	Y180	E245	N246	L122	T123	G247	S248	R249	F250											
L251	S252	T253	L254	Q255	V256	K257	L258	Y259	G260	R261	G262	G263	S264	S265	A266	R267	N206	G207	S208	E209	K210	V211	L212	I213	A214	Q215	E216	R217	I280	S218	V31	A219	V223	Q224	V225	F226	K227	K228	A229	A230	P231	S232	P233	I234	S235	H236	V237	A238	E239	I240	R241	S242	A243	Y180	E245	N246	L122	T123	G247	S248	R249	F250											
E312	M313	L314	K315	P316	C317	V318	E319	D320	G321	F322	V323	I324	Q325	R326	R327	K393	T329	A330	L331	D332	F333	R336	B337	L341	G342	I343	K344	K345	E346	K347	L284	I285	F286	R287	A288	L289	G290	I291	P292	P293	D294	G295	E296	L297	L298	E299	C302	Y303	D304	V305	N306	D307	W308	G309	M310	L311																	
F377	L378	G379	Y380	N381	N382	N383	R384	L385	L386	L387	C388	A389	L390	D391	R392	K393	D394	O395	D396	A400	F401	K404	R405	L406	D407	L408	L412	L413	A414	Q415	L416	F417	K418	T419	K422	K423	L424	T425	L428	F429	R430	Y431	M432	Q433	R434	T435	F436	E437	GLU	ALA	HIS	ASP																					
PHE	ASN	MET	LYS	L446	A447	L448	R451	T452	L453	T454	S455	G456	L457	K458	Y459	A460	A461	A462	G463	Q464	Q465	A466	G467	GLU	GLN	LYS	ARG	A477	Q478	Q479	S480	Q481	W482	L483	N484	R485	Y486	T487	Y488	S490	R496	L497	T498	N499	T500	P501	L502	GLY	ARG	ASP	GLY																						
LYS	LEU	A509	K510	P511	R512	O513	L514	H515	N516	T517	H518	A519	V522	C523	P524	A525	E526	E529	G530	Q531	A532	C533	G534	L535	V536	K537	N538	L539	M542	S543	S546	T549	D550	P551	M552	P553	L554	L555	T556	F557	L558	S559	E560	M561	G562	M563	E564	P565	F566	L566	E567	D568	Y569	V570																			
E571	H572	Q573	S574	P575	D576	E577	L578	R579	V580	F581	V582	V585	W586	H587	G588	V589	H590	R591	N592	P593	A594	R595	L596	M597	E598	R601	T602	N603	R604	R605	K606	G607	D608	L609	P611	E612	V613	S614	M615	L616	R617	D618	L619	R620	G621	K622	E623	L624	K625	L626	F627	P628	D629	R632	G633	V633																	
Y634	R635	F638	I639	V640	E641	D642	G643	E644	S645	L646	G647	H648	K649	E650	L651	K652	ASN	G653	G654	G655	G656	A657	I658	A659	K660	L661	M662	A663	T664	G665	Q666	I667	D668	ILE	GLY	GLY	PHE	ASP	VAL	GLU	VAL	GLU	E678	Y679	T680	W681	S682	S683	L684	L685	V690	G691	K692	I693	D694	A695	E696	E697															
E698	E699	L702	I703	A704	M705	Q706	D707	E708	D709	L710	F711	P712	ALA	GLU	ALA	ASN	GLU	GLU	ASN	ASP	LEU	D722	V723	D724	P725	A726	K727	R728	I729	R730	V731	H732	S733	H734	A735	T736	I737	F738	I743	H744	P745	S746	M747	V751	F752	S754	I755	L756	P757	F758	G759	P760	P761	H762	Q763	S764																	
P765	R766	Y769	M773	G774	K775	Q776	V777	W780	F781	Y785	Y786	G787	R788	M789	D790	T791	N792	F793	N794	F795	L796	Y797	Y798	Q800	L803	G804	T805	R806	A807	O808	M809	E810	Y811	L812	K813	F814	R815	E816	L817	P818	A819	G820	Q821	N822	V825	A826	I827	A828	C829	Y830	S831	G832	S833	E834	E835																		
M834	Q835	E836	R837	M839	I840	M841	N842	K843	S844	S845	I846	D847	R848	G849	L850	F851	R852	S853	L854	F855	F856	R857	S858	Y859	Q862	E863	K864	K865	Y866	G867	M868	S869	I870	A871	T872	F873	F874	K876	P877	Q878	R879	T880	N881	T882	L883	R884	M885	K886	H887	G888	T889	R890	D891	K892	L893	R894	D895																




● Molecule 3: DNA-directed RNA polymerase II 45 kDa polypeptide




● Molecule 4: DNA-directed RNA polymerases I, II, and III 27 kDa polypeptide







• Molecule 5: DNA-directed RNA polymerases I, II, and III 23 kDa polypeptide




• Molecule 6: DNA-directed RNA polymerases I, II, and III 14.5 kDa polypeptide




• Molecule 7: DNA-DIRECTED RNA POLYMERASE II 14.2KD POLYPEPTIDE




• Molecule 8: DNA-directed RNA polymerases I, II, and III 8.3 kDa polypeptide






- Molecule 9: DNA-directed RNA polymerase II 13.6 kDa polypeptide

Chain K: 23% 39% 22% 10% 5%



- Molecule 10: DNA-directed RNA polymerases I, II, and III 7.7 kDa polypeptide

Chain L: 6% 19% 24% 20% 34%



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	123.23Å 223.61Å 374.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 39.76 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.00) 91.6 (39.76-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.90Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.234 , 0.255 0.198 , 0.236	Depositor DCC
R_{free} test set	3424 reflections (2.99%)	wwPDB-VP
Wilson B-factor (Å ²)	56.3	Xtrriage
Anisotropy	0.418	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	27772	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN, ZN, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.17	1078/10811 (10.0%)	2.44	610/14626 (4.2%)
2	B	3.28	922/8860 (10.4%)	2.47	497/11945 (4.2%)
3	C	3.33	245/2133 (11.5%)	2.58	122/2891 (4.2%)
4	E	3.29	172/1796 (9.6%)	2.31	92/2416 (3.8%)
5	F	2.94	59/682 (8.7%)	2.24	31/922 (3.4%)
6	H	3.24	126/1086 (11.6%)	2.37	58/1470 (3.9%)
7	I	3.57	138/1009 (13.7%)	2.46	75/1357 (5.5%)
8	J	2.92	47/533 (8.8%)	2.82	46/715 (6.4%)
9	K	3.18	100/937 (10.7%)	2.62	58/1265 (4.6%)
10	L	3.58	46/366 (12.6%)	2.81	36/485 (7.4%)
All	All	3.24	2933/28213 (10.4%)	2.46	1625/38092 (4.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	28
2	B	0	32
3	C	0	13
4	E	1	7
6	H	0	13
7	I	0	3
9	K	0	2
10	L	0	4
All	All	1	102

The worst 5 of 2933 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	771	GLU	CD-OE2	27.21	1.55	1.25
2	B	552	MET	CG-SD	25.82	2.48	1.81
1	A	728	LYS	CD-CE	23.14	2.09	1.51
10	L	68	GLU	CG-CD	22.53	1.85	1.51
1	A	1005	GLU	CD-OE1	21.89	1.49	1.25

The worst 5 of 1625 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1366	ARG	NE-CZ-NH2	-34.23	103.18	120.30
1	A	774	ARG	NE-CZ-NH1	32.19	136.39	120.30
3	C	34	ARG	NE-CZ-NH2	-31.55	104.53	120.30
2	B	995	ARG	NE-CZ-NH2	-30.46	105.07	120.30
1	A	1366	ARG	NE-CZ-NH1	28.94	134.77	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	E	204	THR	CB

5 of 102 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	31	SER	Peptide
1	A	44	THR	Peptide
1	A	60	SER	Peptide
1	A	70	CYS	Peptide
1	A	74	MET	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10625	0	10693	807	0
2	B	8690	0	8713	507	0
3	C	2095	0	2051	140	0
4	E	1760	0	1788	125	0
5	F	670	0	690	48	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	1068	0	1040	155	0
7	I	990	0	949	70	0
8	J	525	0	535	39	0
9	K	919	0	929	64	0
10	L	364	0	386	51	0
11	A	2	0	0	0	0
11	B	1	0	0	0	0
11	C	1	0	0	0	0
11	I	2	0	0	0	0
11	J	1	0	0	0	0
11	L	1	0	0	0	0
12	A	2	0	0	0	0
13	B	32	0	11	8	0
14	A	11	0	0	3	0
14	B	11	0	0	4	0
14	F	1	0	0	0	0
14	L	1	0	0	1	0
All	All	27772	0	27785	1880	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 1880 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ILE:CA	1:A:61:ILE:CB	1.75	1.64
1:A:941:LYS:CD	1:A:941:LYS:CG	1.74	1.64
10:L:61:THR:CB	10:L:61:THR:CG2	1.75	1.64
5:F:72:LYS:CD	5:F:72:LYS:CE	1.76	1.64
1:A:1405:THR:CB	1:A:1405:THR:CA	1.77	1.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1334/1733 (77%)	1196 (90%)	84 (6%)	54 (4%)	3	17
2	B	1071/1224 (88%)	931 (87%)	104 (10%)	36 (3%)	3	20
3	C	264/318 (83%)	225 (85%)	30 (11%)	9 (3%)	3	20
4	E	213/215 (99%)	181 (85%)	20 (9%)	12 (6%)	2	10
5	F	81/155 (52%)	72 (89%)	8 (10%)	1 (1%)	13	48
6	H	129/146 (88%)	89 (69%)	17 (13%)	23 (18%)	0	0
7	I	119/122 (98%)	108 (91%)	10 (8%)	1 (1%)	19	57
8	J	62/70 (89%)	60 (97%)	2 (3%)	0	100	100
9	K	112/120 (93%)	99 (88%)	10 (9%)	3 (3%)	5	26
10	L	44/70 (63%)	25 (57%)	11 (25%)	8 (18%)	0	0
All	All	3429/4173 (82%)	2986 (87%)	296 (9%)	147 (4%)	2	15

5 of 147 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35	ILE
1	A	45	GLN
1	A	47	ARG
1	A	59	GLY
1	A	62	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1183/1520 (78%)	993 (84%)	190 (16%)	2	12
2	B	947/1061 (89%)	801 (85%)	146 (15%)	2	13
3	C	234/274 (85%)	194 (83%)	40 (17%)	2	10
4	E	197/197 (100%)	153 (78%)	44 (22%)	1	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	F	73/137 (53%)	62 (85%)	11 (15%)	3	14
6	H	117/128 (91%)	74 (63%)	43 (37%)	0	0
7	I	115/116 (99%)	93 (81%)	22 (19%)	1	8
8	J	59/65 (91%)	49 (83%)	10 (17%)	2	11
9	K	99/102 (97%)	83 (84%)	16 (16%)	2	12
10	L	40/57 (70%)	20 (50%)	20 (50%)	0	0
All	All	3064/3657 (84%)	2522 (82%)	542 (18%)	2	9

5 of 542 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	H	109	LYS
7	I	4	PHE
6	H	108	SER
9	K	113	THR
2	B	65	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 78 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1179	GLN
7	I	12	ASN
3	C	73	GLN
4	E	61	GLN
9	K	2	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 10 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
13	GTP	B	3008	2,12	26,34,34	1.38	2 (7%)	32,54,54	1.53	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	GTP	B	3008	2,12	1/1/7/7	8/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	B	3008	GTP	C8-N7	-5.08	1.26	1.35
13	B	3008	GTP	C2-N3	3.35	1.41	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	B	3008	GTP	O4'-C1'-C2'	4.26	113.16	106.93
13	B	3008	GTP	PA-O3A-PB	-3.55	120.63	132.83
13	B	3008	GTP	PB-O3B-PG	-3.43	121.06	132.83
13	B	3008	GTP	C2'-C3'-C4'	2.37	107.25	102.64

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
13	B	3008	GTP	C1'

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
13	B	3008	GTP	C5'-O5'-PA-O3A
13	B	3008	GTP	O4'-C4'-C5'-O5'
13	B	3008	GTP	C3'-C4'-C5'-O5'
13	B	3008	GTP	PB-O3B-PG-O2G
13	B	3008	GTP	C5'-O5'-PA-O2A

There are no ring outliers.

1 monomer is involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
13	B	3008	GTP	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	366:VAL	C	367:PRO	N	1.20

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1351/1733 (77%)	-0.44	23 (1%) 70 41	20, 46, 110, 159	0
2	B	1091/1224 (89%)	-0.34	37 (3%) 45 19	21, 44, 116, 146	0
3	C	266/318 (83%)	-0.51	2 (0%) 86 65	29, 50, 79, 131	0
4	E	215/215 (100%)	-0.35	5 (2%) 60 31	23, 60, 108, 138	0
5	F	83/155 (53%)	-0.50	0 100 100	24, 45, 71, 80	0
6	H	133/146 (91%)	0.06	4 (3%) 50 22	56, 88, 128, 135	0
7	I	121/122 (99%)	-0.44	1 (0%) 86 65	30, 50, 81, 111	0
8	J	64/70 (91%)	-0.59	0 100 100	31, 42, 71, 81	0
9	K	114/120 (95%)	-0.41	0 100 100	31, 58, 79, 86	0
10	L	46/70 (65%)	0.15	4 (8%) 10 3	49, 95, 122, 125	0
All	All	3484/4173 (83%)	-0.38	76 (2%) 62 33	20, 49, 113, 159	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1109	GLY	7.4
2	B	246	LYS	7.4
2	B	882	THR	7.0
1	A	44	THR	6.8
1	A	188	ASP	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
12	MN	A	3010	1/1	0.85	0.13	48,48,48,48	0
13	GTP	B	3008	32/32	0.90	0.23	96,112,119,119	0
11	ZN	A	3006	1/1	0.93	0.12	67,67,67,67	0
11	ZN	L	3005	1/1	0.93	0.09	96,96,96,96	0
11	ZN	B	3007	1/1	0.96	0.11	64,64,64,64	0
12	MN	A	3009	1/1	0.97	0.13	36,36,36,36	0
11	ZN	A	3008	1/1	0.98	0.12	91,91,91,91	0
11	ZN	C	3002	1/1	0.99	0.09	53,53,53,53	0
11	ZN	I	3003	1/1	0.99	0.12	59,59,59,59	0
11	ZN	I	3004	1/1	0.99	0.05	80,80,80,80	0
11	ZN	J	3001	1/1	0.99	0.16	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.