



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 24, 2023 – 08:19 AM EDT

PDB ID : 5U17
Title : Structure of human MR1-DA-6-FP in complex with human MAIT A-F7 TCR
Authors : Keller, A.N.; Rossjohn, J.
Deposited on : 2016-11-28
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	 2% 89% 7% . .
1	C	271	 5% 86% 11% .
2	B	203	 8% 87% 9% .
2	D	203	 2% 91% 8% .
3	E	245	 % 89% 10% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	245	 4% 87% 10% ..
4	F	99	 13% 87% 10% ..
4	H	99	 91% 9%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 13789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major histocompatibility complex class I-related gene protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2157	1379	373	395	10	0	0	0
1	C	262	2152	1375	370	396	11	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP Q95460
A	261	SER	CYS	conflict	UNP Q95460
C	0	MET	-	initiating methionine	UNP Q95460
C	261	SER	CYS	conflict	UNP Q95460

- Molecule 2 is a protein called MAIT T-cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	195	1514	958	244	303	9	0	2	0
2	D	200	1565	986	251	319	9	0	1	0


- Molecule 3 is a protein called MAIT T-cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	244	1920	1207	334	370	9	0	0	0
3	G	241	1899	1195	326	369	9	0	2	0

- Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	97	Total	C	N	O	S	0	0	0
			792	507	135	148	2			
4	H	99	Total	C	N	O	S	0	0	0
			817	520	138	157	2			

- Molecule 5 is 2,4-diaminopteridine-6-carbaldehyde (three-letter code: 7WP) (formula: C₇H₆N₆O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			13	7	6		
5	C	1	Total	C	N	0	0
			13	7	6		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0
6	H	1	Total C O 6 3 3	0	0
6	H	1	Total C O 6 3 3	0	0

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	D	1	Total C O 4 2 2	0	0
7	F	1	Total C O 4 2 2	0	0

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	E	1	Total Na 1 1	0	0
8	G	1	Total Na 1 1	0	0
8	H	1	Total Na 1 1	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	159	Total O 159 159	0	0
9	B	96	Total O 96 96	0	0
9	C	149	Total O 149 149	0	0
9	D	134	Total O 134 134	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	E	141	Total 141	O 141	0	0
9	F	32	Total 32	O 32	0	0
9	G	102	Total 102	O 102	0	0
9	H	69	Total 69	O 69	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major histocompatibility complex class I-related gene protein

Chain A: 





- Molecule 1: Major histocompatibility complex class I-related gene protein

Chain C: 




- Molecule 2: MAIT T-cell receptor alpha chain

Chain B: 




- Molecule 2: MAIT T-cell receptor alpha chain



Chain D: 





- Molecule 3: MAIT T-cell receptor beta chain




- Molecule 3: MAIT T-cell receptor beta chain



- Molecule 4: Beta-2-microglobulin



- Molecule 4: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	215.29Å 69.74Å 142.36Å 90.00° 104.02° 90.00°	Depositor
Resolution (Å)	45.35 – 2.15 45.35 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.35-2.15) 100.0 (45.35-2.15)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.16Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.180 , 0.224 0.180 , 0.224	Depositor DCC
R_{free} test set	5605 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtrriage
Anisotropy	0.266	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13789	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ACT, 7WP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/2222	0.64	1/3022 (0.0%)
1	C	0.49	0/2216	0.59	0/3014
2	B	0.46	0/1553	0.57	0/2108
2	D	0.52	0/1600	0.60	0/2170
3	E	0.50	0/1971	0.61	1/2682 (0.0%)
3	G	0.43	0/1950	0.58	1/2657 (0.0%)
4	F	0.40	0/815	0.58	0/1108
4	H	0.48	0/840	0.60	0/1141
All	All	0.48	0/13167	0.60	3/17902 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	LEU	CA-CB-CG	-7.56	97.92	115.30
3	G	178	LEU	CA-CB-CG	5.45	127.84	115.30
3	E	194	ARG	NE-CZ-NH2	-5.28	117.66	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2157	0	2028	18	0
1	C	2152	0	2016	21	0
2	B	1514	0	1413	15	0
2	D	1565	0	1474	12	0
3	E	1920	0	1812	17	0
3	G	1899	0	1766	18	0
4	F	792	0	745	6	0
4	H	817	0	765	7	0
5	A	13	0	0	0	0
5	C	13	0	0	0	0
6	A	6	0	8	1	0
6	B	6	0	8	2	0
6	C	24	0	32	2	0
6	E	6	0	8	1	0
6	H	12	0	16	1	0
7	D	4	0	3	1	0
7	F	4	0	3	0	0
8	E	1	0	0	0	0
8	G	1	0	0	0	0
8	H	1	0	0	0	0
9	A	159	0	0	5	0
9	B	96	0	0	4	0
9	C	149	0	0	2	0
9	D	134	0	0	1	0
9	E	141	0	0	3	0
9	F	32	0	0	0	0
9	G	102	0	0	2	0
9	H	69	0	0	0	0
All	All	13789	0	12097	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 101 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:206:ARG:HH21	6:E:301:GOL:H31	1.26	0.98
1:A:16:ILE:HG22	1:A:17:HIS:H	1.41	0.85
1:A:1:ARG:O	9:A:401:HOH:O	2.03	0.77
1:C:249:GLN:OE1	9:C:401:HOH:O	2.02	0.77
2:B:150[B]:ASP:OD1	9:B:401:HOH:O	2.03	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/271 (96%)	251 (96%)	9 (4%)	0	100	100
1	C	257/271 (95%)	252 (98%)	5 (2%)	0	100	100
2	B	193/203 (95%)	188 (97%)	5 (3%)	0	100	100
2	D	199/203 (98%)	194 (98%)	5 (2%)	0	100	100
3	E	242/245 (99%)	235 (97%)	7 (3%)	0	100	100
3	G	241/245 (98%)	236 (98%)	5 (2%)	0	100	100
4	F	95/99 (96%)	94 (99%)	1 (1%)	0	100	100
4	H	97/99 (98%)	97 (100%)	0	0	100	100
All	All	1584/1636 (97%)	1547 (98%)	37 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/241 (94%)	221 (97%)	6 (3%)	46	47
1	C	228/241 (95%)	226 (99%)	2 (1%)	78	83
2	B	165/180 (92%)	164 (99%)	1 (1%)	86	90
2	D	176/180 (98%)	175 (99%)	1 (1%)	86	90

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	E	207/211 (98%)	205 (99%)	2 (1%)	76	81
3	G	204/211 (97%)	201 (98%)	3 (2%)	65	69
4	F	87/94 (93%)	85 (98%)	2 (2%)	50	53
4	H	91/94 (97%)	90 (99%)	1 (1%)	73	78
All	All	1385/1452 (95%)	1367 (99%)	18 (1%)	69	74

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	112	LEU
4	H	70	PHE
3	G	194	ARG
1	C	94	ARG
4	F	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	249	GLN
3	E	212	GLN
3	G	17	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 3 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ACT	D	301	-	3,3,3	0.78	0	3,3,3	0.88	0
5	7WP	C	301	1	14,14,15	0.59	0	18,20,21	0.73	0
6	GOL	H	102	-	5,5,5	0.35	0	5,5,5	0.58	0
6	GOL	B	301	-	5,5,5	0.37	0	5,5,5	0.83	0
7	ACT	F	101	-	3,3,3	0.80	0	3,3,3	1.16	0
5	7WP	A	301	1	14,14,15	0.78	0	18,20,21	0.91	0
6	GOL	C	302	-	5,5,5	0.51	0	5,5,5	0.93	0
6	GOL	C	303	-	5,5,5	0.31	0	5,5,5	0.55	0
6	GOL	C	304	-	5,5,5	0.34	0	5,5,5	0.33	0
6	GOL	C	305	-	5,5,5	0.37	0	5,5,5	0.26	0
6	GOL	H	101	-	5,5,5	0.32	0	5,5,5	1.03	1 (20%)
6	GOL	A	302	-	5,5,5	0.33	0	5,5,5	0.53	0
6	GOL	E	301	-	5,5,5	0.53	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	7WP	C	301	1	-	-	0/2/2/2
6	GOL	H	102	-	-	2/4/4/4	-
6	GOL	B	301	-	-	4/4/4/4	-
5	7WP	A	301	1	-	-	0/2/2/2
6	GOL	C	304	-	-	4/4/4/4	-
6	GOL	C	302	-	-	1/4/4/4	-
6	GOL	C	303	-	-	2/4/4/4	-
6	GOL	C	305	-	-	2/4/4/4	-
6	GOL	H	101	-	-	3/4/4/4	-
6	GOL	A	302	-	-	2/4/4/4	-
6	GOL	E	301	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	101	GOL	O2-C2-C1	2.02	118.01	109.12

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	302	GOL	O1-C1-C2-C3
6	C	303	GOL	O1-C1-C2-C3
6	C	304	GOL	C1-C2-C3-O3
6	C	305	GOL	O1-C1-C2-C3
6	E	301	GOL	O1-C1-C2-C3

There are no ring outliers.

7 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	D	301	ACT	1	0
6	B	301	GOL	2	0
6	C	303	GOL	1	0
6	C	304	GOL	1	0
6	H	101	GOL	1	0
6	A	302	GOL	1	0
6	E	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/271 (97%)	0.04	6 (2%) 60 68	23, 32, 64, 106	0
1	C	262/271 (96%)	0.16	13 (4%) 28 37	21, 35, 66, 91	0
2	B	195/203 (96%)	0.45	17 (8%) 10 14	24, 43, 86, 99	0
2	D	200/203 (98%)	0.04	4 (2%) 65 72	24, 34, 61, 79	0
3	E	244/245 (99%)	-0.10	2 (0%) 86 89	25, 36, 59, 87	0
3	G	241/245 (98%)	0.11	10 (4%) 37 46	24, 40, 71, 88	0
4	F	97/99 (97%)	0.69	13 (13%) 3 4	27, 54, 86, 91	0
4	H	99/99 (100%)	0.03	0 100 100	23, 36, 62, 76	0
All	All	1602/1636 (97%)	0.14	65 (4%) 37 46	21, 36, 72, 106	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	195	VAL	7.2
1	C	222	VAL	6.2
1	A	251	SER	5.9
1	C	223	GLN	5.7
1	A	252	ASN	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	ACT	D	301	4/4	0.80	0.19	56,56,60,62	0
6	GOL	A	302	6/6	0.86	0.21	66,67,68,68	0
6	GOL	B	301	6/6	0.87	0.19	47,51,52,52	0
8	NA	H	103	1/1	0.87	0.15	53,53,53,53	0
6	GOL	H	102	6/6	0.88	0.25	72,72,72,73	0
6	GOL	C	303	6/6	0.89	0.17	46,47,48,49	0
7	ACT	F	101	4/4	0.90	0.33	57,58,58,58	0
6	GOL	C	304	6/6	0.93	0.14	40,43,45,47	0
8	NA	G	301	1/1	0.93	0.11	43,43,43,43	0
6	GOL	E	301	6/6	0.93	0.17	44,47,47,49	0
6	GOL	C	302	6/6	0.94	0.13	32,34,43,46	0
6	GOL	C	305	6/6	0.94	0.18	52,54,56,58	0
6	GOL	H	101	6/6	0.95	0.14	31,35,36,36	0
5	7WP	C	301	13/14	0.95	0.12	20,26,31,31	0
5	7WP	A	301	13/14	0.96	0.15	23,28,30,31	0
8	NA	E	302	1/1	0.98	0.08	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.