



wwPDB X-ray Structure Validation Summary Report

Dec 13, 2023 – 03:44 am GMT

PDB ID : 4U2K
Title : X-ray structure uridine phosphorylase from *Vibrio cholerae* in complex with anticancer compound at 2.13 Å resolution
Authors : Prokofev, I.I.; Lashkov, A.A.; Gabdoulkhakov, A.G.; Betzel, C.; Mikhailov, A.M.
Deposited on : 2014-07-17
Resolution : 2.13 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	 12% 77% 20%
1	B	253	 15% 78% 18%
1	C	253	 13% 75% 23%
1	D	253	 16% 80% 15%
1	E	253	 11% 77% 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	253	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	D	301	-	-	X	-
5	PEG	D	302	-	-	X	-
6	M5F	D	303[B]	-	X	-	-
6	M5F	F	302	-	X	-	-
7	TRS	F	301	-	-	-	X

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 11998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	Total 1840	C 1154	N 321	O 352	S 13	0	2	0
1	B	245	Total 1848	C 1158	N 318	O 357	S 15	0	4	0
1	C	251	Total 1920	C 1204	N 336	O 366	S 14	0	6	0
1	D	242	Total 1866	C 1169	N 330	O 354	S 13	0	8	0
1	E	246	Total 1865	C 1170	N 328	O 353	S 14	0	4	0
1	F	251	Total 1918	C 1207	N 330	O 366	S 15	0	7	0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	E	1	Total C O 4 2 2	0	0
2	E	1	Total C O 4 2 2	0	0

- Molecule 3 is 1-[(2R)-2,3-diaminopropyl]-5-fluoropyrimidine-2,4(1H,3H)-dione (three-letter code: 4WR) (formula: C₇H₁₁FN₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C F N O 14 7 1 4 2	0	0
3	D	1	Total C F N O 14 7 1 4 2	0	1

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is 1-[(2S)-2,3-diaminopropyl]-5-fluoropyrimidine-2,4(1H,3H)-dione (three-letter

code: M5F) (formula: C₇H₁₁FN₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	F	N			O
6	D	1	14	7	1	4	2	0	1
6	F	1	14	7	1	4	2	0	0

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	F	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	110	Total	O	0	8
			118	118		
8	B	86	Total	O	0	4
			90	90		
8	C	111	Total	O	0	6
			117	117		
8	D	85	Total	O	0	8
			93	93		
8	E	115	Total	O	0	6
			121	121		
8	F	85	Total	O	0	6
			91	91		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase





- Molecule 1: Uridine phosphorylase









• Molecule 1: Uridine phosphorylase



• Molecule 1: Uridine phosphorylase



• Molecule 1: Uridine phosphorylase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	91.28Å 91.28Å 137.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.77 – 2.13 19.77 – 2.13	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.77-2.13) 100.0 (19.77-2.13)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.81 (at 2.13Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.208 , 0.262 0.209 , 0.262	Depositor DCC
R_{free} test set	2160 reflections (3.02%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.745	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.000 for h,-h-k,-l 0.000 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11998	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5209e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, PEG, EDO, GOL, M5F, 4WR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1874	0.61	0/2539
1	B	0.44	0/1888	0.59	0/2555
1	C	0.45	0/1969	0.58	0/2665
1	D	0.44	0/1916	0.59	0/2591
1	E	0.43	0/1906	0.61	0/2578
1	F	0.44	0/1971	0.62	0/2670
All	All	0.44	0/11524	0.60	0/15598

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1840	0	1856	30	0
1	B	1848	0	1861	29	0
1	C	1920	0	1943	47	0
1	D	1866	0	1888	33	0
1	E	1865	0	1889	37	0
1	F	1918	0	1941	41	0
2	A	8	0	12	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	4	0	6	0	0
2	C	8	0	12	3	0
2	E	8	0	12	2	0
3	B	14	0	11	2	0
3	D	14	0	11	1	0
4	D	6	0	8	4	0
4	E	6	0	8	0	0
5	D	7	0	10	5	0
6	D	14	0	11	2	0
6	F	14	0	11	2	0
7	F	8	0	12	0	0
8	A	118	0	0	2	0
8	B	90	0	0	5	0
8	C	117	0	0	3	0
8	D	93	0	0	5	0
8	E	121	0	0	3	0
8	F	91	0	0	4	0
All	All	11998	0	11502	211	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 211 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:26:ASP:HB3	1:D:29[A]:ARG:HG2	1.63	0.80
1:C:174[B]:ARG:NH1	8:C:402:HOH:O	2.17	0.78
1:D:132:ASP:HB2	5:D:302:PEG:H41	1.66	0.77
1:C:29:ARG:HH11	1:C:32:LYS:HZ2	1.36	0.73
3:B:302:4WR:O4	8:B:401:HOH:O	2.06	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/253 (96%)	236 (97%)	6 (2%)	1 (0%)	34	29
1	B	245/253 (97%)	243 (99%)	1 (0%)	1 (0%)	34	29
1	C	255/253 (101%)	250 (98%)	4 (2%)	1 (0%)	34	29
1	D	246/253 (97%)	241 (98%)	4 (2%)	1 (0%)	34	29
1	E	246/253 (97%)	237 (96%)	8 (3%)	1 (0%)	34	29
1	F	256/253 (101%)	245 (96%)	10 (4%)	1 (0%)	34	29
All	All	1491/1518 (98%)	1452 (97%)	33 (2%)	6 (0%)	34	29

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	TYR
1	B	162	TYR
1	F	162	TYR
1	C	162	TYR
1	E	162	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/203 (97%)	191 (97%)	6 (3%)	41	39
1	B	200/203 (98%)	194 (97%)	6 (3%)	41	39
1	C	207/203 (102%)	200 (97%)	7 (3%)	37	34
1	D	201/203 (99%)	195 (97%)	6 (3%)	41	39
1	E	200/203 (98%)	193 (96%)	7 (4%)	36	33
1	F	208/203 (102%)	198 (95%)	10 (5%)	25	21
All	All	1213/1218 (100%)	1171 (96%)	42 (4%)	37	33

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	195	GLU
1	F	55	GLU
1	E	222[A]	ARG
1	F	48	GLU
1	F	195	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	TRS	F	301	-	7,7,7	0.30	0	9,9,9	0.27	0
2	EDO	B	301	-	3,3,3	0.60	0	2,2,2	0.28	0
3	4WR	B	302	-	12,14,14	2.27	4 (33%)	15,19,19	3.63	9 (60%)
6	M5F	F	302	-	12,14,14	4.60	9 (75%)	15,19,19	3.65	8 (53%)
2	EDO	E	303	-	3,3,3	0.49	0	2,2,2	0.43	0
3	4WR	D	304[A]	-	12,14,14	2.03	2 (16%)	15,19,19	3.56	8 (53%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	E	301	-	5,5,5	0.36	0	5,5,5	0.47	0
2	EDO	A	301	-	3,3,3	0.48	0	2,2,2	0.25	0
4	GOL	D	301	-	5,5,5	0.40	0	5,5,5	0.49	0
2	EDO	C	302	-	3,3,3	0.46	0	2,2,2	0.31	0
2	EDO	E	302	-	3,3,3	0.42	0	2,2,2	0.43	0
6	M5F	D	303[B]	-	12,14,14	4.78	9 (75%)	15,19,19	3.47	8 (53%)
2	EDO	A	302	-	3,3,3	0.43	0	2,2,2	0.36	0
5	PEG	D	302	-	6,6,6	0.41	0	5,5,5	0.57	0
2	EDO	C	301	-	3,3,3	0.48	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TRS	F	301	-	-	3/9/9/9	-
2	EDO	B	301	-	-	1/1/1/1	-
3	4WR	B	302	-	-	1/6/6/6	0/1/1/1
6	M5F	F	302	-	-	3/6/6/6	0/1/1/1
2	EDO	E	303	-	-	0/1/1/1	-
3	4WR	D	304[A]	-	-	2/6/6/6	0/1/1/1
4	GOL	E	301	-	-	2/4/4/4	-
2	EDO	A	301	-	-	1/1/1/1	-
4	GOL	D	301	-	-	4/4/4/4	-
2	EDO	C	302	-	-	1/1/1/1	-
2	EDO	E	302	-	-	1/1/1/1	-
6	M5F	D	303[B]	-	-	2/6/6/6	0/1/1/1
2	EDO	A	302	-	-	0/1/1/1	-
5	PEG	D	302	-	-	2/4/4/4	-
2	EDO	C	301	-	-	1/1/1/1	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	F	302	M5F	O4-C4	8.55	1.39	1.23
6	D	303[B]	M5F	C4-C5	-8.50	1.33	1.44
6	D	303[B]	M5F	O4-C4	7.94	1.38	1.23
6	F	302	M5F	C6-C5	7.59	1.44	1.33
6	D	303[B]	M5F	C6-C5	7.52	1.44	1.33

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	4WR	C5-C4-N3	7.44	119.88	112.56
6	D	303[B]	M5F	C5-C4-N3	6.61	119.06	112.56
3	D	304[A]	4WR	C5-C4-N3	6.60	119.06	112.56
6	F	302	M5F	N3-C2-N1	6.58	120.88	114.86
3	D	304[A]	4WR	N3-C2-N1	6.57	120.87	114.86

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	4WR	NAJ-CAL-CAM-CAB
3	D	304[A]	4WR	NAJ-CAL-CAM-CAB
4	D	301	GOL	O1-C1-C2-C3
4	D	301	GOL	O2-C2-C3-O3
6	D	303[B]	M5F	N1-CAB-CAM-CAL

There are no ring outliers.

9 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	4WR	2	0
6	F	302	M5F	2	0
2	E	303	EDO	2	0
3	D	304[A]	4WR	1	0
4	D	301	GOL	4	0
6	D	303[B]	M5F	2	0
2	A	302	EDO	1	0
5	D	302	PEG	5	0
2	C	301	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/253 (96%)	0.89	30 (12%) 4 5	9, 20, 37, 55	0
1	B	245/253 (96%)	1.01	39 (15%) 1 2	10, 21, 39, 57	0
1	C	251/253 (99%)	1.06	32 (12%) 3 4	9, 19, 41, 72	0
1	D	242/253 (95%)	1.11	41 (16%) 1 1	11, 21, 39, 53	0
1	E	246/253 (97%)	0.87	27 (10%) 5 7	11, 20, 37, 57	0
1	F	251/253 (99%)	1.16	50 (19%) 1 1	10, 22, 44, 70	0
All	All	1480/1518 (97%)	1.02	219 (14%) 2 3	9, 20, 40, 72	0

The worst 5 of 219 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	228	PRO	12.6
1	B	232	THR	12.1
1	F	227	ILE	11.4
1	F	230	HIS	9.3
1	C	231	ALA	9.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	TRS	F	301	8/8	0.59	0.45	44,49,57,58	0
4	GOL	D	301	6/6	0.70	0.28	29,29,38,40	0
3	4WR	D	304[A]	14/14	0.75	0.24	21,24,29,29	14
3	4WR	B	302	14/14	0.77	0.27	15,22,32,34	14
6	M5F	D	303[B]	14/14	0.78	0.22	21,24,29,29	14
6	M5F	F	302	14/14	0.79	0.23	17,26,36,39	14
2	EDO	B	301	4/4	0.83	0.22	14,20,23,46	0
2	EDO	C	301	4/4	0.83	0.20	28,28,35,37	0
2	EDO	E	302	4/4	0.84	0.24	25,28,34,35	0
4	GOL	E	301	6/6	0.86	0.18	27,34,38,40	0
2	EDO	A	302	4/4	0.86	0.29	20,21,26,38	0
2	EDO	C	302	4/4	0.87	0.12	24,27,31,31	0
5	PEG	D	302	7/7	0.89	0.15	10,17,29,29	0
2	EDO	E	303	4/4	0.89	0.16	19,19,26,29	0
2	EDO	A	301	4/4	0.94	0.15	12,17,28,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.