



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2023 – 08:08 AM EDT

PDB ID : 3U5N  
Title : Crystal structure of the complex of TRIM33 PHD-Bromo and H3(1-20)K9me3K14ac histone peptide  
Authors : Wang, Z.; Patel, D.J.  
Deposited on : 2011-10-11  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	
1	B	207	
2	C	20	
2	D	20	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3443 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase TRIM33.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	185	1489	950	246	278	15	0	0	0
1	B	186	1498	957	245	281	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	881	SER	-	expression tag	UNP Q9UPN9
B	881	SER	-	expression tag	UNP Q9UPN9

- Molecule 2 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	10	82	49	19	14	0	0	0
2	D	20	158	96	35	27	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	121	Total	O	0	0
			121	121		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	B	86	Total O 86 86	0	0
4	D	5	Total O 5 5	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase TRIM33



- Molecule 1: E3 ubiquitin-protein ligase TRIM33




- Molecule 2: Histone H3.1



- Molecule 2: Histone H3.1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.59Å 113.74Å 39.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.79 – 1.95 29.79 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.4 (29.79-1.95) 92.0 (29.79-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.59 (at 1.95Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.202 , 0.256 0.198 , 0.252	Depositor DCC
$R_{free}$ test set	1652 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.5	Xtrriage
Anisotropy	0.910	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3443	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, M3L, ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/1523	0.54	0/2057
1	B	0.34	0/1533	0.52	0/2073
2	C	0.37	0/69	0.59	0/90
2	D	0.34	0/133	0.51	0/174
All	All	0.40	0/3258	0.53	0/4394

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1489	0	1450	37	0
1	B	1498	0	1451	45	0
2	C	82	0	97	2	0
2	D	158	0	181	11	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	121	0	0	5	0
4	B	86	0	0	1	0
4	D	5	0	0	0	0
All	All	3443	0	3179	84	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:895:ASN:HD22	1:B:896:GLY:H	1.13	0.96
1:A:977:GLU:CD	1:A:977:GLU:H	1.77	0.87
1:B:1002:ASP:OD1	4:B:174:HOH:O	1.96	0.84
1:B:995:LYS:O	1:B:998:LYS:NZ	2.10	0.83
1:B:904:LYS:HB2	1:B:927:ILE:HD11	1.61	0.82
1:B:895:ASN:ND2	1:B:896:GLY:H	1.81	0.79
1:A:1016:GLN:OE1	2:D:20:LEU:N	2.14	0.79
1:A:1061:GLU:OE1	1:A:1061:GLU:N	2.15	0.78
1:B:895:ASN:HD21	2:D:9:M3L:H	1.30	0.77
1:A:1020:ILE:HG13	1:A:1021:PRO:HD2	1.66	0.76
1:A:1061:GLU:H	1:A:1061:GLU:CD	1.90	0.72
1:B:895:ASN:HD22	1:B:896:GLY:N	1.86	0.70
1:B:1014:HIS:HD2	1:B:1016:GLN:H	1.40	0.68
1:B:1040:GLU:HG2	1:B:1041:MET:HE3	1.75	0.68
1:A:905:CYS:HB2	1:A:906:PRO:HD2	1.76	0.68
1:B:977:GLU:HG3	1:B:978:LEU:HD13	1.76	0.68
1:A:1041:MET:SD	4:A:83:HOH:O	2.51	0.67
1:A:883:ASP:CG	1:A:884:ASP:H	1.96	0.67
1:A:999:LYS:HD2	1:A:1030:LEU:HD21	1.78	0.66
1:B:1020:ILE:HG23	1:B:1021:PRO:HD2	1.76	0.66
1:B:999:LYS:HG3	1:B:1030:LEU:HD21	1.76	0.65
1:A:965:LYS:NZ	4:A:80:HOH:O	2.32	0.62
1:B:932:ARG:O	1:B:968:ARG:NH2	2.34	0.61
1:A:916:PRO:HG3	1:A:944:ASP:HA	1.82	0.61
1:B:1061:GLU:OE1	1:B:1061:GLU:N	2.25	0.60
1:B:980:ILE:HD13	1:B:980:ILE:H	1.68	0.59
1:B:980:ILE:HD13	1:B:980:ILE:N	2.17	0.59
1:A:930:PHE:HA	1:A:968:ARG:HD2	1.85	0.58
1:A:895:ASN:HD22	1:A:896:GLY:H	1.51	0.58
1:A:1015:SER:HB3	2:D:17:ARG:NH2	2.18	0.58
1:B:980:ILE:H	1:B:980:ILE:CD1	2.16	0.58
1:A:977:GLU:CD	1:A:977:GLU:N	2.53	0.57
1:B:1029:ARG:O	1:B:1033:LYS:HG3	2.05	0.57
1:B:977:GLU:O	1:B:980:ILE:HD11	2.05	0.57
1:A:883:ASP:CG	1:A:884:ASP:N	2.58	0.55
1:B:895:ASN:ND2	2:D:9:M3L:H	2.03	0.55
1:A:1067:LYS:NZ	4:A:149:HOH:O	2.40	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1067:LYS:O	1:A:1071:LEU:HD23	2.07	0.55
1:A:888:ASP:O	2:C:6:THR:HG21	2.07	0.54
1:B:1012:LYS:HA	1:B:1017:HIS:CD2	2.42	0.54
1:B:904:LYS:HB2	1:B:927:ILE:CD1	2.33	0.54
1:B:894:GLN:O	2:D:9:M3L:HM22	2.08	0.54
1:A:1020:ILE:HD12	4:A:98:HOH:O	2.06	0.54
1:A:1014:HIS:CD2	1:A:1016:GLN:H	2.27	0.53
1:A:1020:ILE:HD11	4:A:198:HOH:O	2.08	0.52
1:A:895:ASN:ND2	1:A:896:GLY:H	2.07	0.52
1:A:942:ASP:OD2	1:A:942:ASP:N	2.43	0.52
1:A:929:THR:HB	1:A:971:LEU:HD12	1.93	0.51
1:B:903:GLU:HG3	1:B:926:TRP:O	2.12	0.50
1:A:1014:HIS:HD2	1:A:1016:GLN:H	1.60	0.49
1:B:991:PRO:O	1:B:992:ASN:HB2	2.12	0.49
1:B:1067:LYS:HE3	1:B:1067:LYS:HB2	1.71	0.49
1:B:1020:ILE:HG22	1:B:1022:ASP:H	1.77	0.48
1:B:1044:VAL:O	1:B:1047:VAL:HG12	2.12	0.48
1:B:1061:GLU:H	1:B:1061:GLU:CD	2.03	0.48
1:B:1009:LYS:HE2	1:B:1027:ASP:OD2	2.14	0.47
1:B:938:GLU:H	1:B:938:GLU:HG2	1.43	0.47
1:B:889:TRP:CE3	2:D:9:M3L:HM23	2.50	0.46
1:B:888:ASP:OD1	2:D:4:LYS:NZ	2.48	0.46
1:B:933:ASP:HB3	1:B:936:LYS:O	2.15	0.46
2:D:14:ALY:HE2	2:D:14:ALY:HH31	1.66	0.46
1:A:989:SER:O	1:A:991:PRO:HD3	2.16	0.46
1:B:1014:HIS:CD2	1:B:1016:GLN:H	2.27	0.46
2:D:9:M3L:HM22	2:D:9:M3L:HD3	1.46	0.46
1:B:904:LYS:HB2	1:B:927:ILE:CG1	2.47	0.45
1:A:895:ASN:HD22	1:A:896:GLY:N	2.15	0.45
1:A:905:CYS:CB	1:A:906:PRO:HD2	2.45	0.45
1:A:1063:ALA:O	1:A:1067:LYS:HG3	2.17	0.45
1:B:930:PHE:CE2	1:B:971:LEU:HD13	2.52	0.45
1:A:1009:LYS:HE3	1:A:1016:GLN:HB3	1.99	0.45
1:A:919:LEU:N	1:A:919:LEU:HD12	2.33	0.44
1:B:889:TRP:CD2	2:D:9:M3L:HM23	2.52	0.44
1:B:1076:LYS:HD3	1:B:1076:LYS:HA	1.88	0.43
1:A:896:GLY:HA3	2:C:6:THR:HG22	2.00	0.42
1:A:938:GLU:OE1	1:A:938:GLU:N	2.48	0.42
1:A:983:GLN:O	1:A:1004:SER:HB3	2.19	0.42
1:A:1016:GLN:O	1:A:1017:HIS:C	2.57	0.42
1:B:895:ASN:ND2	1:B:896:GLY:N	2.56	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:916:PRO:CG	1:A:944:ASP:HA	2.47	0.42
1:B:963:GLN:O	1:B:967:GLU:HG3	2.20	0.42
1:B:1022:ASP:OD1	1:B:1084:ARG:NH2	2.50	0.42
1:B:896:GLY:O	2:D:8:ARG:NH1	2.53	0.42
1:B:1020:ILE:HG23	1:B:1021:PRO:CD	2.47	0.41
1:B:1074:GLU:O	1:B:1078:THR:HG23	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/207 (86%)	177 (99%)	2 (1%)	0	100	100
1	B	180/207 (87%)	175 (97%)	5 (3%)	0	100	100
2	C	7/20 (35%)	7 (100%)	0	0	100	100
2	D	16/20 (80%)	16 (100%)	0	0	100	100
All	All	382/454 (84%)	375 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/189 (89%)	154 (92%)	14 (8%)	11	3
1	B	169/189 (89%)	150 (89%)	19 (11%)	6	1
2	C	7/13 (54%)	5 (71%)	2 (29%)	0	0
2	D	13/13 (100%)	10 (77%)	3 (23%)	1	0
All	All	357/404 (88%)	319 (89%)	38 (11%)	6	1

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	895	ASN
1	A	900	LEU
1	A	918	LEU
1	A	942	ASP
1	A	968	ARG
1	A	971	LEU
1	A	977	GLU
1	A	978	LEU
1	A	1016	GLN
1	A	1020	ILE
1	A	1022	ASP
1	A	1045	VAL
1	A	1046	GLN
1	A	1061	GLU
2	C	5	GLN
2	C	6	THR
1	B	883	ASP
1	B	895	ASN
1	B	900	LEU
1	B	918	LEU
1	B	919	LEU
1	B	927	ILE
1	B	938	GLU
1	B	971	LEU
1	B	978	LEU
1	B	980	ILE
1	B	995	LYS
1	B	1012	LYS
1	B	1016	GLN
1	B	1020	ILE
1	B	1041	MET
1	B	1062	VAL
1	B	1067	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	1082	SER
1	B	1084	ARG
2	D	5	GLN
2	D	19	GLN
2	D	20	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	895	ASN
1	A	1014	HIS
1	A	1046	GLN
1	B	895	ASN
1	B	983	GLN
1	B	1014	HIS
1	B	1016	GLN
2	D	5	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	M3L	C	9	2	10,11,12	0.80	0	9,14,16	0.58	0
2	M3L	D	9	2	10,11,12	0.80	0	9,14,16	0.52	0
2	ALY	D	14	2	10,11,12	0.39	0	7,12,14	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M3L	C	9	2	-	7/9/10/12	-
2	M3L	D	9	2	-	5/9/10/12	-
2	ALY	D	14	2	-	3/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	9	M3L	N-CA-CB-CG
2	C	9	M3L	C-CA-CB-CG
2	D	14	ALY	OH-CH-NZ-CE
2	D	14	ALY	CH3-CH-NZ-CE
2	D	9	M3L	CG-CD-CE-NZ
2	C	9	M3L	CG-CD-CE-NZ
2	D	9	M3L	CE-CD-CG-CB
2	D	14	ALY	CE-CD-CG-CB
2	D	9	M3L	CD-CE-NZ-CM2
2	C	9	M3L	CD-CE-NZ-CM2
2	D	9	M3L	CD-CE-NZ-CM3
2	D	9	M3L	CD-CE-NZ-CM1
2	C	9	M3L	CD-CE-NZ-CM1
2	C	9	M3L	CD-CE-NZ-CM3

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	9	M3L	6	0
2	D	14	ALY	1	0

## 5.5 Carbohydrates

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/207 (89%)	0.33	8 (4%) 35 45	17, 29, 51, 78	0
1	B	186/207 (89%)	0.77	23 (12%) 4 6	21, 38, 67, 81	0
2	C	9/20 (45%)	1.63	2 (22%) 0 0	36, 49, 69, 85	0
2	D	18/20 (90%)	1.08	4 (22%) 0 0	31, 45, 62, 63	0
All	All	398/454 (87%)	0.60	37 (9%) 8 14	17, 35, 62, 85	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	10	SER	8.3
1	B	1087	ALA	6.8
1	A	942	ASP	6.3
1	B	919	LEU	5.3
1	B	1016	GLN	4.9
1	B	970	LEU	4.1
1	B	971	LEU	4.1
1	A	883	ASP	3.9
2	D	12	GLY	3.7
1	B	1048	TYR	3.6
1	A	943	CYS	3.6
1	B	1061	GLU	3.5
1	A	944	ASP	3.3
1	B	990	ILE	3.3
1	A	1087	ALA	3.2
1	B	1047	VAL	3.2
1	B	936	LYS	3.1
1	B	1013	LYS	3.0
1	B	943	CYS	2.9
1	A	939	VAL	2.9
1	B	1012	LYS	2.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	915	VAL	2.8
1	B	1015	SER	2.7
1	A	923	SER	2.6
2	C	1	ALA	2.6
1	B	939	VAL	2.4
1	B	944	ASP	2.4
1	B	1011	GLN	2.4
1	B	942	ASP	2.3
1	A	884	ASP	2.3
1	B	1010	LEU	2.3
2	D	19	GLN	2.3
1	B	989	SER	2.3
1	B	925	ASP	2.2
1	B	975	CYS	2.1
2	D	11	THR	2.1
2	D	20	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	M3L	C	9	12/13	0.81	0.23	68,76,84,87	0
2	M3L	D	9	12/13	0.91	0.15	27,38,69,69	0
2	ALY	D	14	12/13	0.95	0.12	25,35,41,44	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	B	2	1/1	0.96	0.06	40,40,40,40	0
3	ZN	B	1	1/1	0.98	0.04	37,37,37,37	0
3	ZN	A	1	1/1	0.99	0.03	35,35,35,35	0
3	ZN	A	2	1/1	0.99	0.05	36,36,36,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.