



wwPDB EM Validation Summary Report ⓘ

Feb 5, 2024 – 10:35 PM EST

PDB ID : 7U5C
EMDB ID : EMD-26346
Title : Cryo-EM structure of human CST bound to DNA polymerase alpha-primase in a recruitment state
Authors : Cai, S.W.; Zinder, J.C.; Svetlov, V.; Bush, M.W.; Nudler, E.; Walz, T.; de Lange, T.
Deposited on : 2022-03-02
Resolution : 4.60 Å (reported)
Based on initial models : 5EXR, 6W6W

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	
2	B	512	
3	C	1132	
4	D	598	
5	E	1221	
6	F	368	
7	G	123	
8	H	18	

2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 30077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA primase small subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	386	3243	2087	560	581	15	0	0

- Molecule 2 is a protein called DNA primase large subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	434	3562	2280	616	653	13	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P49643
B	-1	GLY	-	expression tag	UNP P49643
B	0	SER	-	expression tag	UNP P49643

- Molecule 3 is a protein called DNA polymerase alpha catalytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	1057	8546	5479	1433	1578	56	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	331	GLY	-	expression tag	UNP P09884
C	332	PRO	-	expression tag	UNP P09884
C	333	GLY	-	expression tag	UNP P09884
C	334	SER	-	expression tag	UNP P09884

- Molecule 4 is a protein called DNA polymerase alpha subunit B.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	444	Total	C	N	O	S	0	0
			3451	2194	576	666	15		

- Molecule 5 is a protein called CST complex subunit CTC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	1147	Total	C	N	O	S	0	0
			8966	5727	1580	1608	51		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-3	GLY	-	expression tag	UNP Q2NKJ3
E	-2	PRO	-	expression tag	UNP Q2NKJ3
E	-1	GLY	-	expression tag	UNP Q2NKJ3
E	0	SER	-	expression tag	UNP Q2NKJ3

- Molecule 6 is a protein called CST complex subunit STN1.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	155	Total	C	N	O	S	0	0
			1282	827	218	233	4		

- Molecule 7 is a protein called CST complex subunit TEN1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	117	Total	C	N	O	S	0	0
			930	588	167	169	6		

- Molecule 8 is a DNA chain called canonical telomeric DNA sequence.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	4	Total	C	N	O	P	0	0
			85	40	17	24	4		

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
9	A	1	Total	Zn	0
			1	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
9	C	2	2	2	0
9	E	1	1	1	0

- Molecule 10 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
			Total	Fe	S	
10	B	1	8	4	4	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA primase small subunit





- Molecule 2: DNA primase large subunit








• Molecule 3: DNA polymerase alpha catalytic subunit






• Molecule 4: DNA polymerase alpha subunit B





• Molecule 5: CST complex subunit CTC1



PRD	H726	H727	G728	Q729	S730	R731	L732	F733	L734	L735	H736	H737	K738	E739	A740	L741	M742	K743	R744	N745	F746	C747	V748	P749	P750	G751	A752	S753	P754	E755	V756	P757	K758	P759	A760	L761	S762	F763	Y764	V765	L766	G767	S768	W769	L770	G771	G772	T773	Q774	R775	K776	E777	G778	T779	G780	R781	G782	L783	P784
	E785	P786	Q787	G788	N789	D790	D791	N792	D793	Q794	K795	V796	L797	L798	L799	F800	F801	G802	S803	S804	V805	R806	W807	L811	H812	P813	G814	Q815	V816	Y817	R818	L819	I820	A821	P822	G823	P824	A825	T826	P827	M828	L829	F830	E831	K832	D833	G834	S835	S836	C837	I838	S839	R840	R841	P842	L843	E844	A846	
	G847	C848	A849	S850	C851	L852	T853	V854	Q855	D856	N857	W858	T859	L860	E861	L862	S863	S864	S865	Q866	D867	I868	Q869	D870	V871	L872	D873	A874	N875	K876	S877	L878	P879	E880	S881	S882	L883	L886	L887	S888	D889	N890	F891	T892	D893	S894	L895	V896	S897	F898	A900	L903	S904	R905	T906	L907	C908		
	E909	P910	L911	V912	A913	S914	L915	N916	N917	K918	L919	G920	N921	N922	G923	A924	N925	R926	R927	C928	T932	V933	A934	L935	E936	T937	A938	E939	C940	E941	F942	P943	H945	I950	E951	D952	P953	H954	L955	S958	L959	L962	P963	G964	R966	F969	S970	Q971	L972	E973	K974	R975	V976						
	S977	R978	S979	H980	Y983	C984	C985	F986	S987	R988	S989	T990	L995	S996	F997	P998	P999	E1000	T1001	G1002	I1003	S1004	I1005	P1006	L1007	H1009	I1010	E1011	L1012	A1013	E1014	L1015	L1016	Q1017	G1018	G1019	Q1020	S1021	P1022	F1023	Q1024	A1025	T1026	A1027	S1028	C1029	H1030	I1031	V1032	S1033	V1034	F1035	S1036	L1037	Q1038	L1039			
	F1040	W1041	V1042	C1043	A1044	T1045	C1046	T1047	S1048	I1049	V1113	C1050	R1051	Q1052	C1055	T1056	R1057	L1058	GLY	T1061	C1062	P1063	T1064	Q1065	T1066	A1067	I1068	S1069	Q1070	A1071	I1072	I1073	R1074	L1075	L1076	E1078	D1079	T1081	A1082	E1083	A1084	V1085	V1086	T1087	C1088	N1089	H1091	H1092	A1095	A1096	L1097	L1098	C1100	P1101					
	R1102	E1103	W1104	L1107	D1109	V1111	Q1112	P1114	G1115	R1116	V1118	L1119	Q1120	F1121	A1122	G1123	P1124	G1125	Q1126	L1128	E1129	S1130	S1131	A1132	R1133	V1134	D1135	E1136	P1137	M1138	T1139	M1140	F1141	L1145	S1148	C1146	T1147	S1149	S1150	V1151	L1152	R1153	P1154	L1155	V1156	L1157	F1159	E1160	L1161	E1162	R1163	K1164							
	P1165	S1166	K1167	I1168	V1169	P1170	L1171	E1172	P1173	P1174	R1175	L1176	Q1177	R1178	F1179	Q1180	G1181	G1182	E1183	L1184	P1185	F1186	L1187	T1188	H1189	V1190	M1191	P1192	R1193	L1194	R1195	L1196	S1197	C1198	L1199	S1200	I1201	R1202	E1203	S1204	E1205	Y1206	S1207	S1208	S1209	L1210	G1211	I1212	L1213	A1214	S1215	S1216	C1217						

• Molecule 6: CST complex subunit STN1





MET	GLN	PRO	GLY	SER	ARG	CYS	GLU	GLU	E11	T12	L15	L16	W17	GLY	L19	D20	P21	V22	F23	L24	A25	F26	A27	K28	L29	X30	I31	R32	D33	L34	L35	D36	K37	R38	E39	S40	R41	Q42	V43	P44	G45	V46	F47	L48	Y49	N50	G51	H52	P53	I54	K55	O56	V57	D58	V59	L60	G61	
T62	V63	I64	G65	V66	R67	E68	R69	D70	A71	F72	V73	S74	Y75	D78	D79	S80	T81	G82	V83	C86	L87	C88	W89	R90	K91	LEU	ASN	THR	THR	VAL	L36	SER	VAL	SER	TYR	ALA	ALA	PRO	SER	ALA	ALA	ARG	LEU	L109	T110	S111	Q112	L113	K114	K115	L116	Q117	L119	I120	E121	Q122	K123	
T124	K125	I126	E127	L128	G129	D130	T131	I132	R133	V134	R135	G136	S137	I138	R139	T140	R141	R142	E143	E144	R145	E146	I147	H148	A149	T150	T151	Y152	K153	L154	V155	D156	D157	P158	V159	M160	M161	I162	Q163	I164	A165	R166	M167	L168	E169	P171	T172	I173	Y174	R175	K176	V177	V178	D179	F182	H183	SER	
SER	ALA	LEU	GLU	SER	LYS	GLU	ALA	LEU	SER	ALA	THR	SER	ASP	LEU	PRO	LEU	THR	ILE	LEU	LEU	GLU	ILE	GLN	PHE	GLY	MET	GLU	ASN	VAL	PHE	VAL	GLN	SER	PHE	TYR	GLN	ALA	LEU	LEU	GLU	LEU	VAL	VAL	THR	ARG	GLU	SER	ASP	LEU	ALA	LEU	ASN	GLN	PRO	VAL	LYS	VAL	ILE
HIS	SER	ALA	SER	SER	ASP	GLN	VAL	ASN	PHE	LYS	ASP	THR	THR	SER	ALA	ALA	HIS	SER	ILE	PHE	LYS	ASN	ALA	ILE	GLN	LEU	GLU	GLY	LEU	VAL	PHE	GLN	LYS	ASP	GLY	PHE	ASN	LEU	LEU	TYR	TYR	VAL	THR	ARG	GLU	SER	ASP	LEU	ALA	LEU	ASN	GLN	PRO	VAL	LYS	VAL	ILE	

HIS ARG ARG ILE ILE GLN GLN ASP CYS GLN LYS PRO ASN HIS MET MET GLU LYS GLY CYS HIS PHE LEU HIS ILE LEU ALA CYS ARG ARG SER ILE ARG PRO GLY LEU SER GLU ALA VAL LEU GLN VAL LEU LEU LEU LEU LEU LEU LEU ASP GLN SER SER ILE VAL THR MET GLU HIS TYR

TYR THR ALA PHE

● Molecule 7: CST complex subunit TEN1



GLY SER GLN

● Molecule 8: canonical telomeric DNA sequence



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	131850	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	52	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.084	Depositor
Minimum map value	-0.041	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size (\AA)	324.0, 324.0, 324.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.08, 1.08, 1.08	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3327	0.46	0/4488
2	B	0.36	0/3646	0.52	0/4908
3	C	0.37	0/8726	0.51	0/11791
4	D	0.44	0/3529	0.56	0/4795
5	E	0.32	1/9196 (0.0%)	0.53	0/12528
6	F	0.27	0/1309	0.51	0/1771
7	G	0.26	0/948	0.50	0/1284
8	H	0.64	0/95	0.92	0/145
All	All	0.35	1/30776 (0.0%)	0.52	0/41710

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	167	PRO	C-N	8.42	1.50	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3243	0	3221	207	0
2	B	3562	0	3544	197	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	8546	0	8636	377	0
4	D	3451	0	3425	170	0
5	E	8966	0	9050	612	0
6	F	1282	0	1296	107	0
7	G	930	0	941	59	0
8	H	85	0	46	9	0
9	A	1	0	0	0	0
9	C	2	0	0	0	0
9	E	1	0	0	0	0
10	B	8	0	0	0	0
All	All	30077	0	30159	1669	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 1669 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:165:TYR:CD1	5:E:179:LEU:HG	1.24	1.69
5:E:165:TYR:HE1	5:E:179:LEU:CD1	1.24	1.50
5:E:81:CYS:SG	5:E:163:TRP:CZ3	2.05	1.49
5:E:165:TYR:CE1	5:E:179:LEU:CD1	2.09	1.33
5:E:73:LEU:HD13	5:E:165:TYR:CD2	1.61	1.33

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	382/420 (91%)	339 (89%)	43 (11%)	0	100 100
2	B	432/512 (84%)	343 (79%)	88 (20%)	1 (0%)	47 81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	1047/1132 (92%)	886 (85%)	161 (15%)	0	100	100
4	D	442/598 (74%)	357 (81%)	85 (19%)	0	100	100
5	E	1137/1221 (93%)	938 (82%)	197 (17%)	2 (0%)	47	81
6	F	149/368 (40%)	132 (89%)	17 (11%)	0	100	100
7	G	115/123 (94%)	100 (87%)	15 (13%)	0	100	100
All	All	3704/4374 (85%)	3095 (84%)	606 (16%)	3 (0%)	54	85

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	251	ASN
5	E	436	ARG
5	E	78	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/393 (92%)	361 (100%)	0	100	100
2	B	394/460 (86%)	392 (100%)	2 (0%)	88	93
3	C	963/1016 (95%)	955 (99%)	8 (1%)	81	89
4	D	390/527 (74%)	388 (100%)	2 (0%)	88	93
5	E	1005/1066 (94%)	995 (99%)	10 (1%)	76	86
6	F	142/332 (43%)	142 (100%)	0	100	100
7	G	103/107 (96%)	102 (99%)	1 (1%)	76	86
All	All	3358/3901 (86%)	3335 (99%)	23 (1%)	84	90

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	101	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	396	ARG
5	E	106	LEU
5	E	435	SER
3	C	913	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 75 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	386	GLN
6	F	161	ASN
5	E	484	HIS
5	E	708	HIS
3	C	637	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	SF4	B	601	-	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	SF4	B	601	-	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-26346. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 142




Y Index: 151




Z Index: 137


6.3.2 Raw map



X Index: 142



Y Index: 151




Z Index: 136


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 214 nm³; this corresponds to an approximate mass of 193 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.217 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.217 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.60	-	-
Author-provided FSC curve	4.61	7.22	4.70
Unmasked-calculated*	6.15	8.12	6.37

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.15 differs from the reported value 4.6 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-26346 and PDB model 7U5C. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).



















9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 62% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6210	 0.3300
A	 0.5400	 0.2870
B	 0.7910	 0.3640
C	 0.7660	 0.3670
D	 0.8420	 0.4110
E	 0.4640	 0.2950
F	 0.2770	 0.2150
G	 0.1090	 0.2120
H	 0.1650	 0.2390

