



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2023 – 01:32 am GMT

PDB ID : 4U9H
Title : Ultra High Resolution Structure Of The Ni-R State Of [Nife]Hydrogenase From Desulfovibrio Vulgaris Miyazaki F
Authors : Ogata, H.; Nishikawa, K.; Lubitz, W.
Deposited on : 2014-08-06
Resolution : 0.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 0.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1061 (1.04-0.76)
Clashscore	141614	1132 (1.04-0.76)
Ramachandran outliers	138981	1055 (1.04-0.76)
Sidechain outliers	138945	1056 (1.04-0.76)
RSRZ outliers	127900	1028 (1.04-0.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	S	265	 92% 8%
2	L	533	 93% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	L	607	-	-	X	-
5	MPD	L	608	-	-	X	-
5	MPD	L	609	-	-	X	-
5	MPD	L	610	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 14147 atoms, of which 6616 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Periplasmic [NiFe] hydrogenase small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	S	265	4159	1345	2036	355	405	18	25	14	0

- Molecule 2 is a protein called Periplasmic [NiFe] hydrogenase large subunit.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	L	533	8692	2792	4331	756	797	16	80	23	0

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe S		
3	S	1	8	4 4	0	0
3	S	1	8	4 4	0	0

- Molecule 4 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe₃S₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	S	1	Total	Fe S	0	0
			7	3 4		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	S	1	Total	C O	0	0
			8	6 2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	S	1	Total	C	O	0	0
			8	6	2		
5	S	1	Total	C	O	0	0
			8	6	2		
5	S	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		
5	L	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is hydrido[hydridonickel(2+)]bis(hydrocyanato-1kappaC)(hydroxymethyl)iron (three-letter code: NWN) (formula: C₃H₄FeN₂NiO).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	Fe	H	N	Ni			O
6	L	1	9	3	1	1	2	1	1	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	1	Total	Mg	0	0
			1	1		

- Molecule 8 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	L	1	8	4	1	3	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
9	S	320	406	86	320	0	0
9	L	591	753	162	591	2	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Periplasmic [NiFe] hydrogenase small subunit



- Molecule 2: Periplasmic [NiFe] hydrogenase large subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.53Å 97.95Å 125.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.64 – 0.89 77.30 – 0.89	Depositor EDS
% Data completeness (in resolution range)	95.8 (37.64-0.89) 91.0 (77.30-0.89)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.82 (at 0.89Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9pre_1669)	Depositor
R, R_{free}	0.096 , 0.106 0.096 , 0.106	Depositor DCC
R_{free} test set	30295 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	7.6	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 68.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	14147	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, F3S, SF4, NWN, MG, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	S	0.87	10/2181 (0.5%)	0.74	1/2973 (0.0%)
2	L	0.91	16/4476 (0.4%)	0.77	7/6078 (0.1%)
All	All	0.90	26/6657 (0.4%)	0.76	8/9051 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	S	0	1
2	L	0	2
All	All	0	3

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	418	GLU	CD-OE1	17.80	1.45	1.25
2	L	42[A]	GLU	CD-OE1	-16.81	1.07	1.25
2	L	42[B]	GLU	CD-OE1	-16.81	1.07	1.25
2	L	464	GLU	CD-OE2	-15.77	1.08	1.25
1	S	172[A]	GLU	CD-OE1	15.33	1.42	1.25

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	350	ARG	NE-CZ-NH2	-6.39	117.11	120.30
2	L	199	TYR	CB-CG-CD2	-5.93	117.44	121.00
2	L	350	ARG	NE-CZ-NH1	5.70	123.15	120.30
2	L	426	ARG	NE-CZ-NH1	5.62	123.11	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	S	5	ARG	NE-CZ-NH2	-5.49	117.56	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	120	TYR	Sidechain
2	L	79	ARG	Sidechain
1	S	26	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	2123	2036	2020	10	0
2	L	4361	4331	4303	35	0
3	S	16	0	0	0	0
4	S	7	0	0	0	0
5	L	64	0	112	35	0
5	S	32	0	56	3	0
6	L	8	1	0	0	0
7	L	1	0	0	0	0
8	L	8	0	12	5	0
9	L	591	162	0	25	0
9	S	320	86	0	5	0
All	All	7531	6616	6503	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:608:MPD:H52	5:L:608:MPD:HM2	1.35	1.05
5:L:604:MPD:H51	9:L:955:HOH:O	1.56	1.05

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:607:MPD:CM	9:L:777:HOH:O	2.13	0.94
1:S:54[B]:ASP:OD1	9:S:1101:HOH:O	1.90	0.88
2:L:474:PHE:CD2	5:L:607:MPD:H13	2.09	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	277/265 (104%)	271 (98%)	6 (2%)	0	100	100
2	L	554/533 (104%)	538 (97%)	16 (3%)	0	100	100
All	All	831/798 (104%)	809 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	225/211 (107%)	224 (100%)	1 (0%)	91	65
2	L	460/437 (105%)	458 (100%)	2 (0%)	91	65
All	All	685/648 (106%)	682 (100%)	3 (0%)	88	65

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	138	LYS
2	L	161	LYS
2	L	492	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 1 is monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MPD	L	605	-	7,7,7	0.26	0	9,10,10	0.78	0
5	MPD	L	610	-	7,7,7	0.70	0	9,10,10	1.11	1 (11%)
5	MPD	S	1006	-	7,7,7	0.30	0	9,10,10	0.67	0
5	MPD	S	1007	-	7,7,7	0.22	0	9,10,10	0.69	0
5	MPD	L	606	-	7,7,7	0.48	0	9,10,10	0.77	0
5	MPD	S	1004	-	7,7,7	0.41	0	9,10,10	0.73	0
3	SF4	S	1001	1	0,12,12	-	-	-	-	-
3	SF4	S	1002	1	0,12,12	-	-	-	-	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MPD	L	607	-	7,7,7	0.92	1 (14%)	9,10,10	0.99	0
5	MPD	L	608	-	7,7,7	0.24	0	9,10,10	0.70	0
4	F3S	S	1003	1	0,9,9	-	-	-	-	-
6	NWN	L	601	2	1,6,6	8.04	1 (100%)	-	-	-
5	MPD	L	609	-	7,7,7	0.31	0	9,10,10	0.95	0
8	TRS	L	611	-	7,7,7	0.70	0	9,9,9	1.26	1 (11%)
5	MPD	S	1005	-	7,7,7	0.43	0	9,10,10	0.49	0
5	MPD	L	603	-	7,7,7	0.38	0	9,10,10	0.34	0
5	MPD	L	604	-	7,7,7	0.52	0	9,10,10	2.33	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	L	605	-	-	1/5/5/5	-
5	MPD	L	610	-	-	1/5/5/5	-
5	MPD	S	1006	-	-	3/5/5/5	-
5	MPD	S	1007	-	-	0/5/5/5	-
5	MPD	L	606	-	-	1/5/5/5	-
5	MPD	S	1004	-	-	0/5/5/5	-
3	SF4	S	1001	1	-	-	0/6/5/5
5	MPD	L	607	-	-	1/5/5/5	-
5	MPD	L	608	-	-	1/5/5/5	-
3	SF4	S	1002	1	-	-	0/6/5/5
4	F3S	S	1003	1	-	-	0/3/3/3
5	MPD	L	609	-	-	2/5/5/5	-
8	TRS	L	611	-	-	0/9/9/9	-
5	MPD	S	1005	-	-	0/5/5/5	-
5	MPD	L	603	-	-	0/5/5/5	-
5	MPD	L	604	-	-	2/5/5/5	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	L	601	NWN	O3-C3	-8.04	1.15	1.40
5	L	607	MPD	CM-C2	-2.08	1.46	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	604	MPD	O2-C2-C1	-4.80	92.66	108.08
5	L	604	MPD	O2-C2-C3	3.85	124.27	109.80
8	L	611	TRS	O3-C3-C	3.04	120.62	111.00
5	L	604	MPD	CM-C2-C1	-2.82	104.71	110.57
5	L	610	MPD	O2-C2-C1	2.01	114.52	108.08

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	S	1006	MPD	C1-C2-C3-C4
5	S	1006	MPD	O2-C2-C3-C4
5	L	604	MPD	C2-C3-C4-C5
5	L	606	MPD	C2-C3-C4-O4
5	L	609	MPD	C2-C3-C4-O4

There are no ring outliers.

9 monomers are involved in 43 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	605	MPD	3	0
5	L	610	MPD	1	0
5	S	1006	MPD	1	0
5	S	1007	MPD	2	0
5	L	607	MPD	15	0
5	L	608	MPD	6	0
5	L	609	MPD	6	0
8	L	611	TRS	5	0
5	L	604	MPD	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9	
1	S	265/265 (100%)	-0.92	2 (0%)	86 72	4, 9, 19, 31	0
2	L	533/533 (100%)	-0.94	1 (0%)	95 82	4, 9, 18, 31	0
All	All	798/798 (100%)	-0.93	3 (0%)	92 80	4, 9, 19, 31	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	3	GLY	5.2
2	L	161	LYS	3.0
1	S	4	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MPD	L	607	8/8	0.49	0.24	14,19,19,19	5
5	MPD	L	610	8/8	0.67	0.41	20,20,20,20	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MPD	L	604	8/8	0.71	0.25	17,19,20,20	6
5	MPD	L	606	8/8	0.81	0.16	18,19,20,20	8
5	MPD	L	608	8/8	0.85	0.12	17,18,19,19	5
5	MPD	S	1007	8/8	0.86	0.33	19,19,20,20	8
5	MPD	L	609	8/8	0.88	0.12	18,19,19,19	8
5	MPD	S	1006	8/8	0.90	0.28	19,20,20,20	8
5	MPD	L	605	8/8	0.92	0.12	12,18,19,19	3
8	TRS	L	611	8/8	0.92	0.13	14,18,19,19	7
5	MPD	S	1004	8/8	0.95	0.09	12,18,19,20	2
5	MPD	S	1005	8/8	0.96	0.11	13,17,18,19	3
5	MPD	L	603	8/8	0.96	0.09	12,14,16,19	0
3	SF4	S	1002	8/8	1.00	0.04	5,6,6,6	0
4	F3S	S	1003	7/7	1.00	0.04	4,5,5,5	0
6	NWN	L	601	8/8	1.00	0.04	3,4,5,5	0
7	MG	L	602	1/1	1.00	0.04	4,4,4,4	0
3	SF4	S	1001	8/8	1.00	0.04	4,4,4,4	0

6.5 Other polymers [i](#)

There are no such residues in this entry.