



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 06:54 AM EDT

PDB ID : 5UAE  
Title : Crystal structure of the coiled-coil domain from Listeria Innocua Phage Integrase (Trigonal Form)  
Authors : Gupta, K.; Sharp, R.; Yuan, J.B.; Van Duyne, G.D.  
Deposited on : 2016-12-19  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	62	 6% 85% 15%
1	B	62	 5% 68% 23% 10%
1	C	62	 10% 85% 13% .
1	D	62	 5% 74% 21% 5%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2286 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative integrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	62	Total 515	C 322	N 82	O 109	S 2	0	0	0
1	B	56	Total 463	C 289	N 73	O 99	S 2	0	0	0
1	C	61	Total 506	C 316	N 80	O 108	S 2	0	0	0
1	D	59	Total 488	C 305	N 77	O 104	S 2	0	0	0

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 13	C 6	O 7	0	0
2	A	1	Total 13	C 6	O 7	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			13	6	7		
2	D	1	Total	C	O	0	0
			13	6	7		
2	D	1	Total	C	O	0	0
			13	6	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	72	Total	O	0	0
			72	72		
3	B	53	Total	O	0	0
			53	53		
3	C	51	Total	O	0	0
			51	51		
3	D	73	Total	O	0	0
			73	73		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative integrase




- Molecule 1: Putative integrase



- Molecule 1: Putative integrase



- Molecule 1: Putative integrase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.43Å 75.43Å 103.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.72 – 2.75 40.46 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.7 (37.72-2.75) 96.4 (40.46-2.75)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.59 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.216 , 0.250 0.209 , 0.242	Depositor DCC
$R_{free}$ test set	1700 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	87.0	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 91.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.479 for -h,-k,l 0.058 for h,-h-k,-l 0.057 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2286	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/520	0.36	0/694
1	B	0.22	0/468	0.37	0/627
1	C	0.22	0/511	0.42	0/683
1	D	0.22	0/493	0.38	0/660
All	All	0.22	0/1992	0.38	0/2664

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	515	0	500	7	0
1	B	463	0	440	9	0
1	C	506	0	487	5	0
1	D	488	0	468	8	0
2	A	26	0	10	3	0
2	D	39	0	15	0	0
3	A	72	0	0	1	0
3	B	53	0	0	0	0
3	C	51	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	73	0	0	1	0
All	All	2286	0	1920	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:364:ARG:NH1	1:D:370:ILE:O	2.26	0.68
1:B:361:LYS:HD3	1:C:369:TYR:HB3	1.75	0.67
1:A:364:ARG:NH2	1:B:374:TYR:O	2.29	0.65
1:A:396:GLN:O	1:A:400:ASN:ND2	2.30	0.64
1:C:396:GLN:O	1:C:400:ASN:ND2	2.30	0.64
1:A:361:LYS:NZ	2:A:501:FLC:OB2	2.32	0.61
1:B:380:ASP:O	1:B:384:ASN:ND2	2.33	0.61
1:D:361:LYS:NZ	3:D:602:HOH:O	2.37	0.57
1:B:402:GLU:N	1:B:402:GLU:OE2	2.41	0.53
1:B:367:ASP:O	1:B:371:ASN:ND2	2.32	0.52
1:A:357:ILE:HG22	1:A:361:LYS:HE3	1.93	0.50
1:B:347:GLU:O	1:B:350:SER:OG	2.21	0.50
1:D:372:GLY:O	1:D:374:TYR:N	2.40	0.49
1:D:358:GLU:HA	1:D:361:LYS:HD3	1.94	0.49
1:A:374:TYR:OH	1:B:380:ASP:OD1	2.32	0.47
1:D:366:PHE:O	1:D:370:ILE:HG12	2.15	0.47
1:A:362:LYS:NZ	3:A:601:HOH:O	2.48	0.46
1:B:393:TYR:O	1:B:397:ILE:HG12	2.17	0.45
1:B:378:GLU:O	1:B:382:MET:HG3	2.17	0.44
1:D:403:LEU:N	1:D:404:LYS:HA	2.31	0.44
1:A:373:SER:O	2:A:502:FLC:OHB	2.33	0.44
2:A:502:FLC:OHB	2:A:502:FLC:OG2	2.31	0.44
1:C:371:ASN:ND2	1:C:372:GLY:O	2.53	0.42
1:D:355:LEU:HD11	1:D:390:ILE:HG23	2.01	0.42
1:C:355:LEU:HD11	1:C:394:GLU:HG3	2.01	0.42
1:D:378:GLU:O	1:D:382:MET:HG3	2.21	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	60/62 (97%)	56 (93%)	4 (7%)	0	100	100
1	B	54/62 (87%)	51 (94%)	3 (6%)	0	100	100
1	C	59/62 (95%)	57 (97%)	2 (3%)	0	100	100
1	D	57/62 (92%)	54 (95%)	2 (4%)	1 (2%)	8	15
All	All	230/248 (93%)	218 (95%)	11 (5%)	1 (0%)	34	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	373	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	57/57 (100%)	56 (98%)	1 (2%)	59	75
1	B	51/57 (90%)	50 (98%)	1 (2%)	55	72
1	C	56/57 (98%)	56 (100%)	0	100	100
1	D	54/57 (95%)	54 (100%)	0	100	100
All	All	218/228 (96%)	216 (99%)	2 (1%)	78	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	403	LEU
1	B	389	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FLC	D	501	-	12,12,12	1.24	0	17,17,17	1.50	1 (5%)
2	FLC	D	503	-	12,12,12	1.26	0	17,17,17	1.50	1 (5%)
2	FLC	A	501	-	12,12,12	1.25	0	17,17,17	1.41	1 (5%)
2	FLC	A	502	-	12,12,12	1.22	0	17,17,17	1.48	1 (5%)
2	FLC	D	502	-	12,12,12	1.31	0	17,17,17	1.62	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	D	501	-	-	2/16/16/16	-
2	FLC	D	503	-	-	2/16/16/16	-
2	FLC	A	501	-	-	0/16/16/16	-
2	FLC	A	502	-	-	13/16/16/16	-
2	FLC	D	502	-	-	7/16/16/16	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	502	FLC	OB2-CBC-CB	4.13	120.22	113.05
2	D	501	FLC	OB2-CBC-CB	4.11	120.18	113.05
2	A	502	FLC	OB2-CBC-CB	4.03	120.05	113.05
2	D	503	FLC	OB2-CBC-CB	4.01	120.02	113.05
2	A	501	FLC	OB2-CBC-CB	3.79	119.63	113.05
2	D	502	FLC	CB-CG-CGC	-2.88	106.83	113.81

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	FLC	CA-CB-CBC-OB1
2	A	502	FLC	CA-CB-CBC-OB2
2	A	502	FLC	OHB-CB-CBC-OB1
2	A	502	FLC	OHB-CB-CBC-OB2
2	D	502	FLC	CA-CB-CBC-OB1
2	D	502	FLC	CA-CB-CBC-OB2
2	D	502	FLC	OHB-CB-CBC-OB1
2	D	502	FLC	OHB-CB-CBC-OB2
2	D	502	FLC	CA-CB-CG-CGC
2	D	502	FLC	CBC-CB-CG-CGC
2	D	502	FLC	OHB-CB-CG-CGC
2	A	502	FLC	CG-CB-CBC-OB1
2	A	502	FLC	CG-CB-CBC-OB2
2	A	502	FLC	CBC-CB-CG-CGC
2	A	502	FLC	CB-CA-CAC-OA1
2	A	502	FLC	CB-CA-CAC-OA2
2	A	502	FLC	OHB-CB-CG-CGC
2	A	502	FLC	CB-CG-CGC-OG2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	502	FLC	CB-CG-CGC-OG1
2	A	502	FLC	CA-CB-CG-CGC
2	D	501	FLC	CB-CG-CGC-OG1
2	D	501	FLC	CB-CG-CGC-OG2
2	D	503	FLC	CB-CG-CGC-OG1
2	D	503	FLC	CB-CG-CGC-OG2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FLC	1	0
2	A	502	FLC	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	62/62 (100%)	0.74	4 (6%) 18 22	57, 89, 130, 151	0
1	B	56/62 (90%)	0.54	3 (5%) 25 31	57, 85, 108, 122	0
1	C	61/62 (98%)	0.69	6 (9%) 7 8	63, 89, 143, 184	0
1	D	59/62 (95%)	0.65	3 (5%) 28 34	58, 87, 110, 112	0
All	All	238/248 (95%)	0.66	16 (6%) 17 21	57, 88, 132, 184	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	400	ASN	4.2
1	D	351	LEU	3.6
1	B	348	LEU	3.1
1	C	355	LEU	3.0
1	A	344	LYS	2.9
1	D	393	TYR	2.8
1	A	393	TYR	2.7
1	C	399	ALA	2.7
1	D	386	ILE	2.7
1	A	345	GLU	2.6
1	C	392	TYR	2.6
1	C	405	LYS	2.6
1	B	355	LEU	2.5
1	A	351	LEU	2.4
1	C	345	GLU	2.2
1	B	370	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FLC	D	502	13/13	0.74	0.36	149,204,222,225	0
2	FLC	A	502	13/13	0.86	0.25	120,169,187,192	0
2	FLC	A	501	13/13	0.88	0.17	129,143,152,156	0
2	FLC	D	501	13/13	0.90	0.28	100,140,183,190	0
2	FLC	D	503	13/13	0.93	0.10	90,103,119,121	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.