



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 29, 2024 – 09:09 AM EST

PDB ID : 5UDR
Title : LarE, a sulfur transferase involved in synthesis of the cofactor for lactate racemase in complex with nicotinamide mononucleotide NMN
Authors : Fellner, M.; Desguin, B.; Hausinger, R.P.; Hu, J.
Deposited on : 2016-12-28
Resolution : 2.62 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 76%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div>
1	B	286	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 74%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: grey;"></div> </div>
1	C	286	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: grey;"></div> </div>
1	D	286	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 72%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 19%; height: 10px; background-color: grey;"></div> </div>
1	E	286	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 78%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	286	 <p>A horizontal bar chart showing the quality distribution of chain F. The bar is divided into four segments: a small red segment (3%), a large green segment (73%), a yellow segment (14%), and a grey segment (13%).</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-utilizing enzyme of the PP-loopsuperfamily.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	257	Total 1934	C 1219	N 336	O 372	S 7	0	0	0
1	B	243	Total 1863	C 1174	N 324	O 360	S 5	0	0	0
1	C	247	Total 1871	C 1176	N 325	O 364	S 6	0	0	0
1	D	232	Total 1631	C 1017	N 292	O 318	S 4	0	0	0
1	E	260	Total 1923	C 1211	N 332	O 373	S 7	0	0	0
1	F	249	Total 1899	C 1196	N 329	O 368	S 6	0	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	277	ALA	-	expression tag	UNP A0A0G9FES3
A	278	SER	-	expression tag	UNP A0A0G9FES3
A	279	TRP	-	expression tag	UNP A0A0G9FES3
A	280	SER	-	expression tag	UNP A0A0G9FES3
A	281	HIS	-	expression tag	UNP A0A0G9FES3
A	282	PRO	-	expression tag	UNP A0A0G9FES3
A	283	GLN	-	expression tag	UNP A0A0G9FES3
A	284	PHE	-	expression tag	UNP A0A0G9FES3
A	285	GLU	-	expression tag	UNP A0A0G9FES3
A	286	LYS	-	expression tag	UNP A0A0G9FES3
B	277	ALA	-	expression tag	UNP A0A0G9FES3
B	278	SER	-	expression tag	UNP A0A0G9FES3
B	279	TRP	-	expression tag	UNP A0A0G9FES3
B	280	SER	-	expression tag	UNP A0A0G9FES3
B	281	HIS	-	expression tag	UNP A0A0G9FES3
B	282	PRO	-	expression tag	UNP A0A0G9FES3
B	283	GLN	-	expression tag	UNP A0A0G9FES3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	284	PHE	-	expression tag	UNP A0A0G9FES3
B	285	GLU	-	expression tag	UNP A0A0G9FES3
B	286	LYS	-	expression tag	UNP A0A0G9FES3
C	277	ALA	-	expression tag	UNP A0A0G9FES3
C	278	SER	-	expression tag	UNP A0A0G9FES3
C	279	TRP	-	expression tag	UNP A0A0G9FES3
C	280	SER	-	expression tag	UNP A0A0G9FES3
C	281	HIS	-	expression tag	UNP A0A0G9FES3
C	282	PRO	-	expression tag	UNP A0A0G9FES3
C	283	GLN	-	expression tag	UNP A0A0G9FES3
C	284	PHE	-	expression tag	UNP A0A0G9FES3
C	285	GLU	-	expression tag	UNP A0A0G9FES3
C	286	LYS	-	expression tag	UNP A0A0G9FES3
D	277	ALA	-	expression tag	UNP A0A0G9FES3
D	278	SER	-	expression tag	UNP A0A0G9FES3
D	279	TRP	-	expression tag	UNP A0A0G9FES3
D	280	SER	-	expression tag	UNP A0A0G9FES3
D	281	HIS	-	expression tag	UNP A0A0G9FES3
D	282	PRO	-	expression tag	UNP A0A0G9FES3
D	283	GLN	-	expression tag	UNP A0A0G9FES3
D	284	PHE	-	expression tag	UNP A0A0G9FES3
D	285	GLU	-	expression tag	UNP A0A0G9FES3
D	286	LYS	-	expression tag	UNP A0A0G9FES3
E	277	ALA	-	expression tag	UNP A0A0G9FES3
E	278	SER	-	expression tag	UNP A0A0G9FES3
E	279	TRP	-	expression tag	UNP A0A0G9FES3
E	280	SER	-	expression tag	UNP A0A0G9FES3
E	281	HIS	-	expression tag	UNP A0A0G9FES3
E	282	PRO	-	expression tag	UNP A0A0G9FES3
E	283	GLN	-	expression tag	UNP A0A0G9FES3
E	284	PHE	-	expression tag	UNP A0A0G9FES3
E	285	GLU	-	expression tag	UNP A0A0G9FES3
E	286	LYS	-	expression tag	UNP A0A0G9FES3
F	277	ALA	-	expression tag	UNP A0A0G9FES3
F	278	SER	-	expression tag	UNP A0A0G9FES3
F	279	TRP	-	expression tag	UNP A0A0G9FES3
F	280	SER	-	expression tag	UNP A0A0G9FES3
F	281	HIS	-	expression tag	UNP A0A0G9FES3
F	282	PRO	-	expression tag	UNP A0A0G9FES3
F	283	GLN	-	expression tag	UNP A0A0G9FES3
F	284	PHE	-	expression tag	UNP A0A0G9FES3
F	285	GLU	-	expression tag	UNP A0A0G9FES3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	286	LYS	-	expression tag	UNP A0A0G9FES3

- Molecule 2 is BETA-NICOTINAMIDE RIBOSE MONOPHOSPHATE (three-letter code: NMN) (formula: $C_{11}H_{16}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
22	11	2	8	1					
2	B	1	Total	C	N	O	P	0	0
22	11	2	8	1					
2	C	1	Total	C	N	O	P	0	0
22	11	2	8	1					
2	D	1	Total	C	N	O	P	0	0
22	11	2	8	1					
2	E	1	Total	C	N	O	P	0	0
22	11	2	8	1					
2	F	1	Total	C	N	O	P	0	0
22	11	2	8	1					

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
1	1					
3	D	1	Total	Mn	0	0
1	1					


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	19	Total O 19 19	0	0
4	B	51	Total O 51 51	0	0
4	C	15	Total O 15 15	0	0
4	D	14	Total O 14 14	0	0
4	E	17	Total O 17 17	0	0
4	F	20	Total O 20 20	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-utilizing enzyme of the PP-loopsuperfamily




- Molecule 1: ATP-utilizing enzyme of the PP-loopsuperfamily




- Molecule 1: ATP-utilizing enzyme of the PP-loopsuperfamily





- Molecule 1: ATP-utilizing enzyme of the PP-loopsuperfamily







● Molecule 1: ATP-utilizing enzyme of the PP-loopsuperfamily



● Molecule 1: ATP-utilizing enzyme of the PP-loopsuperfamily



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	108.09Å 108.09Å 324.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.42 – 2.62 48.42 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.42-2.62) 99.9 (48.42-2.62)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.11.1-2575	Depositor
R, R_{free}	0.215 , 0.246 0.221 , 0.255	Depositor DCC
R_{free} test set	2880 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	58.4	Xtrriage
Anisotropy	0.033	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11391	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, NMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1963	0.49	0/2666
1	B	0.31	0/1892	0.50	0/2566
1	C	0.30	0/1898	0.48	0/2574
1	D	0.30	0/1658	0.52	1/2260 (0.0%)
1	E	0.34	0/1953	0.52	2/2655 (0.1%)
1	F	0.33	0/1928	0.48	0/2613
All	All	0.32	0/11292	0.50	3/15334 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	165	LEU	CA-CB-CG	5.98	129.06	115.30
1	D	231	ASP	CB-CG-OD2	5.04	122.83	118.30
1	E	33	LEU	CA-CB-CG	5.04	126.88	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1934	0	1855	25	0
1	B	1863	0	1819	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1871	0	1821	21	0
1	D	1631	0	1386	17	0
1	E	1923	0	1811	25	0
1	F	1899	0	1847	27	0
2	A	22	0	14	0	0
2	B	22	0	14	1	0
2	C	22	0	14	0	0
2	D	22	0	14	0	0
2	E	22	0	14	1	0
2	F	22	0	14	0	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	19	0	0	1	0
4	B	51	0	0	5	0
4	C	15	0	0	2	0
4	D	14	0	0	0	0
4	E	17	0	0	3	0
4	F	20	0	0	2	0
All	All	11391	0	10623	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:200:GLU:HG2	1:C:213:VAL:HG23	1.61	0.83
1:A:229:ILE:HG22	1:F:233:LEU:HD11	1.61	0.81
1:F:200:GLU:HG2	1:F:213:VAL:HG23	1.64	0.79
1:D:58:THR:OG1	1:D:61:GLU:HB2	1.87	0.75
1:F:55:GLU:OE2	1:F:190:HIS:ND1	2.21	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/286 (88%)	249 (98%)	4 (2%)	0	100	100
1	B	239/286 (84%)	234 (98%)	5 (2%)	0	100	100
1	C	243/286 (85%)	240 (99%)	3 (1%)	0	100	100
1	D	228/286 (80%)	223 (98%)	5 (2%)	0	100	100
1	E	256/286 (90%)	248 (97%)	8 (3%)	0	100	100
1	F	245/286 (86%)	242 (99%)	3 (1%)	0	100	100
All	All	1464/1716 (85%)	1436 (98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/236 (82%)	193 (100%)	0	100	100
1	B	193/236 (82%)	190 (98%)	3 (2%)	62	81
1	C	193/236 (82%)	188 (97%)	5 (3%)	46	70
1	D	135/236 (57%)	135 (100%)	0	100	100
1	E	188/236 (80%)	184 (98%)	4 (2%)	53	76
1	F	195/236 (83%)	194 (100%)	1 (0%)	88	95
All	All	1097/1416 (78%)	1084 (99%)	13 (1%)	71	86

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	175	SER
1	E	33	LEU
1	F	103	MET
1	E	106	SER
1	E	205	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NMN	A	501	-	22,23,23	3.94	11 (50%)	30,34,34	1.21	4 (13%)
2	NMN	F	501	-	22,23,23	3.96	11 (50%)	30,34,34	1.23	5 (16%)
2	NMN	E	501	-	22,23,23	3.98	11 (50%)	30,34,34	1.16	3 (10%)
2	NMN	B	501	-	22,23,23	3.97	11 (50%)	30,34,34	1.13	3 (10%)
2	NMN	C	501	-	22,23,23	3.96	11 (50%)	30,34,34	1.31	3 (10%)
2	NMN	D	501	-	22,23,23	3.96	11 (50%)	30,34,34	1.23	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NMN	A	501	-	-	7/14/30/30	0/2/2/2
2	NMN	F	501	-	-	5/14/30/30	0/2/2/2
2	NMN	E	501	-	-	7/14/30/30	0/2/2/2
2	NMN	B	501	-	-	7/14/30/30	0/2/2/2
2	NMN	C	501	-	-	2/14/30/30	0/2/2/2
2	NMN	D	501	-	-	7/14/30/30	0/2/2/2

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	501	NMN	C2R-C3R	-10.71	1.24	1.53
2	B	501	NMN	C2R-C3R	-10.63	1.24	1.53
2	F	501	NMN	C2R-C3R	-10.50	1.24	1.53
2	D	501	NMN	C2R-C3R	-10.46	1.24	1.53
2	C	501	NMN	C2R-C3R	-10.45	1.24	1.53

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	NMN	C3R-C2R-C1R	4.42	107.63	100.98
2	D	501	NMN	C3R-C2R-C1R	4.06	107.09	100.98
2	A	501	NMN	C3R-C2R-C1R	3.27	105.91	100.98
2	E	501	NMN	C3R-C2R-C1R	3.27	105.90	100.98
2	F	501	NMN	C3R-C2R-C1R	2.93	105.39	100.98

There are no chirality outliers.

5 of 35 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	NMN	C5R-O5R-P-O3P
2	A	501	NMN	C5R-O5R-P-O1P
2	A	501	NMN	C5R-O5R-P-O2P
2	A	501	NMN	O4R-C4R-C5R-O5R
2	A	501	NMN	C3R-C4R-C5R-O5R


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	501	NMN	1	0
2	B	501	NMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/286 (89%)	0.20	6 (2%) 60 55	40, 63, 81, 96	0
1	B	243/286 (84%)	0.13	4 (1%) 72 68	34, 47, 67, 82	0
1	C	247/286 (86%)	0.17	4 (1%) 72 68	37, 58, 80, 102	0
1	D	232/286 (81%)	0.58	26 (11%) 5 3	39, 79, 107, 113	0
1	E	260/286 (90%)	0.47	18 (6%) 16 12	38, 75, 94, 102	0
1	F	249/286 (87%)	0.26	9 (3%) 42 36	37, 63, 87, 109	0
All	All	1488/1716 (86%)	0.30	67 (4%) 33 27	34, 61, 96, 113	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	138	ALA	7.1
1	D	119	ALA	5.8
1	A	264	ASN	4.4
1	D	105	TYR	4.3
1	D	158	VAL	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	D	502	1/1	0.63	0.13	50,50,50,50	1
3	MN	A	502	1/1	0.83	0.10	47,47,47,47	1
2	NMN	F	501	22/22	0.84	0.24	61,99,105,107	0
2	NMN	C	501	22/22	0.89	0.26	60,98,108,108	0
2	NMN	B	501	22/22	0.89	0.26	49,71,78,79	0
2	NMN	E	501	22/22	0.90	0.22	54,96,118,126	0
2	NMN	A	501	22/22	0.92	0.21	56,92,100,101	0
2	NMN	D	501	22/22	0.92	0.22	54,93,99,100	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NMN C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NMN B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NMN E 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NMN A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.