



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 14, 2026 – 10:16 AM JST

PDB ID : 9UJ5 / pdb\_00009uj5  
Title : Crystal structure of SME-1 E166A in complex with cefprozil  
Authors : Dhankhar, K.; Hazra, S.  
Deposited on : 2025-04-17  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
1	B	275	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	B	303	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4329 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase SME-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	1	0
			2064	1291	368	398	7			
1	B	267	Total	C	N	O	S	0	1	0
			2064	1291	368	398	7			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP P52682
A	0	GLY	-	expression tag	UNP P52682
A	143	ALA	GLU	engineered mutation	UNP P52682
A	268	HIS	-	expression tag	UNP P52682
A	269	HIS	-	expression tag	UNP P52682
A	270	HIS	-	expression tag	UNP P52682
A	271	HIS	-	expression tag	UNP P52682
A	272	HIS	-	expression tag	UNP P52682
A	273	HIS	-	expression tag	UNP P52682
B	-1	MET	-	initiating methionine	UNP P52682
B	0	GLY	-	expression tag	UNP P52682
B	143	ALA	GLU	engineered mutation	UNP P52682
B	268	HIS	-	expression tag	UNP P52682
B	269	HIS	-	expression tag	UNP P52682
B	270	HIS	-	expression tag	UNP P52682
B	271	HIS	-	expression tag	UNP P52682
B	272	HIS	-	expression tag	UNP P52682
B	273	HIS	-	expression tag	UNP P52682

- Molecule 2 is Acylated Cefprozil (CCD ID: A1L87) (formula:  $C_{18}H_{21}N_3O_5S$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	18	3	5	1		
2	B	1	Total	C	N	O	S	0	0
			27	18	3	5	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Cl	0	0
			4	4		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	62	Total	O	0	0
			62	62		
5	B	76	Total	O	0	0
			76	76		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase SME-1



- Molecule 1: Beta-lactamase SME-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	36.07Å 51.24Å 131.15Å 90.00° 93.88° 90.00°	Depositor
Resolution (Å)	21.32 – 2.30 21.32 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (21.32-2.30) 99.8 (21.32-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.63 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.177 , 0.267 0.182 , 0.265	Depositor DCC
$R_{free}$ test set	1058 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.6	Xtriage
Anisotropy	1.064	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 35.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4329	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, A1L87, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2099	1.26	10/2830 (0.4%)
1	B	0.59	0/2099	1.29	14/2830 (0.5%)
All	All	0.58	0/4198	1.28	24/5660 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123	GLU	CB-CG-CD	8.06	126.30	112.60
1	B	174	ASN	CB-CA-C	-7.90	95.91	109.65
1	B	41	ARG	CG-CD-NE	-6.26	98.23	112.00
1	A	17	PHE	CA-CB-CG	6.19	119.99	113.80
1	A	111	THR	CA-CB-OG1	-6.03	100.56	109.60
1	A	153	ASP	CA-CB-CG	5.98	118.58	112.60
1	A	243	ARG	CA-CB-CG	-5.94	102.22	114.10
1	B	197	ARG	CB-CA-C	5.78	121.92	110.42
1	B	243	ARG	CD-NE-CZ	5.75	132.45	124.40
1	B	144	LEU	N-CA-CB	5.61	118.56	110.20
1	B	20	ARG	CD-NE-CZ	5.56	132.18	124.40
1	A	4	ASP	CA-CB-CG	5.53	118.13	112.60
1	A	116	GLU	CB-CG-CD	5.50	121.95	112.60
1	A	243	ARG	CD-NE-CZ	5.45	132.02	124.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	243	ARG	CB-CA-C	5.40	120.14	109.66
1	A	95	LEU	CA-C-N	5.31	125.83	119.94
1	A	95	LEU	C-N-CA	5.31	125.83	119.94
1	A	192	THR	CA-CB-OG1	-5.27	101.69	109.60
1	B	60	VAL	N-CA-CB	5.26	116.70	110.55
1	B	203	PRO	CA-C-N	5.20	127.50	120.38
1	B	203	PRO	C-N-CA	5.20	127.50	120.38
1	B	243	ARG	CA-CB-CG	-5.18	103.74	114.10
1	B	66	ASP	CA-CB-CG	5.10	117.70	112.60
1	B	79	GLU	CB-CG-CD	5.07	121.22	112.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	243	ARG	Sidechain
1	A	260[A]	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2064	0	2055	20	0
1	B	2064	0	2055	28	0
2	A	27	0	0	0	0
2	B	27	0	0	2	0
3	A	4	0	0	2	0
3	B	1	0	0	2	0
4	B	4	0	6	0	0
5	A	62	0	0	1	0
5	B	76	0	0	2	0
All	All	4329	0	4116	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ASN:O	1:A:116:GLU:HG2	1.67	0.94
1:B:112:ASN:O	1:B:116:GLU:HG2	1.71	0.91
1:B:145:GLU:HG2	5:B:461:HOH:O	1.75	0.85
1:A:66:ASP:H	1:A:69:GLN:HE21	1.34	0.75
1:A:82:SER:O	1:A:86:THR:HG23	1.87	0.73
1:A:20:ARG:NH2	1:A:41:ARG:O	2.22	0.73
1:A:17:PHE:CD2	1:A:243:ARG:HD2	2.25	0.72
1:B:66:ASP:H	1:B:69:GLN:HE21	1.40	0.70
1:B:74:GLU:H	1:B:74:GLU:CD	2.00	0.69
1:A:63:LYS:NZ	5:A:401:HOH:O	2.23	0.65
1:A:76:ARG:HD3	1:A:78:LEU:HD23	1.79	0.65
1:B:226:VAL:HG22	1:B:236:ILE:HD13	1.79	0.65
1:B:79:GLU:HB3	1:B:144:LEU:HD12	1.79	0.64
1:B:17:PHE:CD2	1:B:243:ARG:HD2	2.33	0.62
1:A:139:LEU:HD11	1:A:146:LEU:HD22	1.84	0.59
1:B:123:GLU:H	1:B:123:GLU:CD	2.15	0.55
1:B:147:ASN:ND2	3:B:303:CL:CL	2.73	0.54
1:B:226:VAL:HG22	1:B:236:ILE:CD1	2.38	0.53
1:B:143:ALA:O	3:B:303:CL:CL	2.66	0.51
1:A:123:GLU:HG3	1:A:124:GLY:H	1.77	0.50
1:A:143:ALA:O	3:A:305:CL:CL	2.67	0.50
1:B:199:ARG:HG3	5:B:423:HOH:O	2.10	0.49
1:A:123:GLU:HG3	1:A:124:GLY:N	2.28	0.49
2:B:301:A1L87:C3	2:B:301:A1L87:O4	2.61	0.48
1:B:84:ILE:HG23	1:B:88:TYR:HE2	1.80	0.47
1:A:62:GLN:HG2	1:B:28:THR:O	2.15	0.46
1:A:243:ARG:HE	1:A:248:ASP:HB3	1.80	0.46
1:A:63:LYS:HB3	1:A:63:LYS:HE2	1.76	0.46
1:A:170:LEU:O	3:A:302:CL:CL	2.71	0.45
1:B:202:VAL:HG11	1:B:227:ILE:HD13	1.98	0.45
1:B:103:LEU:C	1:B:103:LEU:HD23	2.41	0.45
1:B:17:PHE:CE2	1:B:243:ARG:CD	3.00	0.44
1:A:17:PHE:CE2	1:A:243:ARG:HD2	2.52	0.44
1:B:249:LYS:HD2	1:B:249:LYS:HA	1.88	0.44
1:B:227:ILE:O	1:B:234:PRO:HA	2.18	0.43
1:A:244:LYS:HD3	1:A:244:LYS:HA	1.81	0.42
1:A:66:ASP:O	1:A:69:GLN:HG2	2.18	0.42
1:A:120:GLY:O	1:A:123:GLU:HG3	2.19	0.42
1:B:150:ILE:O	1:B:151:PRO:C	2.61	0.42
1:B:52:LEU:HD13	1:B:111:THR:HG23	2.01	0.42
1:B:17:PHE:CE2	1:B:243:ARG:HD2	2.54	0.42
1:B:66:ASP:H	1:B:69:GLN:NE2	2.12	0.42

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:146:LEU:C	1:B:146:LEU:HD12	2.44	0.42
1:B:218:TYR:CD1	1:B:246:LYS:HA	2.55	0.42
1:B:82:SER:HB3	1:B:85:THR:OG1	2.20	0.41
1:B:215:CYS:O	2:B:301:A1L87:N3	2.52	0.41
1:A:186:ASN:N	1:A:186:ASN:HD22	2.18	0.41
1:B:201:SER:HB2	1:B:260[B]:ARG:HG2	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/275 (97%)	258 (97%)	8 (3%)	0	100	100
1	B	266/275 (97%)	262 (98%)	4 (2%)	0	100	100
All	All	532/550 (97%)	520 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/223 (96%)	203 (94%)	12 (6%)	19	28
1	B	215/223 (96%)	202 (94%)	13 (6%)	17	25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	430/446 (96%)	405 (94%)	25 (6%)	18	26

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LYS
1	A	17	PHE
1	A	70	LYS
1	A	74	GLU
1	A	87	LYS
1	A	119	LEU
1	A	123	GLU
1	A	168	ASN
1	A	210	ASP
1	A	214	SER
1	A	244	LYS
1	A	253	LYS
1	B	11	LYS
1	B	17	PHE
1	B	20	ARG
1	B	70	LYS
1	B	74	GLU
1	B	87	LYS
1	B	106	SER
1	B	119	LEU
1	B	123	GLU
1	B	144	LEU
1	B	174	ASN
1	B	214	SER
1	B	253	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	147	ASN
1	A	168	ASN
1	A	186	ASN
1	B	69	GLN
1	B	81	HIS
1	B	168	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	A1L87	A	301	1	22,28,28	1.00	2 (9%)	21,38,38	2.06	3 (14%)
4	EDO	B	302	-	3,3,3	0.10	0	2,2,2	0.01	0
2	A1L87	B	301	1	22,28,28	0.86	1 (4%)	21,38,38	1.62	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1L87	A	301	1	-	9/18/38/38	0/1/2/2
4	EDO	B	302	-	-	1/1/1/1	-
2	A1L87	B	301	1	-	10/18/38/38	0/1/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	A1L87	C5-N1	3.27	1.36	1.28
2	B	301	A1L87	C5-N1	2.37	1.34	1.28
2	A	301	A1L87	C5-C18	2.29	1.53	1.48

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	A1L87	C17-S1-C6	-6.67	81.39	94.47
2	A	301	A1L87	C10-C9-N2	4.86	120.70	114.78
2	B	301	A1L87	C17-S1-C6	-4.44	85.76	94.47
2	B	301	A1L87	C10-C9-N2	3.50	119.04	114.78
2	A	301	A1L87	O2-C9-C10	-3.07	116.96	120.99
2	B	301	A1L87	O4-C18-O5	2.94	130.32	123.61
2	B	301	A1L87	O5-C18-C5	-2.60	114.01	120.86

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	A1L87	C2-C3-C4-C5
2	A	301	A1L87	C2-C3-C4-C17
2	A	301	A1L87	O4-C18-C5-N1
2	A	301	A1L87	O5-C18-C5-N1
2	A	301	A1L87	O5-C18-C5-C4
2	A	301	A1L87	N3-C10-C9-N2
2	A	301	A1L87	C1-C2-C3-C4
2	B	301	A1L87	C2-C3-C4-C5
2	B	301	A1L87	C2-C3-C4-C17
2	B	301	A1L87	O4-C18-C5-N1
2	B	301	A1L87	O5-C18-C5-N1
2	B	301	A1L87	O5-C18-C5-C4
2	B	301	A1L87	C1-C2-C3-C4
2	A	301	A1L87	N3-C10-C9-O2
2	B	301	A1L87	N3-C10-C9-N2
2	B	301	A1L87	N3-C10-C11-C12
2	B	301	A1L87	N3-C10-C11-C16
2	B	301	A1L87	N3-C10-C9-O2
4	B	302	EDO	O1-C1-C2-O2
2	A	301	A1L87	C8-C7-N2-C9


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	A1L87	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	267/275 (97%)	-0.53	1 (0%) 88 89	9, 17, 29, 60	1 (0%)
1	B	267/275 (97%)	-0.50	0 100 100	7, 14, 30, 47	1 (0%)
All	All	534/550 (97%)	-0.52	1 (0%) 91 91	7, 16, 30, 60	2 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1L87	A	301	27/27	0.92	0.08	16,21,25,29	0
2	A1L87	B	301	27/27	0.93	0.08	14,18,20,23	0
3	CL	A	304	1/1	0.97	0.09	45,45,45,45	0
3	CL	A	305	1/1	0.97	0.17	55,55,55,55	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	B	302	4/4	0.97	0.05	27,27,27,28	0
3	CL	A	303	1/1	0.98	0.05	39,39,39,39	0
3	CL	B	303	1/1	0.98	0.12	44,44,44,44	0
3	CL	A	302	1/1	0.98	0.15	45,45,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around A1L87 A 301:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A1L87 B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CL A 304:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CL A 305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around EDO B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CL A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CL B 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CL A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.