

wwPDB NMR Structure Validation Summary Report (i)

Oct 23, 2023 – 05:01 PM EDT

:	8ULM
:	31111
:	Chickpea (Cicer arientinum) nodule-specific cysteine-rich peptide NCR13: So-
	lution NMR structure of the isomer with C4:C23, C15:C30, and C10:C28 disul-
	fide bonds
:	Buchko, G.W.; Zhou, M.; Shah, D.M.; Velivelli, S.L.S.
:	2023-10-16
	:

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 71%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	NMR archive $(\#Futrics)$
	(#Entries)	$(\# { m Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	А	33	67%	27%	6%		



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:2-A:32 (31)	0.54	1		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 10, 11, 17, 18, 20
2	2, 7, 8, 9, 12, 13, 14, 16
Single-model clusters	15; 19



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 533 atoms, of which 277 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Nodule cysteine-rich protein 13.

Mol	Chain	Residues	Atoms				Trace		
1	٨	9 9	Total	С	Η	Ν	Ο	S	0
	A	33	533	159	277	49	42	6	U



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Nodule cysteine-rich protein 13



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: Nodule cysteine-rich protein 13

Chain A:		58%	36%	6%
8 8 8 8 9 8 9 8 8 8 8 8 8 8 8 8 8 8 8 8	K12 F13 F13 F13 C23 C23 C23 C23 C28 C23 C28 C28 C28 C32 C32 C32 C32 C32 C32 C32 C32 C32 C32			



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: torsion angle dynamics.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: target function.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	2.1
CNS	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	321
Number of shifts mapped to atoms	321
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	71%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	А	$0.0{\pm}0.0$	0.2 ± 0.4
All	All	0	5

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	А	12	LYS	Peptide	5

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	244	263	257	6 ± 2
All	All	4880	5260	5140	123

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

5 of 53 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:15:CYS:SG	1:A:19:LYS:HD2	0.79	2.17	14	1
1:A:19:LYS:HE3	1:A:30:CYS:SG	0.73	2.24	19	1

Continued on next page...



Atom 1	Atom 2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:2:LYS:HD2	1:A:10:CYS:SG	0.71	2.25	8	1
1:A:2:LYS:NZ	1:A:10:CYS:SG	0.70	2.64	9	3
1:A:4:CYS:SG	1:A:5:GLN:N	0.67	2.67	13	20

Continued from previous page...

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Favoured Allowed		Outliers	Percentiles		
1	А	30/33~(91%)	$25 \pm 1 (84 \pm 4\%)$	3 ± 1 (12 $\pm4\%$)	2 ± 0 (5 $\pm2\%$)		4	25
All	All	600/660~(91%)	501 (84%)	69 (12%)	30~(5%)		4	25

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	А	13	PHE	20
1	А	3	PRO	9
1	А	19	LYS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Outliers	Perce	\mathbf{ntiles}
1	А	29/30~(97%)	$28 \pm 1 \ (96 \pm 3\%)$	$1 \pm 1 (4 \pm 3\%)$	34	82
All	All	580/600~(97%)	556 (96%)	24 (4%)	34	82

5 of 8 unique residues with a non-rotameric side chain are listed below. They are sorted by the frequency of occurrence in the ensemble.



Mol	Chain	Res	Type	Models (Total)
1	А	32	ARG	5
1	А	2	LYS	4
1	А	8	LYS	4
1	А	11	LYS	4
1	А	17	LYS	4

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 71% for the well-defined parts and 72% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chemical_shifts_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	321
Number of shifts mapped to atoms	321
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction}\pm{\rm precision},ppm$	Suggested action
$^{13}C_{\alpha}$	31	0.17 ± 0.53	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	31	-0.21 ± 0.64	None needed (< 0.5 ppm)
$^{13}C'$	0		None (insufficient data)
¹⁵ N	29	0.18 ± 0.75	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 71%, i.e. 304 atoms were assigned a chemical shift out of a possible 426. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	117/150~(78%)	60/60~(100%)	29/62~(47%)	28/28~(100%)
Sidechain	179/256~(70%)	124/162~(77%)	53/78~(68%)	2/16~(12%)

Continued on next page...



	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Aromatic	8/20 (40%)	8/10 (80%)	0/10~(0%)	$0/0 \ (\%)$
Overall	304/426~(71%)	192/232~(83%)	82/150~(55%)	30/44~(68%)

Continued from previous page...

7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	А	1	THR	HG1	4.15	0.08 - 2.19	14.3

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	316
Intra-residue $(i-j =0)$	98
Sequential (i-j =1)	107
Medium range ($ i-j >1$ and $ i-j <5$)	34
Long range $(i-j \ge 5)$	54
Inter-chain	0
Hydrogen bond restraints	20
Disulfide bond restraints	3
Total dihedral-angle restraints	48
Number of unmapped restraints	0
Number of restraints per residue	11.0
Number of long range restraints per residue ¹	2.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	5.4	0.2
0.2-0.5 (Medium)	5.7	0.49
>0.5 (Large)	6.0	1.67



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max $(^{\circ})$
1.0-10.0 (Small)	8.2	5.11
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Bestroints type	Count	Count $\%^1$		$olated^3$		Consis	tently	Violated ⁴
Restraints type	Count	70-	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^{1}$
Intra-residue (i-j =0)	98	31.0	4	4.1	1.3	2	2.0	0.6
Backbone-Backbone	2	0.6	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	86	27.2	4	4.7	1.3	2	2.3	0.6
Sidechain-Sidechain	10	3.2	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	107	33.9	5	4.7	1.6	2	1.9	0.6
Backbone-Backbone	27	8.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	72	22.8	5	6.9	1.6	2	2.8	0.6
Sidechain-Sidechain	8	2.5	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	34	10.8	4	11.8	1.3	0	0.0	0.0
Backbone-Backbone	11	3.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	16	5.1	3	18.8	0.9	0	0.0	0.0
Sidechain-Sidechain	7	2.2	1	14.3	0.3	0	0.0	0.0
Long range $(i-j \ge 5)$	54	17.1	11	20.4	3.5	1	1.9	0.3
Backbone-Backbone	12	3.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	24	7.6	10	41.7	3.2	1	4.2	0.3
Sidechain-Sidechain	18	5.7	1	5.6	0.3	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	20	6.3	0	0.0	0.0	0	0.0	0.0
Disulfide bond	3	0.9	3	100.0	0.9	3	100.0	0.9
Total	316	100.0	27	8.5	8.5	8	2.5	2.5
Backbone-Backbone	52	16.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	218	69.0	22	10.1	7.0	5	2.3	1.6
Sidechain-Sidechain	46	14.6	5	10.9	1.6	3	6.5	0.9

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID		Nun	nber o	f viola	ations	3	Maan (Å)	Max (Å)	SD^6 (Å)	Madian (Å)
Model ID	IR^{1}	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (Å)	Max (A)	$SD^{*}(A)$	Median (Å)
1	3	3	2	7	0	15	0.52	1.3	0.38	0.4
2	3	3	2	7	0	15	0.42	1.15	0.31	0.34
3	3	3	2	8	0	16	0.4	1.22	0.3	0.38
4	3	3	2	9	0	17	0.41	1.45	0.34	0.28
5	3	4	3	8	0	18	0.6	1.59	0.46	0.42
6	3	3	1	10	0	17	0.45	1.67	0.37	0.41
7	3	5	1	8	0	17	0.46	1.44	0.4	0.27
8	3	3	1	10	0	17	0.41	1.63	0.37	0.3
9	3	3	1	7	0	14	0.43	1.21	0.3	0.37
10	3	5	3	8	0	19	0.57	1.37	0.43	0.43
11	3	5	2	9	0	19	0.46	1.19	0.31	0.43

Continued on next page...




Madal ID		Nun	nber o	f viola	ations	5	Mean (Å)		SD^6 (Å)	Madian (Å)
Model ID	IR^{1}	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (A)	Max (Å)	SD^{6} (Å)	Median (Å)
12	3	3	1	7	0	14	0.55	1.47	0.41	0.5
13	3	3	2	8	0	16	0.52	1.65	0.4	0.41
14	3	5	2	8	0	18	0.44	1.67	0.37	0.42
15	3	3	3	7	0	16	0.52	1.15	0.34	0.44
16	3	5	3	9	0	20	0.5	1.48	0.36	0.41
17	3	3	2	8	0	16	0.51	1.37	0.39	0.42
18	3	5	2	7	0	17	0.49	1.39	0.39	0.37
19	3	5	1	11	0	20	0.47	1.32	0.35	0.36
20	3	5	2	10	0	20	0.52	1.49	0.41	0.4

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation





The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right




9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 269(IR:94, SQ:102, MR:30, LR:43, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restra	aints	Fractio	n of the ensemble
IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Count^6	%
1	0	0	0	0	1	1	5.0
0	0	1	1	0	2	2	10.0
0	0	0	1	0	1	3	15.0
0	0	0	1	0	1	4	20.0
0	0	0	2	0	2	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	1	0	1	7	35.0
0	0	0	0	0	0	8	40.0
0	2	1	0	0	3	9	45.0
0	0	0	0	0	0	10	50.0
0	0	1	0	0	1	11	55.0
0	0	0	2	0	2	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	1	0	0	1	16	80.0
0	0	0	1	0	1	17	85.0
0	0	0	0	0	0	18	90.0
1	1	0	1	0	3	19	95.0
2	2	0	1	0	5	20	100.0

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations






9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	20	1.38	0.18	1.36
(1,129)	1:14:A:ALA:H	1:15:A:CYS:HB3	20	0.83	0.13	0.83
(1,165)	1:23:A:CYS:H	1:24:A:ILE:HG13	20	0.63	0.18	0.58
(1,173)	1:24:A:ILE:H	1:24:A:ILE:HG13	20	0.61	0.13	0.65
(1,123)	1:27:A:PHE:H	1:27:A:PHE:HB3	20	0.44	0.02	0.44
(3,2)	1:15:A:CYS:SG	1:30:A:CYS:SG	20	0.13	0.0	0.13
(3,1)	1:4:A:CYS:SG	1:23:A:CYS:SG	20	0.13	0.0	0.13
(3,3)	1:10:A:CYS:SG	1:28:A:CYS:SG	20	0.12	0.01	0.13
(1,213)	1:19:A:LYS:HB2	1:31:A:VAL:H	19	0.43	0.13	0.39
(1,86)	1:9:A:ASP:H	1:10:A:CYS:HB3	19	0.39	0.09	0.37


 $^1\mathrm{Number}$ of violated models, $^2\mathrm{Standard}$ deviation



9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	6	1.67
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	14	1.67
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	13	1.65
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	8	1.63
(1,215)	1:19:A:LYS:HG3	1:31:A:VAL:H	5	1.59
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	20	1.49
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	16	1.48
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	12	1.47
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	4	1.45
(1,16)	1:6:A:SER:HA	1:23:A:CYS:HB3	7	1.44



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

	Count	$\%^1$	Vie	olated	3	Consis	stently	r Violated ⁴
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
PSI	24	50.0	6	25.0	12.5	2	8.3	4.2
PHI	24	50.0	9	37.5	18.8	3	12.5	6.2
Total	48	100.0	15	31.2	31.2	5	10.4	10.4

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber o	of violations	Maan (°)	M_{OV} (°)	SD (°)	Median (°)
Model ID	PSI	PHI	Total	Mean ($^{\circ}$)	$Max (^{\circ})$	$SD(^{\circ})$	Median ()
1	3	5	8	2.27	3.68	0.94	2.64
2	2	4	6	3.61	4.44	0.99	4.07
3	4	6	10	2.84	4.49	1.03	2.66
4	4	7	11	2.28	4.25	1.0	1.77
5	3	5	8	2.4	3.35	0.89	2.8
6	3	7	10	2.38	4.92	1.33	1.95
7	3	4	7	2.76	4.03	1.04	3.16
8	4	4	8	2.4	3.89	1.1	2.58
9	4	5	9	2.34	3.88	1.11	2.59
10	3	4	7	2.87	5.06	1.44	2.58
11	2	5	7	2.54	3.66	0.94	2.65
12	3	4	7	2.98	4.52	1.22	3.55
13	3	4	7	2.72	3.92	0.97	3.05
14	3	4	7	3.09	4.53	1.23	3.55
15	3	4	7	3.02	5.11	1.49	2.65
16	4	7	11	2.14	4.03	0.97	1.86
17	3	7	10	2.46	4.11	1.09	2.12
18	4	6	10	2.08	3.91	1.0	1.6
19	2	4	6	2.6	3.79	0.83	2.7
20	3	6	9	2.15	3.7	0.98	2.3





10.2.1 Bar graph : Dihedral violation statistics for each model (i)

The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fractio	n of the ensemble
PSI	PHI	Total	Count^1	%
1	1	2	1	5.0
0	1	1	2	10.0
1	0	1	3	15.0
0	0	0	4	20.0
0	0	0	5	25.0
0	2	2	6	30.0
0	0	0	7	35.0
0	1	1	8	40.0
1	0	1	9	45.0
1	0	1	10	50.0
0	0	0	11	55.0


Continued on next page...




Continuea from prettous page								
Nun	nber o	of violated restraints	Fraction of the ensemble					
PSI	PHI	Total	Count^1	%				
0	0	0	12	60.0				
0	0	0	13	65.0				
0	0	0	14	70.0				
0	0	0	15	75.0				
0	0	0	16	80.0				
0	0	0	17	85.0				
0	0	0	18	90.0				
0	1	1	19	95.0				
2	3	5	20	100.0				

Continued from previous page...

 1 Number of models with violations







10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models







10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	$Models^1$	Mean	\mathbf{SD}^2	Median
(1,16)	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	1:12:A:LYS:N	20	3.86	0.59	3.92
(1,14)	1:10:A:CYS:N	1:10:A:CYS:CA	1:10:A:CYS:C	1:11:A:LYS:N	20	3.63	0.69	3.62
(1,15)	1:10:A:CYS:C	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	20	3.25	0.85	3.03
(1,17)	1:11:A:LYS:C	1:12:A:LYS:N	1:12:A:LYS:CA	1:12:A:LYS:C	20	3.18	0.64	3.22
(1,3)	1:3:A:PRO:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	20	2.77	0.26	2.72
(1,41)	1:27:A:PHE:C	1:28:A:CYS:N	1:28:A:CYS:CA	1:28:A:CYS:C	19	1.47	0.25	1.37
(1,40)	1:27:A:PHE:N	1:27:A:PHE:CA	1:27:A:PHE:C	1:28:A:CYS:N	10	1.26	0.2	1.2
(1,20)	1:13:A:PHE:N	1:13:A:PHE:CA	1:13:A:PHE:C	1:14:A:ALA:N	9	1.26	0.16	1.22
(1,19)	1:12:A:LYS:C	1:13:A:PHE:N	1:13:A:PHE:CA	1:13:A:PHE:C	8	1.23	0.18	1.19
(1,9)	1:7:A:ASP:C	1:8:A:LYS:N	1:8:A:LYS:CA	1:8:A:LYS:C	6	1.29	0.2	1.23


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation ($^{\circ}$)
(1,14)	1:10:A:CYS:N	1:10:A:CYS:CA	1:10:A:CYS:C	1:11:A:LYS:N	15	5.11
(1,15)	1:10:A:CYS:C	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	15	5.1
(1,14)	1:10:A:CYS:N	1:10:A:CYS:CA	1:10:A:CYS:C	1:11:A:LYS:N	10	5.06
(1,16)	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	1:12:A:LYS:N	6	4.92
(1,15)	1:10:A:CYS:C	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	10	4.83
(1,16)	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	1:12:A:LYS:N	14	4.53
(1,17)	1:11:A:LYS:C	1:12:A:LYS:N	1:12:A:LYS:CA	1:12:A:LYS:C	6	4.52
(1,16)	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	1:12:A:LYS:N	12	4.52
(1,16)	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	1:12:A:LYS:N	3	4.49
(1,16)	1:11:A:LYS:N	1:11:A:LYS:CA	1:11:A:LYS:C	1:12:A:LYS:N	2	4.44

