



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2023 – 03:25 PM EST

PDB ID : 5UPQ  
Title : Acyl-CoA synthetase PtmA2 from *Streptomyces platensis* in complex with SBNP465 ligand  
Authors : Osipiuk, J.; Hatzos-Skintges, C.; Endres, M.; Babnigg, G.; Rudolf, J.D.; Chang, C.Y.; Ma, M.; Shen, B.; Phillips Jr., G.N.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG); Enzyme Discovery for Natural Product Biosynthesis (NatPro)  
Deposited on : 2017-02-03  
Resolution : 2.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	525	88% 10% .
1	B	525	83% 10% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	702	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA synthetase PtmA2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	512	3924	2451	727	729	7	10	0	0	0
1	B	488	3746	2342	693	695	6	10	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A0A0A0V031
A	-1	ASN	-	expression tag	UNP A0A0A0V031
A	0	ALA	-	expression tag	UNP A0A0A0V031
A	1	LEU	-	expression tag	UNP A0A0A0V031
A	141	PRO	ALA	conflict	UNP A0A0A0V031
A	246	ASP	GLY	conflict	UNP A0A0A0V031
B	-2	SER	-	expression tag	UNP A0A0A0V031
B	-1	ASN	-	expression tag	UNP A0A0A0V031
B	0	ALA	-	expression tag	UNP A0A0A0V031
B	1	LEU	-	expression tag	UNP A0A0A0V031
B	141	PRO	ALA	conflict	UNP A0A0A0V031
B	246	ASP	GLY	conflict	UNP A0A0A0V031

- Molecule 2 is 5'-O-[(R)-{[(7beta,8alpha,9beta,10alpha,13alpha,16beta)-7,16-dihydroxy-18-oxokauran-18-yl]oxy}(hydroxy)phosphoryl]adenosine (three-letter code: 8JJ) (formula: C<sub>30</sub>H<sub>44</sub>N<sub>5</sub>O<sub>10</sub>P).



*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0
4	B	1	Total Cl 1 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	119	Total O 119 119	0	1
6	B	93	Total O 93 93	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acyl-CoA synthetase PtmA2



- Molecule 1: Acyl-CoA synthetase PtmA2





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.61Å 145.74Å 145.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.50 – 2.42 38.43 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.4 (38.50-2.42) 99.4 (38.43-2.41)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.42 (at 2.42Å)	Xtrriage
Refinement program	REFMAC 5.8.0069	Depositor
R, $R_{free}$	0.232 , 0.285 0.233 , 0.284	Depositor DCC
$R_{free}$ test set	2820 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.0	Xtrriage
Anisotropy	0.241	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 20.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	0.346 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8048	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, 8JJ, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/4010	0.68	0/5444
1	B	0.46	0/3827	0.68	0/5187
All	All	0.47	0/7837	0.68	0/10631

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3924	0	3807	29	0
1	B	3746	0	3635	31	0
2	A	46	0	0	0	0
2	B	46	0	0	1	0
3	A	30	0	0	0	0
3	B	35	0	0	3	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
5	B	6	0	8	0	0
6	A	119	0	0	3	0
6	B	93	0	0	1	0
All	All	8048	0	7450	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:LEU:HD21	1:A:372:TRP:HB2	1.79	0.63
1:B:321:VAL:HG13	1:B:335:ARG:C	2.19	0.63
1:A:410:MSE:HE3	1:A:417:ASN:HD22	1.65	0.61
1:A:1:LEU:HD21	1:B:385:GLY:HA2	1.86	0.58
1:A:1:LEU:HD13	1:B:382:PHE:CD1	2.40	0.56
1:A:271:ARG:NH1	1:A:290:GLU:OE1	2.39	0.56
1:B:340:LEU:HD11	1:B:362:ALA:HB1	1.88	0.54
1:A:340:LEU:HD11	1:A:362:ALA:HB1	1.89	0.54
1:B:310:GLY:O	2:B:701:8JJ:N4	2.42	0.52
1:B:504:ASP:O	1:B:509:GLY:N	2.42	0.52
1:A:395:ARG:NH1	1:B:12:GLU:OE2	2.44	0.50
1:B:21:THR:OG1	3:B:702:SO4:O3	2.27	0.50
1:B:475:ILE:HB	1:B:479:LYS:HE3	1.93	0.50
1:A:259:MSE:HB2	6:A:913:HOH:O	2.12	0.49
1:A:212:PRO:HB2	1:A:214:PHE:CE2	2.49	0.48
1:B:111:GLU:HB3	1:B:112:ILE:HG23	1.96	0.48
1:B:29:ARG:NH1	3:B:702:SO4:O4	2.47	0.47
1:A:258:LEU:C	1:A:259:MSE:HE2	2.34	0.47
1:A:410:MSE:CE	1:A:417:ASN:HD22	2.27	0.47
1:A:503:LEU:O	1:A:504:ASP:C	2.53	0.46
1:A:309:ARG:NH1	1:A:331:GLY:O	2.50	0.45
1:B:409:ARG:HH12	1:B:439:ILE:HG22	1.81	0.45
1:B:409:ARG:NH1	1:B:439:ILE:HG22	2.32	0.45
1:A:320:ALA:HB1	1:A:340:LEU:HD22	1.98	0.45
1:B:476:ALA:H	1:B:479:LYS:HE2	1.82	0.45
1:B:20:ARG:NH2	6:B:806:HOH:O	2.49	0.45
1:A:474:ARG:O	1:A:474:ARG:HG2	2.17	0.44
1:A:432:ALA:HB1	1:A:460:ALA:HB1	1.98	0.44
1:A:281:ARG:HA	1:A:294:THR:O	2.18	0.44
1:B:284:VAL:O	1:B:285:ALA:C	2.56	0.44
1:B:85:TYR:CZ	1:B:87:ARG:HB2	2.53	0.44
1:B:241:ALA:HB2	1:B:265:GLU:OE1	2.17	0.44
1:A:42:LEU:HD21	1:A:70:LEU:HD23	1.99	0.44
1:A:270:ASN:OD1	1:A:275:HIS:HB2	2.18	0.44
1:B:381:ARG:NE	1:B:388:HIS:O	2.45	0.44
1:A:410:MSE:HE3	1:A:417:ASN:ND2	2.32	0.43
1:B:267:VAL:HG21	1:B:288:LEU:O	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:VAL:HG21	1:A:289:TRP:HA	2.00	0.43
1:A:206:ALA:HB3	1:A:253:CYS:HA	2.00	0.43
1:B:192:ALA:HB1	1:B:225:PRO:HA	2.01	0.42
1:A:22:ALA:HB2	1:A:32:TRP:CD2	2.54	0.42
1:B:345:LEU:HD23	1:B:351:GLU:HA	2.02	0.42
1:A:180:GLY:O	1:A:371:TYR:HA	2.20	0.42
1:B:53:ARG:HG3	3:B:703:SO4:O1	2.20	0.42
1:A:23:LEU:HD13	1:A:65:PHE:HB2	2.02	0.42
1:B:491:PRO:CD	1:B:503:LEU:HD23	2.50	0.42
1:B:158:LEU:HD23	1:B:158:LEU:HA	1.93	0.41
1:B:56:ARG:HB3	1:B:100:PHE:CZ	2.56	0.41
1:A:495:SER:HB3	6:A:896:HOH:O	2.20	0.41
1:A:126:ASP:HA	6:A:851:HOH:O	2.19	0.41
1:A:311:TYR:CG	1:A:312:GLY:N	2.89	0.41
1:B:252:GLU:OE2	1:B:279[B]:ARG:NE	2.54	0.41
1:B:213:MSE:HE1	1:B:235:ILE:HD12	2.03	0.40
1:A:410:MSE:HE3	1:A:417:ASN:CB	2.51	0.40
1:B:335:ARG:NH2	1:B:398:ASP:O	2.51	0.40
1:B:63:ASN:HB3	1:B:213:MSE:HE3	2.02	0.40
1:B:85:TYR:HB3	1:B:88:TRP:CE2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/525 (97%)	489 (96%)	18 (4%)	1 (0%)	47	61
1	B	479/525 (91%)	449 (94%)	30 (6%)	0	100	100
All	All	987/1050 (94%)	938 (95%)	48 (5%)	1 (0%)	51	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	416	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	397/393 (101%)	388 (98%)	9 (2%)	50	68
1	B	379/393 (96%)	372 (98%)	7 (2%)	59	75
All	All	776/786 (99%)	760 (98%)	16 (2%)	53	71

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	279	ARG
1	A	319	PHE
1	A	368	HIS
1	A	426	CYS
1	A	444	THR
1	A	471	CYS
1	A	480	LYS
1	A	483	SER
1	B	2	GLN
1	B	134	HIS
1	B	319	PHE
1	B	368	HIS
1	B	409	ARG
1	B	459	ASP
1	B	480	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	417	ASN
1	B	2	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	388	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 3 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	704	-	4,4,4	0.41	0	6,6,6	0.24	0
2	8JJ	A	701	-	48,52,52	1.14	4 (8%)	67,86,86	1.53	8 (11%)
3	SO4	B	705	-	4,4,4	0.41	0	6,6,6	0.17	0
3	SO4	A	703	-	4,4,4	0.31	0	6,6,6	0.39	0
3	SO4	B	706	-	4,4,4	0.38	0	6,6,6	0.19	0
2	8JJ	B	701	-	48,52,52	1.08	5 (10%)	67,86,86	1.59	14 (20%)
3	SO4	B	702	-	4,4,4	0.40	0	6,6,6	0.22	0
3	SO4	B	707	-	4,4,4	0.35	0	6,6,6	0.26	0
3	SO4	B	708	-	4,4,4	0.37	0	6,6,6	0.35	0
3	SO4	B	704	-	4,4,4	0.36	0	6,6,6	0.27	0
3	SO4	A	707	-	4,4,4	0.53	0	6,6,6	0.36	0
3	SO4	B	703	-	4,4,4	0.34	0	6,6,6	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	706	-	4,4,4	0.33	0	6,6,6	0.40	0
3	SO4	A	705	-	4,4,4	0.39	0	6,6,6	0.23	0
3	SO4	A	702	-	4,4,4	0.36	0	6,6,6	0.22	0
5	GOL	B	709	-	5,5,5	0.51	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8JJ	B	701	-	-	4/15/101/101	0/8/7/7
2	8JJ	A	701	-	-	0/15/101/101	0/8/7/7
5	GOL	B	709	-	-	4/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	8JJ	P-O7	2.92	1.61	1.50
2	B	701	8JJ	C20-N3	2.81	1.36	1.32
2	A	701	8JJ	C5-C1	-2.69	1.52	1.55
2	A	701	8JJ	C18-C21	2.57	1.47	1.40
2	B	701	8JJ	C18-C21	2.40	1.47	1.40
2	B	701	8JJ	C1-C2	-2.33	1.51	1.54
2	B	701	8JJ	P-O7	2.23	1.58	1.50
2	A	701	8JJ	P-O2	2.13	1.65	1.60
2	B	701	8JJ	C25-C8	-2.02	1.53	1.56

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	8JJ	O2-C11-C9	6.34	121.36	111.36
2	B	701	8JJ	O2-C11-C9	4.65	118.70	111.36
2	A	701	8JJ	O9-C11-C9	-4.36	119.11	125.21
2	A	701	8JJ	N3-C20-N2	-3.79	122.76	128.68
2	B	701	8JJ	C5-C4-C27	3.67	112.80	108.87
2	B	701	8JJ	N3-C20-N2	-3.61	123.03	128.68
2	B	701	8JJ	C21-C18-N1	-3.31	105.95	109.40
2	B	701	8JJ	C3-C2-C1	3.10	103.91	101.34
2	A	701	8JJ	C1-C5-C4	-2.83	105.20	107.34
2	B	701	8JJ	O9-C11-C9	-2.83	121.25	125.21
2	B	701	8JJ	C29-C2-C1	-2.76	112.07	114.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	8JJ	O8-P-O2	2.61	112.65	104.14
2	B	701	8JJ	C10-C9-C22	-2.48	105.51	109.43
2	A	701	8JJ	C5-C1-C2	2.29	105.18	103.58
2	B	701	8JJ	O-C1-C5	2.25	113.53	108.97
2	B	701	8JJ	C3-C4-C27	-2.24	108.82	111.90
2	B	701	8JJ	O2-P-O3	2.22	109.40	102.92
2	B	701	8JJ	O3-C12-C13	2.21	116.60	108.99
2	A	701	8JJ	C21-C18-N1	-2.20	107.11	109.40
2	B	701	8JJ	C29-C28-C27	2.11	117.20	112.43
2	A	701	8JJ	C3-C2-C1	2.10	103.08	101.34
2	A	701	8JJ	O8-P-O2	2.00	110.68	104.14

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	701	8JJ	O3-C12-C13-C14
2	B	701	8JJ	C12-O3-P-O7
2	B	701	8JJ	O3-C12-C13-O4
2	B	701	8JJ	C12-O3-P-O2
5	B	709	GOL	O1-C1-C2-C3
5	B	709	GOL	O1-C1-C2-O2
5	B	709	GOL	O2-C2-C3-O3
5	B	709	GOL	C1-C2-C3-O3

There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	701	8JJ	1	0
3	B	702	SO4	2	0
3	B	703	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 8JJ A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 8JJ B 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.