



wwPDB X-ray Structure Validation Summary Report

Apr 28, 2024 – 07:09 am BST

PDB ID : 2UVP
Title : Crystal structure of HobA (HP1230) from Helicobacter pylori
Authors : Terradot, L.; Natrajan, G.; Thompson, A.C.; Hall, D.R.
Deposited on : 2007-03-13
Resolution : 1.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	 81% 13% . .
2	B	186	 75% 18% . . .
2	C	186	 75% 20% . .
2	D	186	 74% 18% . 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	1181	-	-	X	-
3	ACT	A	1182	-	-	X	-
3	ACT	C	1183	-	-	X	-
5	GOL	B	1182	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6928 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HOBA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1502	983	239	277	3	0	2	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	ILE	VAL	conflict	UNP O25828

- Molecule 2 is a protein called HOBA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	181	1525	997	243	280	5	0	4	0
2	C	186	1545	1008	249	284	4	0	2	0
2	D	176	1475	970	236	266	3	0	2	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Ca 2 2	0	0
4	C	3	Total Ca 3 3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	191	Total O 191 191	0	0
6	B	232	Total O 232 232	0	0
6	C	235	Total O 235 235	0	0
6	D	194	Total O 194 194	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HOBA




- Molecule 2: HOBA




- Molecule 2: HOBA



- Molecule 2: HOBA





4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	112.23Å 112.23Å 61.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	97.13 – 1.70 38.23 – 1.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (97.13-1.70) 99.9 (38.23-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.3.0011	Depositor
R, R_{free}	0.180 , 0.214 0.180 , 0.210	Depositor DCC
R_{free} test set	4808 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l 0.487 for h,-h-k,-l 0.022 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6928	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1539	0.68	1/2083 (0.0%)
2	B	0.61	0/1562	0.78	3/2112 (0.1%)
2	C	0.61	0/1581	0.85	3/2137 (0.1%)
2	D	0.58	0/1512	0.67	0/2045
All	All	0.59	0/6194	0.75	7/8377 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	1	0
2	C	0	1
All	All	1	1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	58	ARG	NE-CZ-NH2	-11.53	114.53	120.30
2	C	58	ARG	NE-CZ-NH1	9.31	124.96	120.30
2	B	146	LEU	CA-CB-CG	7.21	131.88	115.30
2	B	62	LEU	CB-CG-CD2	5.76	120.80	111.00
2	C	146	LEU	CA-CB-CG	5.67	128.34	115.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	97	ASN	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	93	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1502	0	1494	27	0
2	B	1525	0	1519	54	0
2	C	1545	0	1543	30	0
2	D	1475	0	1477	28	0
3	A	8	0	6	6	0
3	C	4	0	3	4	0
4	B	2	0	0	0	0
4	C	3	0	0	0	0
5	B	6	0	8	4	0
5	D	6	0	8	1	0
6	A	191	0	0	7	0
6	B	232	0	0	11	0
6	C	235	0	0	10	0
6	D	194	0	0	4	0
All	All	6928	0	6058	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:146:LEU:CD1	5:B:1182:GOL:H31	1.76	1.14
2:B:151:HIS:HD2	6:B:2205:HOH:O	1.45	1.00
1:A:107:TYR:OH	3:A:1182:ACT:H1	1.63	0.96
2:B:110:PRO:HG2	2:B:131:ILE:HD11	1.45	0.96
2:C:177:LYS:HD2	6:C:2231:HOH:O	1.64	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/186 (97%)	175 (97%)	4 (2%)	1 (1%)	25	11
2	B	183/186 (98%)	176 (96%)	4 (2%)	3 (2%)	9	1
2	C	186/186 (100%)	179 (96%)	4 (2%)	3 (2%)	9	1
2	D	174/186 (94%)	171 (98%)	2 (1%)	1 (1%)	25	11
All	All	723/744 (97%)	701 (97%)	14 (2%)	8 (1%)	14	3

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	94	ASN
2	B	95	ASP
2	C	95	ASP
2	C	0	SER
2	D	142	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/171 (98%)	158 (95%)	9 (5%)	22	7
2	B	170/171 (99%)	156 (92%)	14 (8%)	11	2
2	C	172/171 (101%)	155 (90%)	17 (10%)	8	1
2	D	165/171 (96%)	152 (92%)	13 (8%)	12	2
All	All	674/684 (98%)	621 (92%)	53 (8%)	13	2

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	58	ARG
2	C	127	LEU
2	D	131	ILE
2	C	62	LEU
2	C	86	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	114	HIS
2	D	108	GLN
2	C	43	HIS
2	D	39	GLN
2	B	151	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	A	1181	-	3,3,3	0.95	0	3,3,3	0.80	0
3	ACT	A	1182	-	3,3,3	0.70	0	3,3,3	1.15	0
5	GOL	D	1181	-	5,5,5	0.29	0	5,5,5	0.96	0
5	GOL	B	1182	-	5,5,5	0.29	0	5,5,5	0.81	0
3	ACT	C	1183	-	3,3,3	0.92	0	3,3,3	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	1182	-	-	2/4/4/4	-
5	GOL	D	1181	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	1181	GOL	C1-C2-C3-O3
5	D	1181	GOL	O2-C2-C3-O3
5	B	1182	GOL	O2-C2-C3-O3
5	B	1182	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1181	ACT	2	0
3	A	1182	ACT	4	0
5	D	1181	GOL	1	0
5	B	1182	GOL	4	0
3	C	1183	ACT	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/186 (96%)	-0.40	1 (0%) 89 91	9, 19, 35, 44	2 (1%)
2	B	181/186 (97%)	-0.39	2 (1%) 80 83	8, 16, 33, 46	0
2	C	186/186 (100%)	-0.35	2 (1%) 80 83	9, 16, 40, 57	0
2	D	176/186 (94%)	-0.40	0 100 100	10, 19, 35, 40	1 (0%)
All	All	723/744 (97%)	-0.39	5 (0%) 87 90	8, 17, 35, 57	3 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	0	SER	3.9
2	C	31	SER	3.2
2	B	31	SER	2.7
2	C	0	SER	2.6
1	A	93	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ACT	A	1181	4/4	0.57	0.18	29,30,31,32	0
5	GOL	B	1182	6/6	0.80	0.12	34,34,36,37	0
5	GOL	D	1181	6/6	0.84	0.16	23,30,33,35	0
3	ACT	A	1182	4/4	0.90	0.09	11,14,16,16	0
3	ACT	C	1183	4/4	0.94	0.08	7,11,14,14	0
4	CA	B	1183	1/1	0.99	0.04	25,25,25,25	0
4	CA	C	1184	1/1	0.99	0.04	25,25,25,25	0
4	CA	B	1181	1/1	1.00	0.04	16,16,16,16	0
4	CA	C	1181	1/1	1.00	0.04	11,11,11,11	1
4	CA	C	1182	1/1	1.00	0.04	15,15,15,15	1

6.5 Other polymers [i](#)

There are no such residues in this entry.