

Full wwPDB X-ray Structure Validation Report (i)

Nov 6, 2023 – 06:26 AM EST

PDB ID	:	6V1O
Title	:	Structure of OXA-48 bound to $QPX7728$ at 1.80 A
Authors	:	Pemberton, O.A.; Chen, Y.
Deposited on		
Resolution	:	1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	:::::::::::::::::::::::::::::::::::::::	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	5950(1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	263	86%	6%	8%
1	В	263	% • 88%	•	8%
1	С	263	3% 90%	•	7%
1	D	263	% 86%	6%	8%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 9528 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	243	Total	С	Ν	0	\mathbf{S}	0	1	0
	A	240	1993	1267	352	367	7	0		0
1	В	242	Total	С	Ν	0	S	0	0	0
	D	242	1978	1259	349	363	$\overline{7}$			
1	С	245	Total	С	Ν	0	S	0	2	0
		240	2026	1291	357	371	7	0	2	0
1	П	242	Total	С	Ν	0	S	0	2	0
			1994	1268	353	366	7		2	0

• Molecule 1 is a protein called OXA family beta-lactamase.

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	3	MET	-	expression tag	UNP A0A482LRD5
А	4	LYS	-	expression tag	UNP A0A482LRD5
А	5	HIS	-	expression tag	UNP A0A482LRD5
А	6	HIS	-	expression tag	UNP A0A482LRD5
А	7	HIS	-	expression tag	UNP A0A482LRD5
А	8	HIS	-	expression tag	UNP A0A482LRD5
А	9	HIS	-	expression tag	UNP A0A482LRD5
А	10	HIS	-	expression tag	UNP A0A482LRD5
А	11	MET	-	expression tag	UNP A0A482LRD5
А	12	HIS	-	expression tag	UNP A0A482LRD5
А	13	HIS	-	expression tag	UNP A0A482LRD5
А	14	HIS	-	expression tag	UNP A0A482LRD5
A	15	HIS	-	expression tag	UNP A0A482LRD5
А	16	HIS	-	expression tag	UNP A0A482LRD5
А	17	HIS	-	expression tag	UNP A0A482LRD5
А	18	GLU	-	expression tag	UNP A0A482LRD5
А	19	ASN	-	expression tag	UNP A0A482LRD5
А	20	LEU	-	expression tag	UNP A0A482LRD5
А	21	TYR	-	expression tag	UNP A0A482LRD5
А	22	PHE	-	expression tag	UNP A0A482LRD5
А	23	GLN	-	expression tag	UNP A0A482LRD5



6V	10
----	----

Continu	ed from pre	vious page			
Chain	Residue	Modelled	Actual	Comment	Reference
А	24	GLY	-	expression tag	UNP A0A482LRD5
В	3	MET	-	expression tag	UNP A0A482LRD5
В	4	LYS	-	expression tag	UNP A0A482LRD5
В	5	HIS	-	expression tag	UNP A0A482LRD5
В	6	HIS	-	expression tag	UNP A0A482LRD5
В	7	HIS	-	expression tag	UNP A0A482LRD5
В	8	HIS	-	expression tag	UNP A0A482LRD5
В	9	HIS	-	expression tag	UNP A0A482LRD5
В	10	HIS	-	expression tag	UNP A0A482LRD5
В	11	MET	-	expression tag	UNP A0A482LRD5
В	12	HIS	-	expression tag	UNP A0A482LRD5
В	13	HIS	-	expression tag	UNP A0A482LRD5
В	14	HIS	-	expression tag	UNP A0A482LRD5
В	15	HIS	-	expression tag	UNP A0A482LRD5
В	16	HIS	-	expression tag	UNP A0A482LRD5
В	17	HIS	-	expression tag	UNP A0A482LRD5
В	18	GLU	-	expression tag	UNP A0A482LRD5
В	19	ASN	-	expression tag	UNP A0A482LRD5
В	20	LEU	-	expression tag	UNP A0A482LRD5
В	21	TYR	-	expression tag	UNP A0A482LRD5
В	22	PHE	-	expression tag	UNP A0A482LRD5
В	23	GLN	-	expression tag	UNP A0A482LRD5
В	24	GLY	-	expression tag	UNP A0A482LRD5
С	3	MET	-	expression tag	UNP A0A482LRD5
С	4	LYS	-	expression tag	UNP A0A482LRD5
С	5	HIS	-	expression tag	UNP A0A482LRD5
С	6	HIS	-	expression tag	UNP A0A482LRD5
С	7	HIS	-	expression tag	UNP A0A482LRD5
С	8	HIS	-	expression tag	UNP A0A482LRD5
С	9	HIS	-	expression tag	UNP A0A482LRD5
С	10	HIS	-	expression tag	UNP A0A482LRD5
С	11	MET	-	expression tag	UNP A0A482LRD5
С	12	HIS	-	expression tag	UNP A0A482LRD5
С	13	HIS	-	expression tag	UNP A0A482LRD5
С	14	HIS	-	expression tag	UNP A0A482LRD5
С	15	HIS	-	expression tag	UNP A0A482LRD5
С	16	HIS	-	expression tag	UNP A0A482LRD5
С	17	HIS	-	expression tag	UNP A0A482LRD5
С	18	GLU	-	expression tag	UNP A0A482LRD5
C	19	ASN	-	expression tag	UNP A0A482LRD5
C	20	LEU	-	expression tag	UNP A0A482LRD5
С	21	TYR	-	expression tag	UNP A0A482LRD5




Chain	Residue	Modelled	Actual	Comment	Reference
			Actual		
С	22	PHE	-	expression tag	UNP A0A482LRD5
С	23	GLN	-	expression tag	UNP A0A482LRD5
С	24	GLY	-	expression tag	UNP A0A482LRD5
D	3	MET	-	expression tag	UNP A0A482LRD5
D	4	LYS	-	expression tag	UNP A0A482LRD5
D	5	HIS	-	expression tag	UNP A0A482LRD5
D	6	HIS	-	expression tag	UNP A0A482LRD5
D	7	HIS	-	expression tag	UNP A0A482LRD5
D	8	HIS	-	expression tag	UNP A0A482LRD5
D	9	HIS	-	expression tag	UNP A0A482LRD5
D	10	HIS	-	expression tag	UNP A0A482LRD5
D	11	MET	-	expression tag	UNP A0A482LRD5
D	12	HIS	-	expression tag	UNP A0A482LRD5
D	13	HIS	-	expression tag	UNP A0A482LRD5
D	14	HIS	-	expression tag	UNP A0A482LRD5
D	15	HIS	-	expression tag	UNP A0A482LRD5
D	16	HIS	_	expression tag	UNP A0A482LRD5
D	17	HIS	_	expression tag	UNP A0A482LRD5
D	18	GLU	_	expression tag	UNP A0A482LRD5
D	19	ASN	_	expression tag	UNP A0A482LRD5
D	20	LEU	-	expression tag	UNP A0A482LRD5
D	21	TYR	-	expression tag	UNP A0A482LRD5
D	22	PHE	-	expression tag	UNP A0A482LRD5
D	23	GLN	-	expression tag	UNP A0A482LRD5
D	24	GLY	-	expression tag	UNP A0A482LRD5

Continued from previous page...

• Molecule 2 is (1aR,7bS)-5-fluoro-2-hydroxy-1,1a,2,7b-tetrahydrocyclopropa[c][1,2]benzo xaborinine-4-carboxylic acid (three-letter code: RM9) (formula: C₁₀H₈BFO₄) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Δ	1	Total	В	С	F	0	0	0
	Л	1	16	1	10	1	4	0	0
2	В	1	Total	В	С	F	Ο	0	0
	2 D	1	16	1	10	1	4	0	0
2	С	1	Total	В	С	F	Ο	0	0
	U	1	16	1	10	1	4	0	0
0	Л	1	Total	В	С	F	Ο	0	0
	D	1	16	1	10	1	4	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	2	Total Mg 2 2	0	0
3	В	2	Total Mg 2 2	0	0
3	С	1	Total Mg 1 1	0	0
3	D	3	Total Mg 3 3	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	А	1	Total C 1 1	1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	1	Total Cl 1 1	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	385	Total O 385 385	0	0
5	В	388	Total O 388 388	0	0
5	С	329	Total O 329 329	0	0
5	D	361	Total O 361 361	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: OXA family beta-lactamase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	58.44Å 105.55 Å 94.40 Å	Depositor
a, b, c, α , β , γ	90.00° 107.92° 90.00°	Depositor
Resolution (Å)	38.85 - 1.80	Depositor
Resolution (A)	38.85 - 1.80	EDS
% Data completeness	96.5(38.85 - 1.80)	Depositor
(in resolution range)	85.2 (38.85-1.80)	EDS
R _{merge}	0.17	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$0.91 (at 1.81 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.158 , 0.224	Depositor
II, II, <i>free</i>	0.159 , 0.222	DCC
R_{free} test set	1959 reflections (2.02%)	wwPDB-VP
Wilson B-factor $(Å^2)$	8.0	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36, 59.5	EDS
L-test for twinning ²	$< L > = 0.49, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9528	wwPDB-VP
Average B, all atoms $(Å^2)$	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: RM9, CL, KCX, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.43	0/2029	0.56	0/2743
1	В	0.43	0/2014	0.56	0/2723
1	С	0.42	0/2065	0.53	0/2792
1	D	0.43	0/2031	0.57	0/2746
All	All	0.43	0/8139	0.56	0/11004

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1993	0	1940	8	0
1	В	1978	0	1928	6	0
1	С	2026	0	1964	3	0
1	D	1994	0	1938	7	0
2	А	16	0	0	0	0
2	В	16	0	0	1	0
2	С	16	0	0	0	0
2	D	16	0	0	0	0
3	А	2	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	2	0	0	0	0
3	С	1	0	0	0	0
3	D	3	0	0	0	0
4	А	1	0	0	0	0
4	С	1	0	0	0	0
5	А	385	0	0	0	3
5	В	388	0	0	0	0
5	С	329	0	0	0	4
5	D	361	0	0	0	1
All	All	9528	0	7770	24	4

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A + 1	A + 0	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:23:GLN:HA	1:A:52:GLN:O	1.91	0.70
1:A:69:ALA:HB2	1:A:212:SER:HB2	1.83	0.60
1:D:216:GLU:OE1	1:D:218:LYS:NZ	2.33	0.59
1:B:115:MET:HG2	1:B:123:TYR:OH	2.08	0.52
1:C:31:TRP:HB2	1:C:57:ASN:HB3	1.92	0.52
1:A:115:MET:HG2	1:A:123:TYR:OH	2.09	0.51
1:D:115:MET:HG2	1:D:123:TYR:OH	2.11	0.51
1:D:31:TRP:HB2	1:D:57:ASN:HB3	1.94	0.49
1:B:31:TRP:HB2	1:B:57:ASN:HB3	1.93	0.49
1:A:31:TRP:HB2	1:A:57:ASN:HB3	1.98	0.45
1:A:177:TYR:CZ	1:A:232:VAL:HG21	2.51	0.45
1:A:73:KCX:CX	1:A:120:VAL:HG22	2.46	0.44
1:C:220:GLY:O	1:C:238:ASN:HA	2.18	0.44
1:B:220:GLY:O	1:B:238:ASN:HA	2.19	0.43
1:A:146:ASN:OD1	1:A:162:ILE:HD12	2.19	0.43
1:D:59:LEU:HD23	1:D:59:LEU:HA	1.79	0.43
1:D:27:GLU:HA	1:D:56:THR:O	2.20	0.42
1:D:177:TYR:CZ	1:D:232:VAL:HG21	2.54	0.42
1:C:27:GLU:HA	1:C:56:THR:O	2.20	0.42
1:A:67:LEU:HD12	1:A:215:ILE:HG12	2.02	0.42
1:B:48:ASN:HB2	1:B:233:TRP:CH2	2.55	0.42
1:B:180:LYS:HA	1:B:180:LYS:HD3	1.90	0.41
1:D:220:GLY:O	1:D:238:ASN:HA	2.21	0.41
1:B:70:SER:CB	2:B:301:RM9:O13	2.69	0.40



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:700:HOH:O	5:C:688:HOH:O[1_455]	1.71	0.49
5:A:606:HOH:O	5:C:630:HOH:O[1_455]	2.13	0.07
5:A:633:HOH:O	5:C:674:HOH:O[1_455]	2.16	0.04
5:C:635:HOH:O	5:D:645:HOH:O[1 655]	2.17	0.03

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	241/263~(92%)	235~(98%)	6(2%)	0	100	100
1	В	239/263~(91%)	235~(98%)	4(2%)	0	100	100
1	С	244/263~(93%)	239~(98%)	5(2%)	0	100	100
1	D	241/263~(92%)	236~(98%)	5(2%)	0	100	100
All	All	965/1052~(92%)	945 (98%)	20~(2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	211/230~(92%)	210 (100%)	1 (0%)	88	87
1	В	209/230~(91%)	209 (100%)	0	100	100



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		ntiles
1	С	214/230~(93%)	214 (100%)	0		100	100
1	D	211/230 (92%)	210 (100%)	1 (0%)		88	87
All	All	845/920~(92%)	843 (100%)	2~(0%)		93	92

Continued from previous page...

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	118	SER
1	D	118	SER

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	В	ond leng	gths	Bond angles		
	Type	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
1	KCX	С	73	1	9,11,12	2.39	2 (22%)	5,12,14	1.25	1 (20%)
1	KCX	А	73	1	9,11,12	2.21	2 (22%)	5,12,14	1.75	1 (20%)
1	KCX	В	73	1	9,11,12	2.28	2 (22%)	5,12,14	1.41	1 (20%)
1	KCX	D	73	1	9,11,12	2.32	2 (22%)	5,12,14	1.40	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	С	73	1	-	0/9/10/12	-
1	KCX	А	73	1	-	0/9/10/12	-
1	KCX	В	73	1	-	0/9/10/12	-
1	KCX	D	73	1	-	0/9/10/12	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	С	73	KCX	CX-NZ	6.22	1.46	1.35
1	D	73	KCX	CX-NZ	6.06	1.45	1.35
1	В	73	KCX	CX-NZ	5.63	1.45	1.35
1	А	73	KCX	CX-NZ	5.60	1.45	1.35
1	В	73	KCX	OQ1-CX	3.28	1.27	1.21
1	С	73	KCX	OQ1-CX	3.05	1.27	1.21
1	А	73	KCX	OQ1-CX	2.99	1.27	1.21
1	D	73	KCX	OQ1-CX	2.57	1.26	1.21

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
1	А	73	KCX	OQ1-CX-NZ	-3.59	119.39	124.96
1	D	73	KCX	OQ1-CX-NZ	-3.01	120.29	124.96
1	В	73	KCX	OQ1-CX-NZ	-2.64	120.86	124.96
1	С	73	KCX	OQ1-CX-NZ	-2.55	121.00	124.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	А	73	KCX	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.



6V1O

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bond lengths			Bond angles		
INIOI	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	RM9	В	301	1	15,18,18	1.62	4 (26%)	19,28,28	1.39	4 (21%)
2	RM9	D	301	1	15,18,18	1.93	4 (26%)	19,28,28	1.23	2 (10%)
2	RM9	С	301	1	15,18,18	1.83	5 (33%)	19,28,28	1.20	2 (10%)
2	RM9	А	301	1	15,18,18	2.11	4 (26%)	19,28,28	1.41	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RM9	В	301	1	-	2/4/21/21	0/2/3/3
2	RM9	D	301	1	-	2/4/21/21	0/2/3/3
2	RM9	С	301	1	-	2/4/21/21	0/2/3/3
2	RM9	А	301	1	-	2/4/21/21	0/2/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	А	301	RM9	C09-C16	-5.35	1.44	1.52
2	D	301	RM9	C04-C05	4.28	1.46	1.39
2	А	301	RM9	C04-C05	3.84	1.45	1.39
2	D	301	RM9	C09-C16	-3.48	1.47	1.52
2	В	301	RM9	C09-C16	-3.35	1.47	1.52
2	С	301	RM9	C04-C02	3.25	1.58	1.50
2	С	301	RM9	C09-C16	-3.13	1.47	1.52
2	С	301	RM9	C04-C05	3.03	1.44	1.39
2	В	301	RM9	C04-C05	2.99	1.43	1.39
2	С	301	RM9	O01-C02	-2.62	1.22	1.30
2	D	301	RM9	O01-C02	-2.55	1.22	1.30
2	В	301	RM9	O01-C02	-2.50	1.22	1.30
2	D	301	RM9	C04-C02	2.41	1.56	1.50
2	А	301	RM9	C07-C05	2.33	1.42	1.37



	Chain	-	1 0	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	С	301	RM9	C07-C05	2.24	1.42	1.37
2	А	301	RM9	O01-C02	-2.18	1.23	1.30
2	В	301	RM9	C15-C16	-2.18	1.46	1.50

Continued from previous page...

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	301	RM9	C09-C16-C14	2.66	122.62	118.34
2	А	301	RM9	C07-C05-C04	-2.54	119.04	123.58
2	А	301	RM9	O03-C02-C04	-2.34	115.39	121.16
2	D	301	RM9	C07-C05-C04	-2.30	119.48	123.58
2	В	301	RM9	C07-C05-C04	-2.19	119.66	123.58
2	В	301	RM9	C05-C04-C02	-2.18	117.66	121.86
2	В	301	RM9	O03-C02-C04	-2.17	115.79	121.16
2	С	301	RM9	C07-C05-C04	-2.14	119.77	123.58
2	А	301	RM9	C08-C09-C16	2.09	126.64	121.81
2	С	301	RM9	B12-O11-C10	-2.06	116.23	119.82
2	D	301	RM9	O03-C02-C04	-2.03	116.16	121.16
2	В	301	RM9	C08-C09-C16	2.01	126.45	121.81

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	301	RM9	O03-C02-C04-C05
2	С	301	RM9	O01-C02-C04-C05
2	В	301	RM9	O03-C02-C04-C05
2	С	301	RM9	O03-C02-C04-C05
2	А	301	RM9	O01-C02-C04-C05
2	В	301	RM9	O01-C02-C04-C05
2	D	301	RM9	O01-C02-C04-C05
2	D	301	RM9	O03-C02-C04-C05

There are no ring outliers.


1 monomer is involved in 1 short contact:

[Mol	Chain	Res	Type	Clashes	Symm-Clashes
	2	В	301	RM9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In




addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	242/263~(92%)	-0.59	1 (0%) 92 90	5, 11, 25, 56	0
1	В	241/263~(91%)	-0.57	2 (0%) 86 84	5, 10, 26, 35	0
1	С	244/263~(92%)	-0.19	9 (3%) 41 36	9, 17, 35, 68	0
1	D	241/263~(91%)	-0.49	2 (0%) 86 84	6, 12, 30, 47	0
All	All	968/1052~(92%)	-0.46	14 (1%) 75 72	5, 13, 30, 68	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	21	TYR	6.1
1	С	97	GLY	4.1
1	С	95	TRP	3.3
1	С	22	PHE	3.2
1	С	99	THR	2.7
1	D	36	THR	2.7
1	С	102	ILE	2.5
1	А	23	GLN	2.4
1	D	39	LYS	2.3
1	В	101	ASP	2.3
1	С	87	LYS	2.2
1	С	24	GLY	2.2
1	С	23	GLN	2.1
1	В	245	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
1	KCX	С	73	12/13	0.94	0.09	$10,\!13,\!17,\!18$	0
1	KCX	В	73	12/13	0.97	0.08	5,6,9,10	0
1	KCX	D	73	12/13	0.97	0.07	$5,\!6,\!8,\!9$	0
1	KCX	А	73	12/13	0.98	0.06	$3,\!5,\!7,\!7$	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
3	MG	В	303	1/1	0.85	0.14	43,43,43,43	0
3	MG	D	303	1/1	0.88	0.14	41,41,41,41	1
3	MG	D	304	1/1	0.90	0.15	41,41,41,41	1
2	RM9	С	301	16/16	0.93	0.09	13,16,20,20	0
3	MG	С	302	1/1	0.94	0.08	24,24,24,24	0
2	RM9	D	301	16/16	0.95	0.08	6,9,11,13	0
2	RM9	А	301	16/16	0.96	0.08	6,8,11,11	0
3	MG	А	302	1/1	0.96	0.14	34,34,34,34	0
2	RM9	В	301	16/16	0.97	0.07	$7,\!12,\!15,\!16$	0
3	MG	А	303	1/1	0.97	0.07	$15,\!15,\!15,\!15$	1
3	MG	D	302	1/1	0.98	0.04	11, 11, 11, 11	0
4	CL	С	303	1/1	0.98	0.04	16, 16, 16, 16	0
4	CL	А	304	1/1	0.99	0.06	11,11,11,11	0
3	MG	В	302	1/1	0.99	0.05	$19,\!19,\!19,\!19$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















6.5 Other polymers (i)

There are no such residues in this entry.

