



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2023 – 05:00 pm GMT

PDB ID : 2V2G
Title : Crystal Structure of the C45S mutant of the Peroxiredoxin 6 of *Arenicola Marina*. Monoclinic form
Authors : Smeets, A.; Declercq, J.P.
Deposited on : 2007-06-05
Resolution : 1.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	233	86% 9% 6%
1	B	233	80% 12% 6%
1	C	233	88% 6% 6%
1	D	233	85% 9% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	C	1224	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7908 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEROXIREDOXIN 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	220	1703	1087	288	319	9	0	0	0
1	B	218	1694	1082	285	318	9	0	1	0
1	C	220	1703	1087	288	319	9	0	0	0
1	D	219	1697	1084	287	317	9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	SER	CYS	engineered mutation	UNP Q1AN22
B	45	SER	CYS	engineered mutation	UNP Q1AN22
C	45	SER	CYS	engineered mutation	UNP Q1AN22
D	45	SER	CYS	engineered mutation	UNP Q1AN22

- Molecule 2 is BENZOIC ACID (three-letter code: BEZ) (formula: C₇H₆O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C O	0	0
			9	7 2		
2	B	1	Total	C O	0	0
			9	7 2		
2	C	1	Total	C O	0	0
			9	7 2		
2	D	1	Total	C O	0	0
			9	7 2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	252	Total	O	0	0
			252	252		
4	B	255	Total	O	0	0
			255	255		
4	C	261	Total	O	0	0
			261	261		
4	D	283	Total	O	0	0
			283	283		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PEROXIREDOXIN 6

Chain A:  86% 9% 6%



- Molecule 1: PEROXIREDOXIN 6

Chain B:  80% 12% 6%




- Molecule 1: PEROXIREDOXIN 6

Chain C:  88% 6% 6%



- Molecule 1: PEROXIREDOXIN 6

Chain D:  85% 9% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	127.25Å 83.23Å 98.35Å 90.00° 100.71° 90.00°	Depositor
Resolution (Å)	13.96 – 1.60 13.96 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.4 (13.96-1.60) 99.4 (13.96-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.160 , 0.197 0.169 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7908	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: BEZ, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	0/1742	0.88	1/2362 (0.0%)
1	B	0.93	2/1736 (0.1%)	0.92	2/2355 (0.1%)
1	C	0.90	0/1742	0.90	4/2362 (0.2%)
1	D	0.87	0/1736	0.93	6/2354 (0.3%)
All	All	0.90	2/6956 (0.0%)	0.91	13/9433 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	86	CYS	CB-SG	-6.22	1.71	1.82
1	B	105	GLU	CG-CD	5.90	1.60	1.51

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	40	ASP	CB-CG-OD1	7.38	124.95	118.30
1	A	15	ASP	CB-CG-OD1	6.92	124.53	118.30
1	D	181	ASP	CB-CG-OD2	-6.29	112.64	118.30
1	D	15	ASP	CB-CG-OD1	5.68	123.42	118.30
1	C	15	ASP	CB-CG-OD1	5.53	123.27	118.30
1	B	40	ASP	CB-CG-OD2	-5.31	113.52	118.30
1	C	101	ASP	CB-CG-OD2	-5.31	113.53	118.30
1	C	136	ASP	CB-CG-OD1	5.25	123.03	118.30
1	D	39	ARG	NE-CZ-NH2	-5.09	117.76	120.30
1	D	40	ASP	CB-CG-OD2	-5.09	113.72	118.30
1	C	40	ASP	CB-CG-OD1	5.07	122.86	118.30
1	D	181	ASP	CB-CG-OD1	5.07	122.86	118.30
1	B	62	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1703	0	1723	11	0
1	B	1694	0	1711	24	0
1	C	1703	0	1723	12	0
1	D	1697	0	1718	17	0
2	A	9	0	5	2	0
2	B	9	0	5	2	0
2	C	9	0	5	2	0
2	D	9	0	5	2	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
3	C	12	0	16	4	0
4	A	252	0	0	2	0
4	B	255	0	0	5	0
4	C	261	0	0	2	0
4	D	283	0	0	5	0
All	All	7908	0	6927	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:153:PHE:HD1	4:D:2208:HOH:O	1.43	1.00
1:C:142:SER:OG	3:C:1224:GOL:H11	1.65	0.97
1:B:46:THR:HG22	1:B:87:LEU:HD12	1.51	0.93
1:C:40:ASP:OD2	1:C:77:HIS:HD2	1.66	0.78
1:B:40:ASP:OD2	1:B:77:HIS:HD2	1.67	0.78
2:B:1220:BEZ:H2	4:B:2080:HOH:O	1.84	0.78
2:C:1222:BEZ:H2	4:C:2069:HOH:O	1.84	0.77
2:D:1221:BEZ:H2	4:D:2083:HOH:O	1.84	0.77
1:B:9:PHE:CD2	1:B:10:PRO:HD2	2.23	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:ASP:OD2	1:A:77:HIS:HD2	1.71	0.73
1:D:46:THR:HG22	1:D:87:LEU:HD12	1.70	0.72
1:D:40:ASP:OD2	1:D:77:HIS:HD2	1.73	0.72
2:A:1222:BEZ:H2	4:A:2066:HOH:O	1.92	0.70
1:C:193:GLU:O	1:C:196:THR:HG22	1.92	0.68
1:C:142:SER:OG	3:C:1224:GOL:C1	2.43	0.66
1:B:153:PHE:HD1	4:B:2193:HOH:O	1.80	0.65
1:D:153:PHE:CD1	4:D:2208:HOH:O	2.27	0.62
1:B:9:PHE:CG	1:B:10:PRO:HD2	2.34	0.61
3:C:1224:GOL:H32	1:D:144:LEU:HD12	1.82	0.61
1:B:60:LYS:HD2	4:B:2045:HOH:O	2.02	0.59
1:B:11:ASN:HD21	1:B:22:LYS:HE2	1.70	0.57
1:A:117:GLU:CD	2:A:1222:BEZ:H4	2.26	0.56
1:B:118:ARG:NE	4:B:2177:HOH:O	2.38	0.56
1:B:196:THR:CG2	1:B:197:LEU:N	2.69	0.56
1:D:153:PHE:CD2	1:D:156:ILE:HD12	2.41	0.56
1:D:203:VAL:HG22	1:D:214:ARG:HG2	1.89	0.55
1:B:61:LYS:NZ	1:B:62:ARG:HH12	2.05	0.54
3:C:1224:GOL:H32	1:D:144:LEU:CD1	2.38	0.53
1:B:153:PHE:CD2	1:B:156:ILE:HD12	2.44	0.53
1:D:169:LYS:HE3	1:D:220:LYS:HG3	1.92	0.52
1:C:40:ASP:OD2	1:C:77:HIS:CD2	2.56	0.52
1:C:153:PHE:CD1	1:C:156:ILE:HD12	2.45	0.52
1:B:196:THR:HG22	1:B:197:LEU:N	2.25	0.51
1:A:40:ASP:OD2	1:A:77:HIS:CD2	2.58	0.51
1:D:114:ASP:CG	1:D:115:PRO:HD2	2.31	0.51
1:B:11:ASN:ND2	1:B:22:LYS:HE2	2.26	0.50
1:D:219:PRO:O	1:D:220:LYS:HB2	2.12	0.50
1:D:22:LYS:NZ	4:D:2042:HOH:O	2.36	0.49
1:C:15:ASP:HB2	4:C:2025:HOH:O	2.12	0.49
1:B:130:VAL:HB	1:B:143:ILE:HG12	1.94	0.49
1:A:169:LYS:HE3	1:A:220:LYS:O	2.13	0.48
1:A:114:ASP:CG	1:A:115:PRO:HD2	2.34	0.47
1:A:153:PHE:CD1	1:A:156:ILE:HD12	2.49	0.47
1:B:40:ASP:OD2	1:B:77:HIS:CD2	2.58	0.46
1:A:13:GLU:HG2	1:A:22:LYS:HD3	1.97	0.46
1:B:117:GLU:CD	2:B:1220:BEZ:H4	2.35	0.46
1:B:203:VAL:HG22	1:B:214:ARG:HG2	1.98	0.46
1:B:71:CYS:O	1:B:102[A]:GLU:HG2	2.17	0.45
1:A:105:GLU:CD	4:A:2139:HOH:O	2.55	0.45
1:C:117:GLU:CD	2:C:1222:BEZ:H4	2.38	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:196:THR:CG2	1:C:197:LEU:N	2.80	0.44
1:A:13:GLU:HG2	1:A:22:LYS:CD	2.47	0.43
1:B:21:LEU:HD12	1:B:21:LEU:C	2.37	0.43
1:C:170:LYS:NZ	1:C:193:GLU:OE2	2.39	0.43
1:A:34:LEU:HD13	1:A:67:ILE:HG23	2.01	0.43
1:D:77:HIS:HE1	1:D:99:ILE:O	2.02	0.43
1:D:61:LYS:HD2	4:D:2100:HOH:O	2.19	0.42
1:D:40:ASP:OD2	1:D:77:HIS:CD2	2.63	0.42
1:D:13:GLU:HG2	1:D:22:LYS:HD3	2.02	0.42
1:B:61:LYS:HZ3	1:B:62:ARG:NH1	2.17	0.41
1:B:114:ASP:CG	1:B:115:PRO:HD2	2.41	0.41
1:D:117:GLU:CD	2:D:1221:BEZ:H4	2.41	0.41
1:B:77:HIS:HE1	1:B:99:ILE:O	2.03	0.41
1:A:206:VAL:HG11	1:A:213:LEU:HD22	2.02	0.41
1:C:153:PHE:CD1	1:C:156:ILE:CD1	3.03	0.41
1:B:118:ARG:CZ	4:B:2177:HOH:O	2.68	0.40
1:C:196:THR:HG23	1:C:197:LEU:N	2.35	0.40
1:B:158:ARG:HD2	1:B:177:TRP:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/233 (94%)	213 (98%)	5 (2%)	0	100	100
1	B	217/233 (93%)	213 (98%)	4 (2%)	0	100	100
1	C	218/233 (94%)	215 (99%)	3 (1%)	0	100	100
1	D	217/233 (93%)	213 (98%)	4 (2%)	0	100	100
All	All	870/932 (93%)	854 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/200 (94%)	186 (98%)	3 (2%)	62	41
1	B	188/200 (94%)	185 (98%)	3 (2%)	62	41
1	C	189/200 (94%)	187 (99%)	2 (1%)	73	57
1	D	188/200 (94%)	187 (100%)	1 (0%)	88	80
All	All	754/800 (94%)	745 (99%)	9 (1%)	71	54

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	GLU
1	A	87	LEU
1	A	196	THR
1	B	15	ASP
1	B	87	LEU
1	B	196	THR
1	C	87	LEU
1	C	220	LYS
1	D	87	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	168	GLN
1	A	178	GLN
1	B	77	HIS
1	B	178	GLN
1	C	77	HIS
1	C	168	GLN
1	C	178	GLN
1	D	77	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BEZ	C	1222	-	9,9,9	1.12	1 (11%)	11,11,11	2.38	5 (45%)
2	BEZ	D	1221	-	9,9,9	1.28	1 (11%)	11,11,11	2.33	4 (36%)
3	GOL	C	1224	-	5,5,5	0.78	0	5,5,5	1.86	1 (20%)
2	BEZ	A	1222	-	9,9,9	1.20	1 (11%)	11,11,11	2.04	4 (36%)
3	GOL	A	1223	-	5,5,5	0.51	0	5,5,5	0.30	0
2	BEZ	B	1220	-	9,9,9	0.87	0	11,11,11	2.18	4 (36%)
3	GOL	B	1221	-	5,5,5	0.38	0	5,5,5	0.65	0
3	GOL	C	1223	-	5,5,5	0.37	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEZ	C	1222	-	-	1/4/4/4	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEZ	D	1221	-	-	1/4/4/4	0/1/1/1
3	GOL	C	1224	-	-	4/4/4/4	-
2	BEZ	A	1222	-	-	0/4/4/4	0/1/1/1
3	GOL	A	1223	-	-	2/4/4/4	-
2	BEZ	B	1220	-	-	0/4/4/4	0/1/1/1
3	GOL	B	1221	-	-	2/4/4/4	-
3	GOL	C	1223	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1222	BEZ	O1-C	3.05	1.31	1.22
2	D	1221	BEZ	O1-C	2.77	1.30	1.22
2	C	1222	BEZ	O1-C	2.66	1.30	1.22

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1222	BEZ	O1-C-C1	-5.15	107.72	121.45
2	A	1222	BEZ	O1-C-C1	-4.30	110.00	121.45
2	D	1221	BEZ	O1-C-C1	-3.81	111.29	121.45
2	B	1220	BEZ	O1-C-C1	-3.71	111.58	121.45
2	C	1222	BEZ	O2-C-O1	3.55	131.22	123.35
2	D	1221	BEZ	O2-C-O1	3.53	131.17	123.35
2	D	1221	BEZ	C6-C1-C2	3.45	123.49	118.59
2	B	1220	BEZ	C6-C1-C2	3.31	123.31	118.59
2	B	1220	BEZ	O2-C-O1	3.26	130.58	123.35
3	C	1224	GOL	O1-C1-C2	2.90	124.11	110.20
2	D	1221	BEZ	C3-C2-C1	-2.83	116.99	120.34
2	A	1222	BEZ	O2-C-O1	2.70	129.33	123.35
2	A	1222	BEZ	C6-C1-C2	2.66	122.37	118.59
2	C	1222	BEZ	O2-C-C1	2.36	120.98	114.85
2	C	1222	BEZ	C6-C1-C2	2.30	121.86	118.59
2	A	1222	BEZ	O2-C-C1	2.23	120.64	114.85
2	B	1220	BEZ	C3-C2-C1	-2.13	117.83	120.34
2	C	1222	BEZ	C6-C1-C	-2.03	116.39	120.39

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1221	GOL	O1-C1-C2-C3
3	C	1224	GOL	O1-C1-C2-C3
3	B	1221	GOL	O1-C1-C2-O2
3	C	1224	GOL	O1-C1-C2-O2
3	A	1223	GOL	O1-C1-C2-C3
3	C	1224	GOL	C1-C2-C3-O3
3	A	1223	GOL	O1-C1-C2-O2
3	C	1224	GOL	O2-C2-C3-O3
2	D	1221	BEZ	O1-C-C1-C2
2	C	1222	BEZ	O1-C-C1-C2

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1222	BEZ	2	0
2	D	1221	BEZ	2	0
3	C	1224	GOL	4	0
2	A	1222	BEZ	2	0
2	B	1220	BEZ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.