



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 05:27 pm GMT

PDB ID : 4V2G  
Title : Tetracycline repressor TetR(D) bound to chlortetracycline and iso- chlortetracycline  
Authors : Werten, S.; Orth, P.; Saenger, W.; Hinrichs, W.  
Deposited on : 2014-10-09  
Resolution : 2.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	 4% 88% 10% •
1	B	207	 2% 84% 10% • 5%






### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TETRACYCLINE REPRESSOR PROTEIN CLASS D

Chain A:  4% 88% 10%



- Molecule 1: TETRACYCLINE REPRESSOR PROTEIN CLASS D

Chain B:  2% 84% 10% 5%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.25Å 69.25Å 182.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.74 – 2.71 19.74 – 2.71	Depositor EDS
% Data completeness (in resolution range)	99.0 (64.74-2.71) 99.4 (19.74-2.71)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.15 (at 2.71Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.175 , 0.252 0.180 , 0.251	Depositor DCC
$R_{free}$ test set	1260 reflections (9.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.5	Xtrriage
Anisotropy	0.216	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3302	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 77.99 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.5940e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CTC, ITC, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/1670	0.79	2/2261 (0.1%)
1	B	0.66	0/1587	0.80	1/2144 (0.0%)
All	All	0.64	0/3257	0.79	3/4405 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	180	ASP	CB-CG-OD1	6.01	123.71	118.30
1	A	171	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	B	180	ASP	CB-CG-OD1	5.18	122.96	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	151	HIS	Peptide
1	B	165	ASN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1641	0	1633	11	0
1	B	1561	0	1558	9	0
2	A	33	0	21	0	0
3	B	33	0	20	1	0
4	B	1	0	0	0	0
5	A	20	0	0	1	0
5	B	13	0	0	2	0
All	All	3302	0	3232	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:222:CTC:H6'1	3:B:222:CTC:CL7	2.11	0.87
1:A:148:GLN:O	1:A:152:THR:HG23	1.91	0.70
1:A:16:LEU:HD23	1:A:25:LEU:HD12	1.77	0.66
1:B:16:LEU:HD23	1:B:25:LEU:HD12	1.76	0.66
1:A:16:LEU:HD23	1:A:25:LEU:CD1	2.32	0.59
1:B:16:LEU:HD23	1:B:25:LEU:CD1	2.32	0.59
1:A:158:ARG:HD2	1:B:53:ASP:OD2	2.04	0.57
1:A:104:ARG:CG	1:A:105:PRO:HD2	2.39	0.53
1:A:104:ARG:HG2	1:A:105:PRO:HD2	1.91	0.53
1:B:98:LYS:NZ	5:B:2002:HOH:O	2.36	0.49
1:A:87:ARG:NH2	5:A:2008:HOH:O	2.46	0.45
1:A:104:ARG:HH11	1:B:151:HIS:CD2	2.34	0.45
1:B:148:GLN:HG3	1:B:181:ASP:O	2.17	0.45
1:B:129:ASP:HB2	5:B:2010:HOH:O	2.18	0.44
1:A:21:GLY:HA2	1:A:94:ARG:HG2	2.00	0.44
1:A:101:LEU:HG	1:B:150:GLU:HG3	2.01	0.42
1:B:92:ARG:HH11	1:B:92:ARG:HG2	1.85	0.42
1:A:148:GLN:OE1	1:A:181:ASP:O	2.38	0.42

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/207 (99%)	193 (94%)	12 (6%)	0	100	100
1	B	192/207 (93%)	186 (97%)	6 (3%)	0	100	100
All	All	397/414 (96%)	379 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/173 (100%)	158 (91%)	15 (9%)	10	23
1	B	165/173 (95%)	153 (93%)	12 (7%)	14	31
All	All	338/346 (98%)	311 (92%)	27 (8%)	12	26

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	55	LEU
1	A	66	TYR
1	A	79	LEU
1	A	87	ARG
1	A	94	ARG
1	A	103	THR
1	A	104	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	107	GLU
1	A	142	LEU
1	A	152	THR
1	A	157	ASP
1	A	179	SER
1	A	201	LEU
1	A	206	GLN
1	B	3	ARG
1	B	4	LEU
1	B	14	LEU
1	B	55	LEU
1	B	66	TYR
1	B	94	ARG
1	B	107	GLU
1	B	128	ARG
1	B	131	LEU
1	B	165	ASN
1	B	175	GLN
1	B	201	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	200	GLN
1	B	151	HIS
1	B	165	ASN
1	B	200	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CTC	B	222	4	34,36,36	4.23	13 (38%)	42,60,60	2.22	19 (45%)
2	ITC	A	222	-	32,36,36	4.02	11 (34%)	36,59,59	3.10	17 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CTC	B	222	4	-	0/8/74/74	0/4/4/4
2	ITC	A	222	-	-	0/14/71/71	0/4/4/4

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	222	CTC	C4B-C12	-13.54	1.40	1.52
3	B	222	CTC	C6-C6A	-12.24	1.42	1.53
2	A	222	ITC	C4B-C4A	-10.42	1.44	1.53
2	A	222	ITC	C4B-C1	-10.26	1.41	1.55
2	A	222	ITC	C5B-C12	-9.44	1.36	1.50
3	B	222	CTC	C5A-C5B	-8.46	1.38	1.52
2	A	222	ITC	C6B-C11	-7.95	1.33	1.47
2	A	222	ITC	O6-C6	-6.99	1.41	1.48
3	B	222	CTC	C4B-C4A	-6.99	1.47	1.53
3	B	222	CTC	C4B-C1	-6.76	1.46	1.55
3	B	222	CTC	C4-C3	-6.25	1.39	1.51
2	A	222	ITC	C4-C3	-5.50	1.40	1.51
2	A	222	ITC	C6B-C6A	-3.89	1.34	1.40
2	A	222	ITC	C5-C5A	3.85	1.60	1.53
3	B	222	CTC	C5-C4A	-3.38	1.48	1.53
2	A	222	ITC	C6B-C10	-3.26	1.36	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	222	CTC	C5B-C11	-3.15	1.39	1.47
3	B	222	CTC	C7-CL7	3.10	1.81	1.73
3	B	222	CTC	C5B-C12	2.82	1.39	1.36
3	B	222	CTC	O4B-C4B	2.57	1.46	1.42
2	A	222	ITC	C2-C2'	-2.39	1.42	1.47
3	B	222	CTC	C5-C5A	2.36	1.58	1.54
2	A	222	ITC	C5-C4A	-2.22	1.50	1.53
3	B	222	CTC	C6B-C11	-2.21	1.40	1.46

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	222	ITC	O6-C11-C6B	10.72	116.51	108.20
2	A	222	ITC	O4B-C4B-C4A	-6.30	102.35	110.09
2	A	222	ITC	C10-C6B-C11	4.91	133.28	129.90
2	A	222	ITC	O11-C11-C6B	-4.86	121.90	131.04
2	A	222	ITC	C4A-C4B-C1	4.57	116.30	111.05
2	A	222	ITC	C6B-C6A-C7	-4.41	118.16	120.23
3	B	222	CTC	C11-C5B-C12	4.27	122.18	118.80
3	B	222	CTC	C1-C4B-C12	3.93	114.48	109.88
3	B	222	CTC	O11-C11-C5B	-3.82	114.92	120.78
2	A	222	ITC	O3-C3-C2	-3.71	116.45	122.96
3	B	222	CTC	C2'-C2-C1	3.45	125.06	120.97
2	A	222	ITC	C8-C9-C10	-3.25	117.16	120.50
3	B	222	CTC	C8-C9-C10	-3.19	117.22	120.50
2	A	222	ITC	O12-C12-C5B	-3.14	116.81	121.79
3	B	222	CTC	O3-C3-C2	-3.12	117.48	122.96
2	A	222	ITC	C4A-C5-C5A	-3.09	105.69	111.32
3	B	222	CTC	C9-C10-C6B	3.09	124.14	120.17
3	B	222	CTC	C4B-C4A-C4	-3.07	107.44	111.64
3	B	222	CTC	O2'-C2'-C2	3.04	125.90	120.67
2	A	222	ITC	C5A-C5B-C12	-3.04	105.31	113.43
3	B	222	CTC	O12-C12-C5B	-2.92	119.90	123.90
2	A	222	ITC	C6A-C6B-C11	-2.84	105.88	108.59
3	B	222	CTC	O11-C11-C6B	-2.83	116.64	121.99
3	B	222	CTC	O4B-C4B-C12	-2.81	105.65	110.14
3	B	222	CTC	C4B-C1-C2	2.73	120.09	115.75
3	B	222	CTC	O2'-C2'-N2'	-2.60	116.81	122.88
2	A	222	ITC	O1-C1-C2	-2.49	118.31	123.55
3	B	222	CTC	C8-C7-CL7	-2.43	113.53	118.41
3	B	222	CTC	O6-C6-C6A	2.30	113.04	108.96
2	A	222	ITC	C4B-C1-C2	2.29	119.39	115.75

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	222	CTC	C5-C5A-C5B	2.28	114.83	109.53
3	B	222	CTC	C10-C6B-C6A	-2.23	117.07	119.47
2	A	222	ITC	C8-C7-CL7	-2.19	114.01	118.41
3	B	222	CTC	C6'-C6-C6A	-2.17	106.78	111.60
2	A	222	ITC	O6-C6-C6'	2.06	108.71	106.89
2	A	222	ITC	O4B-C4B-C1	-2.05	100.42	106.40

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	222	CTC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/207 (100%)	-0.29	8 (3%) 39 39	20, 40, 90, 124	0
1	B	196/207 (94%)	-0.47	4 (2%) 65 67	19, 35, 82, 107	0
All	All	403/414 (97%)	-0.37	12 (2%) 50 51	19, 38, 88, 124	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	153	ALA	5.2
1	A	159	PRO	4.8
1	B	152	THR	4.2
1	A	160	ALA	3.9
1	A	156	THR	3.6
1	A	164	GLU	3.0
1	A	107	GLU	3.0
1	B	175	GLN	2.9
1	A	104	ARG	2.7
1	A	162	PRO	2.7
1	A	157	ASP	2.4
1	B	165	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CTC	B	222	33/33	0.93	0.14	23,28,43,53	0
2	ITC	A	222	33/33	0.94	0.14	33,40,61,62	0
4	MG	B	223	1/1	0.97	0.09	20,20,20,20	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.