



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 11:42 pm BST

PDB ID : 4V3Y
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with N-2-(2-(1H-imidazol-1-yl)pyrimidin-4-yl)ethyl-3-(3-chlorophenyl)propan-1-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-10-20
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, BRAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	408	Total 3323	C 2127	N 567	O 607	S 22	0	3	1
1	B	411	Total 3357	C 2148	N 574	O 614	S 21	0	3	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is 3-(3-chlorophenyl)-N-{2-[2-(1H-imidazol-1-yl)pyrimidin-4-yl]ethyl}propan-1-amine (three-letter code: HLW) (formula: $C_{18}H_{20}ClN_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	Cl	N		
4	A	1	24	18	1	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	Cl	N		
4	B	1	24	18	1	5	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	B	1	4	2	2	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	178	178	178	0	0
7	B	228	228	228	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



- Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.85Å 111.05Å 164.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.48 – 1.96 38.49 – 1.96	Depositor EDS
% Data completeness (in resolution range)	78.4 (38.48-1.96) 78.4 (38.49-1.96)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 1.97Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.191 , 0.248 0.191 , 0.249	Depositor DCC
R_{free} test set	2656 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtrriage
Anisotropy	0.053	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7263	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HLW, HEM, ZN, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3422	0.54	0/4643
1	B	0.41	0/3459	0.54	0/4689
All	All	0.40	0/6881	0.54	0/9332

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3323	0	3232	35	0
1	B	3357	0	3275	14	0
2	A	43	0	30	2	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	24	0	20	0	0
4	B	24	0	20	0	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	178	0	0	3	0
7	B	228	0	0	1	0
All	All	7263	0	6643	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:750:HEM:HBB2	2:A:750:HEM:HHC	1.70	0.74
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.73	0.70
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	1.75	0.67
1:B:322:LEU:HD13	1:B:699:ARG:HH21	1.59	0.67
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.77	0.67
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.77	0.67
2:B:750:HEM:HBC2	2:B:750:HEM:HMC2	1.76	0.66
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.79	0.65
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.80	0.63
1:A:652:HIS:ND1	7:A:2152:HOH:O	2.31	0.59
1:A:393:THR:OG1	1:A:394:TYR:N	2.35	0.58
1:A:496:PRO:HA	1:A:499:VAL:HG23	1.86	0.57
1:B:487:GLN:OE1	1:B:514:ARG:NH2	2.38	0.56
1:B:447:LYS:HD2	1:B:540:LEU:HD11	1.87	0.56
1:A:360:GLU:O	1:A:364:GLN:HG3	2.07	0.54
1:A:351:LYS:NZ	1:A:389:GLU:O	2.37	0.53
1:B:593:ILE:HA	1:B:597:ASP:HB2	1.91	0.53
1:A:416:VAL:HG23	1:A:679:ILE:HG23	1.91	0.52
1:A:554:PHE:HB3	7:A:2114:HOH:O	2.10	0.52
1:B:414:ARG:HD3	1:B:678:TRP:CD2	2.44	0.52
1:A:351:LYS:HE2	1:A:392:SER:HA	1.94	0.49
1:A:555:LYS:N	7:A:2114:HOH:O	2.46	0.48
1:A:465:ARG:NH2	1:A:471:ASP:OD2	2.46	0.48
1:B:407:HIS:CE1	1:B:410:ARG:HH11	2.31	0.47
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.97	0.46
1:A:659:ILE:HG13	1:A:689:PRO:HB2	1.97	0.45
1:A:645:LYS:HB2	1:A:645:LYS:HE3	1.72	0.44
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.53	0.44
1:A:299:ARG:HE	1:A:318:LEU:HD13	1.83	0.44
1:A:350:THR:N	1:A:353:GLN:OE1	2.50	0.44
1:B:388:ILE:O	1:B:392:SER:N	2.43	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:516:ARG:HD2	7:B:2140:HOH:O	2.18	0.43
1:A:545:PRO:HG2	1:A:547:ARG:NH1	2.33	0.43
1:A:455:LEU:HD12	1:A:587:TRP:HB3	2.00	0.43
1:A:618:MET:HG2	1:A:625:TRP:CD2	2.54	0.43
1:A:414:ARG:HD3	1:A:678:TRP:CD2	2.54	0.43
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.54	0.42
1:A:445:HIS:C	1:A:445:HIS:CD2	2.92	0.42
1:B:415:CYS:HB2	2:B:750:HEM:ND	2.34	0.42
1:A:553:TRP:HZ3	1:A:557:LEU:HD11	1.85	0.42
1:A:321:THR:HG23	1:A:322:LEU:HG	2.02	0.42
1:A:608:GLU:HG3	1:A:618:MET:HE1	2.02	0.42
1:B:563:GLY:O	1:B:565:PRO:HD3	2.20	0.41
1:A:696:LEU:HA	1:A:696:LEU:HD23	1.95	0.41
1:A:685:GLY:O	1:A:691:PHE:HB2	2.21	0.41
1:A:709:ASP:HA	1:A:710:PRO:HD3	1.89	0.41
1:A:495:ASP:HA	1:A:496:PRO:HD3	1.93	0.41
1:A:500:GLN:O	1:A:503:GLU:HB2	2.21	0.40
1:A:595:VAL:O	1:A:599:CYS:HB2	2.21	0.40
1:A:549:PRO:HB3	1:A:639:TYR:CE1	2.56	0.40
1:B:462:PHE:HB2	1:B:581:ALA:HB3	2.02	0.40
1:A:357:LEU:HD13	1:A:576:GLY:HA2	2.03	0.40
1:B:595:VAL:HG13	1:B:630:LEU:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/422 (96%)	392 (97%)	14 (3%)	0	100	100
1	B	410/422 (97%)	396 (97%)	14 (3%)	0	100	100
All	All	816/844 (97%)	788 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/377 (97%)	355 (97%)	10 (3%)	44	34
1	B	369/377 (98%)	359 (97%)	10 (3%)	44	34
All	All	734/754 (97%)	714 (97%)	20 (3%)	44	34

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	299	ARG
1	A	393	THR
1	A	469	LYS
1	A	476	ASN
1	A	523	LEU
1	A	547	ARG
1	A	550	LYS
1	A	555	LYS
1	A	645	LYS
1	A	715	VAL
1	B	320	SER
1	B	321	THR
1	B	392	SER
1	B	423	LYS
1	B	516	ARG
1	B	540	LEU
1	B	547	ARG
1	B	550	LYS
1	B	619	ARG
1	B	717	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	407	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	A	860	-	1,3,3	1.47	0	0,3,3	0.00	-
5	ACT	B	860	-	1,3,3	1.24	0	0,3,3	0.00	-
3	H4B	B	760	-	16,18,18	0.98	0	11,26,26	2.47	4 (36%)
4	HLW	A	800	2	25,26,26	1.24	2 (8%)	30,33,33	1.96	8 (26%)
4	HLW	B	800	2	25,26,26	1.18	2 (8%)	30,33,33	1.71	5 (16%)
2	HEM	B	750	1,4	27,50,50	2.22	6 (22%)	17,82,82	1.95	3 (17%)
3	H4B	A	760	-	16,18,18	0.93	1 (6%)	11,26,26	2.94	6 (54%)
2	HEM	A	750	1,4	27,50,50	2.12	5 (18%)	17,82,82	2.29	7 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	750	1,4	-	0/6/54/54	-
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	HLW	A	800	2	-	3/9/13/13	0/3/3/3
4	HLW	B	800	2	-	1/9/13/13	0/3/3/3
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
2	HEM	A	750	1,4	-	0/6/54/54	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	750	HEM	C3D-C2D	5.26	1.53	1.37
2	B	750	HEM	C3D-C2D	4.97	1.52	1.37
2	B	750	HEM	C3B-C2B	-4.77	1.33	1.40
2	A	750	HEM	C3B-C2B	-4.35	1.34	1.40
2	B	750	HEM	C3C-C2C	-4.15	1.34	1.40
2	A	750	HEM	C3C-C2C	-4.12	1.34	1.40
2	B	750	HEM	C3C-CAC	3.96	1.55	1.47
2	A	750	HEM	C3C-CAC	3.64	1.55	1.47
2	A	750	HEM	C3B-CAB	3.60	1.55	1.47
4	A	800	HLW	C02-N03	-3.59	1.32	1.36
2	B	750	HEM	C3B-CAB	3.54	1.55	1.47
4	A	800	HLW	C04-N03	-3.22	1.34	1.39
4	B	800	HLW	C02-N03	-2.93	1.33	1.36
4	B	800	HLW	C04-N03	-2.83	1.35	1.39
2	B	750	HEM	CAA-C2A	2.09	1.55	1.52
3	A	760	H4B	C4-N3	2.06	1.36	1.33

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	760	H4B	C4-C4A-N5	5.94	124.11	119.12
3	B	760	H4B	C4-C4A-C8A	5.85	119.76	114.57
4	A	800	HLW	C16-N11-C12	5.36	121.14	114.04
2	A	750	HEM	CBD-CAD-C3D	-5.24	102.82	112.48
4	B	800	HLW	C16-N11-C12	5.15	120.86	114.04
2	B	750	HEM	CBD-CAD-C3D	-5.12	103.04	112.48
4	A	800	HLW	C15-C16-N11	-4.80	118.00	123.96
4	B	800	HLW	C15-C16-N11	-3.98	119.02	123.96
3	A	760	H4B	N3-C2-N1	-3.83	119.42	125.42
3	A	760	H4B	C4-C4A-C8A	3.83	117.97	114.57
2	A	750	HEM	CBA-CAA-C2A	-3.82	105.44	112.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	C1D-C2D-C3D	-3.76	104.38	107.00
2	A	750	HEM	C1D-C2D-C3D	-3.58	104.51	107.00
3	A	760	H4B	C2-N1-C8A	3.31	121.95	114.54
4	A	800	HLW	N13-C12-N03	3.11	118.81	114.78
4	B	800	HLW	N11-C12-N03	3.08	119.58	114.81
3	B	760	H4B	C4-N3-C2	3.06	120.79	115.93
4	A	800	HLW	C20-C21-C22	-2.98	107.64	112.95
3	A	760	H4B	C4-N3-C2	2.90	120.54	115.93
4	B	800	HLW	N11-C12-N13	-2.90	122.79	126.08
2	A	750	HEM	CAD-CBD-CGD	-2.88	107.84	112.67
2	B	750	HEM	CBA-CAA-C2A	-2.68	107.54	112.49
4	A	800	HLW	C04-N03-C12	-2.56	122.61	125.50
3	A	760	H4B	N2-C2-N3	2.46	121.08	117.25
2	A	750	HEM	CMA-C3A-C4A	-2.41	124.76	128.46
3	B	760	H4B	N3-C2-N1	-2.41	121.64	125.42
2	A	750	HEM	CMC-C2C-C3C	2.36	129.09	124.68
2	A	750	HEM	C4A-C3A-C2A	2.32	108.61	107.00
4	A	800	HLW	N11-C12-N13	-2.23	123.56	126.08
4	B	800	HLW	N13-C12-N03	2.20	117.63	114.78
4	A	800	HLW	C05-N01-C02	2.18	112.72	106.20
3	B	760	H4B	C2-N1-C8A	2.07	119.17	114.54
4	A	800	HLW	C17-C14-N13	2.01	118.95	115.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	800	HLW	C2'-C1'-C22-C21
4	A	800	HLW	C6'-C1'-C22-C21
4	A	800	HLW	N19-C20-C21-C22
4	B	800	HLW	C17-C18-N19-C20


There are no ring outliers.


2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	750	HEM	3	0
2	A	750	HEM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/422 (96%)	0.64	48 (11%) 4 7	13, 35, 71, 104	0
1	B	411/422 (97%)	0.05	16 (3%) 39 49	11, 26, 53, 93	0
All	All	819/844 (97%)	0.34	64 (7%) 13 20	11, 30, 68, 104	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	717	LYS	7.7
1	B	348	VAL	7.3
1	A	715	VAL	7.2
1	B	300	PHE	6.3
1	A	488	PRO	6.0
1	A	716	TRP	5.5
1	B	350	THR	4.9
1	A	352	ASP	4.4
1	A	355	PHE	3.9
1	A	388	ILE	3.7
1	A	351	LYS	3.7
1	B	619	ARG	3.6
1	A	486	LYS	3.5
1	B	718	GLY	3.5
1	A	506	ILE	3.5
1	B	620	LYS	3.4
1	B	351	LYS	3.4
1	A	551	PHE	3.4
1	A	712	ASN	3.4
1	A	391	THR	3.4
1	A	350	THR	3.4
1	A	300	PHE	3.2
1	A	490	GLY	3.2
1	A	508	GLN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	299	ARG	3.1
1	A	491	SER	3.0
1	B	302	LYS	3.0
1	A	321	THR	3.0
1	A	489	ASP	3.0
1	A	389	GLU	3.0
1	A	487	GLN	2.9
1	A	390	SER	2.9
1	B	352	ASP	2.9
1	A	553	TRP	2.9
1	A	511	LYS	2.8
1	A	311	VAL	2.7
1	A	386	LYS	2.7
1	A	619	ARG	2.7
1	A	493	LEU	2.6
1	B	299	ARG	2.6
1	A	514	ARG	2.6
1	A	713	THR	2.4
1	A	507	GLN	2.4
1	A	504	ILE	2.4
1	A	354	LEU	2.3
1	A	714	HIS	2.3
1	B	667	ARG	2.2
1	B	349	ARG	2.2
1	B	616	LEU	2.2
1	A	567	VAL	2.2
1	A	393	THR	2.1
1	A	711	TRP	2.1
1	B	611	ALA	2.1
1	A	392	SER	2.1
1	B	615	ASP	2.1
1	A	415	CYS	2.1
1	A	643	SER	2.1
1	A	552	ASP	2.1
1	B	617	ASP	2.1
1	A	385	ASN	2.1
1	A	584	PHE	2.0
1	A	550	LYS	2.0
1	A	681	PRO	2.0
1	A	682	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ACT	A	860	4/4	0.92	0.18	37,41,42,46	0
4	HLW	A	800	24/24	0.94	0.20	10,26,55,77	0
4	HLW	B	800	24/24	0.95	0.15	11,23,51,67	0
5	ACT	B	860	4/4	0.96	0.13	27,33,34,41	0
3	H4B	B	760	17/17	0.97	0.14	11,18,25,26	0
3	H4B	A	760	17/17	0.97	0.13	10,18,26,27	0
2	HEM	B	750	43/43	0.98	0.14	5,16,25,31	0
2	HEM	A	750	43/43	0.98	0.16	9,19,28,30	0
6	ZN	A	1717	1/1	1.00	0.06	25,25,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HLW A 800:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around HLW B 800:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 750:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.