



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2023 – 07:27 PM EST

PDB ID : 1V4W  
Title : Crystal structure of bluefin tuna hemoglobin deoxy form at pH7.5  
Authors : Yokoyama, T.; Chong, K.T.; Miyazaki, Y.; Nakatsukasa, T.; Unzai, S.;  
Miyazaki, G.; Morimoto, H.; Jeremy, R.H.T.; Park, S.Y.  
Deposited on : 2003-11-19  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	144	
1	C	144	
2	B	146	
2	D	146	



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	76	Total	O	0	0
			76	76		
4	C	48	Total	O	0	0
			48	48		
4	D	33	Total	O	0	0
			33	33		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: hemoglobin alpha chain

Chain A:  88% 12%




- Molecule 1: hemoglobin alpha chain

Chain C:  84% 13%




- Molecule 2: hemoglobin beta chain

Chain B:  86% 12%



- Molecule 2: hemoglobin beta chain

Chain D:  83% 13%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.77Å 58.64Å 174.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	13.84 – 1.70	Depositor
% Data completeness (in resolution range)	100.0 (13.84-1.70)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.1.22	Depositor
R, $R_{free}$	0.174 , 0.202	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4843	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/1110	0.81	1/1499 (0.1%)
1	C	0.59	0/1110	0.91	7/1499 (0.5%)
2	B	0.66	0/1168	0.86	6/1585 (0.4%)
2	D	0.57	0/1168	0.76	2/1585 (0.1%)
All	All	0.60	0/4556	0.83	16/6168 (0.3%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	108	ASP	CB-CG-OD2	7.07	124.66	118.30
1	C	28	ASP	CB-CG-OD2	7.03	124.63	118.30
1	C	49	ASP	CB-CG-OD2	6.98	124.58	118.30
2	B	94	ASP	CB-CG-OD2	6.80	124.42	118.30
2	B	106	LEU	CB-CG-CD2	6.60	122.22	111.00
1	A	49	ASP	CB-CG-OD2	6.48	124.13	118.30
2	B	120	ASP	CB-CG-OD2	5.96	123.67	118.30
1	C	34	LEU	CB-CG-CD1	5.88	121.00	111.00
1	C	107	ILE	CG1-CB-CG2	5.58	123.68	111.40
1	C	128	ASP	CB-CG-OD2	5.49	123.24	118.30
2	B	52	ASP	CB-CG-OD2	5.33	123.10	118.30
2	D	79	ASP	CB-CG-OD2	5.32	123.08	118.30
1	C	76	ASP	CB-CG-OD2	5.24	123.01	118.30
1	C	84	ASP	CB-CG-OD2	5.18	122.97	118.30
2	B	68	LEU	CB-CG-CD2	5.17	119.80	111.00
2	D	101	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1088	0	1120	20	0
1	C	1088	0	1120	23	0
2	B	1143	0	1135	11	0
2	D	1143	0	1135	18	0
3	A	43	0	30	1	0
3	B	43	0	30	2	0
3	C	43	0	30	2	0
3	D	43	0	30	1	0
4	A	52	0	0	0	0
4	B	76	0	0	1	0
4	C	48	0	0	1	0
4	D	33	0	0	4	0
All	All	4843	0	4630	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:LEU:HD21	1:C:0:ACE:H1	1.33	1.10
1:A:136:LEU:CD2	1:C:0:ACE:H1	1.84	1.06
1:A:136:LEU:CG	1:C:0:ACE:H1	1.93	0.99
1:A:136:LEU:HD21	1:C:0:ACE:CH3	2.00	0.91
2:D:139:PHE:CE2	2:D:143:ARG:HD3	2.14	0.83
1:A:136:LEU:HG	1:C:0:ACE:H1	1.70	0.72
2:D:143:ARG:NE	2:D:143:ARG:HA	2.11	0.65
2:B:69:HIS:HD2	2:B:72:ASP:OD2	1.81	0.63
1:A:32:ARG:HH11	2:B:127:GLN:HE21	1.49	0.61
2:D:91:LEU:HD12	2:D:95:LYS:HD3	1.83	0.60
2:D:125:GLU:OE2	4:D:159:HOH:O	2.16	0.60
3:C:144:HEM:HBB2	3:C:144:HEM:HHC	1.83	0.60
1:A:2:THR:HG23	1:C:140:GLU:OE1	2.03	0.58
1:C:113:LYS:HE3	4:D:169:HOH:O	2.02	0.58
1:C:105:HIS:HD2	4:D:151:HOH:O	1.87	0.57

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:32:ARG:HH11	2:D:127:GLN:HE21	1.51	0.57
1:A:136:LEU:CD2	1:C:0:ACE:CH3	2.66	0.56
2:B:47:ASP:OD1	2:B:49:SER:OG	2.23	0.56
1:C:32:ARG:HD3	2:D:127:GLN:HE22	1.71	0.56
2:D:19:ASN:HD22	2:D:19:ASN:C	2.09	0.56
1:A:32:ARG:HH11	2:B:127:GLN:NE2	2.03	0.56
1:C:101:LYS:HE2	1:C:101:LYS:H	1.71	0.56
2:B:69:HIS:HE1	4:B:158:HOH:O	1.90	0.55
1:C:59:ALA:O	1:C:62:LYS:HG3	2.08	0.54
2:D:139:PHE:O	2:D:143:ARG:HD2	2.08	0.53
1:A:125:VAL:HG13	2:B:34:VAL:HA	1.91	0.53
1:A:32:ARG:HD3	2:B:127:GLN:HE22	1.73	0.53
1:A:60:HIS:ND1	1:A:63:LYS:HE2	2.23	0.52
2:B:6:GLN:HG2	2:B:125:GLU:OE2	2.08	0.52
1:A:136:LEU:HG	1:C:0:ACE:CH3	2.39	0.52
3:B:147:HEM:HBC2	3:B:147:HEM:HMC1	1.91	0.52
1:A:2:THR:HG23	1:C:140:GLU:CD	2.32	0.50
1:C:0:ACE:C	4:C:174:HOH:O	2.49	0.49
1:C:79:THR:HG23	1:C:140:GLU:HG3	1.95	0.49
3:C:144:HEM:HHC	3:C:144:HEM:CBB	2.43	0.49
2:B:74:ALA:HB2	2:B:85:TYR:HE1	1.77	0.49
1:C:97:PRO:O	1:C:101:LYS:CE	2.62	0.48
1:C:101:LYS:HE2	1:C:101:LYS:N	2.28	0.47
2:D:143:ARG:HA	2:D:143:ARG:CZ	2.44	0.47
1:C:32:ARG:HH11	2:D:127:GLN:NE2	2.12	0.46
2:D:103:PHE:CD1	3:D:147:HEM:HBB2	2.51	0.45
1:A:79:THR:HG23	1:A:140:GLU:HG3	1.98	0.45
3:B:147:HEM:HBC2	3:B:147:HEM:CMC	2.47	0.44
1:A:60:HIS:HE1	3:A:144:HEM:CHA	2.29	0.44
1:A:2:THR:HG23	1:C:140:GLU:OE2	2.18	0.43
2:D:74:ALA:HB2	2:D:85:TYR:CE1	2.54	0.43
1:A:118:GLU:H	1:A:118:GLU:HG2	1.62	0.43
2:D:5:GLN:H	2:D:5:GLN:CD	2.21	0.42
2:B:74:ALA:HB2	2:B:85:TYR:CE1	2.54	0.42
1:A:136:LEU:CD1	1:C:0:ACE:H1	2.47	0.42
2:D:57:ASN:ND2	2:D:59:LYS:H	2.18	0.41
2:D:57:ASN:ND2	2:D:60:ILE:H	2.19	0.41
1:A:11:VAL:HG21	1:A:130:PHE:HB2	2.03	0.41
1:C:0:ACE:H2	1:C:1:THR:HA	1.41	0.41
2:D:18:LEU:CD2	2:D:23:ILE:HG13	2.50	0.41
2:D:26:LYS:NZ	4:D:148:HOH:O	2.54	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:37:TRP:O	2:B:40:ARG:HB3	2.22	0.40
2:D:18:LEU:HD23	2:D:18:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/144 (99%)	140 (99%)	2 (1%)	0	100	100
1	C	142/144 (99%)	140 (99%)	2 (1%)	0	100	100
2	B	144/146 (99%)	142 (99%)	2 (1%)	0	100	100
2	D	144/146 (99%)	143 (99%)	1 (1%)	0	100	100
All	All	572/580 (99%)	565 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/117 (100%)	110 (94%)	7 (6%)	19	6
1	C	117/117 (100%)	108 (92%)	9 (8%)	13	3
2	B	119/119 (100%)	112 (94%)	7 (6%)	19	6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	119/119 (100%)	107 (90%)	12 (10%)	7	1
All	All	472/472 (100%)	437 (93%)	35 (7%)	13	3

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LYS
1	A	21	SER
1	A	88	LEU
1	A	92	LYS
1	A	101	LYS
1	A	122	ASP
1	A	125	VAL
2	B	2	GLU
2	B	40	ARG
2	B	49	SER
2	B	68	LEU
2	B	95	LYS
2	B	104	ARG
2	B	106	LEU
1	C	20	LYS
1	C	30	LEU
1	C	34	LEU
1	C	62	LYS
1	C	78	LEU
1	C	85	LEU
1	C	88	LEU
1	C	101	LYS
1	C	107	ILE
2	D	5	GLN
2	D	18	LEU
2	D	19	ASN
2	D	57	ASN
2	D	68	LEU
2	D	80	ASN
2	D	86	SER
2	D	87	GLU
2	D	118	LEU
2	D	125	GLU
2	D	134	LEU
2	D	143	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	69	HIS
2	B	127	GLN
1	C	60	HIS
1	C	99	ASN
1	C	105	HIS
2	D	5	GLN
2	D	17	ASN
2	D	19	ASN
2	D	57	ASN
2	D	63	HIS
2	D	80	ASN
2	D	102	ASN
2	D	127	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HEM	A	144	1	41,50,50	1.86	7 (17%)	45,82,82	1.48	6 (13%)
3	HEM	C	144	1	41,50,50	1.83	5 (12%)	45,82,82	1.40	6 (13%)
3	HEM	D	147	2	41,50,50	1.93	8 (19%)	45,82,82	1.75	8 (17%)
3	HEM	B	147	2	41,50,50	1.80	8 (19%)	45,82,82	1.77	14 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	144	1	-	0/12/54/54	-
3	HEM	C	144	1	-	2/12/54/54	-
3	HEM	D	147	2	-	2/12/54/54	-
3	HEM	B	147	2	-	0/12/54/54	-

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	147	HEM	C3D-C2D	7.51	1.52	1.36
3	A	144	HEM	C3D-C2D	7.03	1.51	1.36
3	C	144	HEM	C3D-C2D	6.89	1.51	1.36
3	B	147	HEM	C3D-C2D	5.83	1.49	1.36
3	D	147	HEM	C3C-C2C	-5.11	1.33	1.40
3	B	147	HEM	C3C-C2C	-4.95	1.33	1.40
3	C	144	HEM	C3C-C2C	-4.31	1.34	1.40
3	A	144	HEM	C3C-C2C	-3.96	1.34	1.40
3	C	144	HEM	C3C-CAC	3.78	1.55	1.47
3	A	144	HEM	C3C-CAC	3.66	1.55	1.47
3	D	147	HEM	FE-ND	3.24	2.12	1.96
3	D	147	HEM	C3C-CAC	3.04	1.54	1.47
3	A	144	HEM	CAB-C3B	2.91	1.55	1.47
3	B	147	HEM	CAA-C2A	2.83	1.56	1.52
3	B	147	HEM	FE-NB	2.80	2.10	1.96
3	C	144	HEM	CAB-C3B	2.68	1.54	1.47
3	D	147	HEM	CAB-C3B	2.59	1.54	1.47
3	B	147	HEM	C3C-CAC	2.55	1.53	1.47
3	C	144	HEM	C3B-C2B	-2.47	1.32	1.37
3	A	144	HEM	CAA-C2A	2.47	1.55	1.52
3	B	147	HEM	CMB-C2B	2.41	1.55	1.50
3	B	147	HEM	C1B-NB	-2.32	1.36	1.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	147	HEM	CMB-C2B	2.27	1.55	1.50
3	B	147	HEM	CAB-C3B	2.26	1.53	1.47
3	A	144	HEM	FE-ND	2.07	2.07	1.96
3	D	147	HEM	CMD-C2D	2.03	1.55	1.50
3	D	147	HEM	CAA-C2A	2.01	1.55	1.52
3	A	144	HEM	C2C-C1C	2.01	1.47	1.42

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	147	HEM	C4D-ND-C1D	4.35	109.57	105.07
3	B	147	HEM	C4D-ND-C1D	4.29	109.51	105.07
3	B	147	HEM	CMA-C3A-C4A	-4.09	122.18	128.46
3	D	147	HEM	CAD-CBD-CGD	-3.92	105.17	113.60
3	A	144	HEM	CMA-C3A-C4A	-3.82	122.59	128.46
3	B	147	HEM	CHC-C4B-NB	3.51	128.24	124.43
3	B	147	HEM	C1B-NB-C4B	3.50	108.69	105.07
3	A	144	HEM	CBD-CAD-C3D	-3.46	103.02	112.63
3	D	147	HEM	CMA-C3A-C4A	-3.31	123.38	128.46
3	A	144	HEM	C4D-ND-C1D	3.30	108.48	105.07
3	D	147	HEM	C3B-C2B-C1B	3.14	108.82	106.49
3	C	144	HEM	C4D-ND-C1D	3.08	108.25	105.07
3	C	144	HEM	CMA-C3A-C4A	-3.07	123.75	128.46
3	B	147	HEM	CBA-CAA-C2A	-3.02	107.46	112.62
3	D	147	HEM	C1D-C2D-C3D	-2.97	103.84	106.96
3	D	147	HEM	C4C-CHD-C1D	2.93	126.43	122.56
3	B	147	HEM	C2C-C3C-C4C	2.81	108.86	106.90
3	C	144	HEM	CBD-CAD-C3D	-2.62	105.36	112.63
3	B	147	HEM	CHA-C4D-ND	2.49	127.45	124.38
3	A	144	HEM	CMA-C3A-C2A	2.41	129.49	124.94
3	A	144	HEM	CAD-C3D-C4D	2.41	128.86	124.66
3	B	147	HEM	C4B-CHC-C1C	2.32	125.61	122.56
3	C	144	HEM	CHD-C1D-ND	2.30	126.93	124.43
3	C	144	HEM	C4A-C3A-C2A	2.28	108.58	107.00
3	B	147	HEM	C4A-C3A-C2A	2.26	108.57	107.00
3	B	147	HEM	CBD-CAD-C3D	-2.22	106.47	112.63
3	B	147	HEM	CMA-C3A-C2A	2.22	129.12	124.94
3	A	144	HEM	C3B-C2B-C1B	2.20	108.12	106.49
3	D	147	HEM	CMC-C2C-C3C	2.18	128.75	124.68
3	D	147	HEM	C1B-NB-C4B	2.15	107.29	105.07
3	B	147	HEM	CAD-CBD-CGD	-2.14	109.00	113.60
3	B	147	HEM	CHD-C1D-ND	2.06	126.67	124.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	147	HEM	CAD-C3D-C4D	2.05	128.24	124.66
3	C	144	HEM	CBA-CAA-C2A	-2.05	109.12	112.62

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	144	HEM	CAA-CBA-CGA-O1A
3	C	144	HEM	CAA-CBA-CGA-O2A
3	D	147	HEM	CAD-CBD-CGD-O1D
3	D	147	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.


4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	144	HEM	1	0
3	C	144	HEM	2	0
3	D	147	HEM	1	0
3	B	147	HEM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.