



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 12:16 PM JST

PDB ID : 7V4X  
Title : Structure of cyclohexanone monooxygenase mutant from *Acinetobacter calcoaceticus*  
Authors : Wu, Y.; Yu, H.  
Deposited on : 2021-08-15  
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4468 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative flavin-binding monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	529	4204	2679	706	800	19	0	0	0

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	4	GLN	LYS	conflict	UNP A0A0A8XFY0
A	246	TYR	PHE	engineered mutation	UNP A0A0A8XFY0
A	326	CYS	LYS	engineered mutation	UNP A0A0A8XFY0
A	386	SER	ASN	engineered mutation	UNP A0A0A8XFY0
A	388	LYS	ILE	engineered mutation	UNP A0A0A8XFY0
A	390	ILE	MET	engineered mutation	UNP A0A0A8XFY0
A	426	PHE	LEU	engineered mutation	UNP A0A0A8XFY0
A	432	LEU	PHE	engineered mutation	UNP A0A0A8XFY0
A	433	ALA	THR	engineered mutation	UNP A0A0A8XFY0
A	435	SER	LEU	engineered mutation	UNP A0A0A8XFY0
A	438	ILE	SER	engineered mutation	UNP A0A0A8XFY0
A	488	LYS	GLU	engineered mutation	UNP A0A0A8XFY0
A	489	CYS	SER	engineered mutation	UNP A0A0A8XFY0
A	490	ARG	TRP	engineered mutation	UNP A0A0A8XFY0
A	505	LEU	PHE	engineered mutation	UNP A0A0A8XFY0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	48	21	7	17	3	0	0


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	163	Total 163	O 163	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative flavin-binding monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.59Å 53.00Å 100.49Å 90.00° 96.64° 90.00°	Depositor
Resolution (Å)	46.81 – 2.33 46.81 – 2.34	Depositor EDS
% Data completeness (in resolution range)	90.7 (46.81-2.33) 90.7 (46.81-2.34)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.53 (at 2.34Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, $R_{free}$	0.187 , 0.235 0.186 , 0.235	Depositor DCC
$R_{free}$ test set	1101 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.1	Xtrriage
Anisotropy	0.090	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 40.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4468	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.26	0/4298	0.49	1/5815 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	323	LEU	CA-CB-CG	5.48	127.91	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4204	0	4123	47	0
2	A	53	0	31	0	0
3	A	48	0	25	3	0
4	A	163	0	0	5	0
All	All	4468	0	4179	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:ASP:HB2	3:A:602:NAP:H72N	1.28	0.98
1:A:381:ASP:OD1	4:A:701:HOH:O	1.98	0.81
1:A:313:PHE:O	4:A:703:HOH:O	2.04	0.74
1:A:206:GLN:NE2	4:A:705:HOH:O	2.22	0.72
1:A:357:LYS:NZ	1:A:368:GLU:OE2	2.21	0.72
1:A:96:ASP:OD1	1:A:101:ARG:NH1	2.24	0.68
1:A:150:PRO:HB2	1:A:152:ILE:HG13	1.82	0.61
1:A:79:ARG:HD3	4:A:702:HOH:O	2.04	0.57
1:A:7:PHE:HA	1:A:32:LYS:HD2	1.85	0.57
1:A:395:LYS:O	1:A:398:ILE:HG22	2.05	0.56
1:A:208:SER:OG	3:A:602:NAP:O3X	2.24	0.55
1:A:71:LEU:HD21	1:A:94:VAL:HG22	1.90	0.53
1:A:484:PHE:HB3	1:A:491:ILE:HD13	1.91	0.52
1:A:57:ASP:HB2	3:A:602:NAP:N7N	2.11	0.49
1:A:49:ASN:O	1:A:84:GLN:HG3	2.11	0.49
1:A:40:THR:HG22	1:A:107:GLU:HG3	1.95	0.49
1:A:298:GLN:NE2	1:A:323:LEU:O	2.44	0.49
1:A:45:THR:HA	1:A:144:LEU:HD22	1.95	0.49
1:A:88:LEU:HG	1:A:92:LYS:HE3	1.95	0.49
1:A:114:TYR:CE1	1:A:393:ARG:HG3	2.49	0.48
1:A:120:SER:OG	1:A:460:GLN:OE1	2.27	0.48
1:A:294:ASN:OD1	1:A:298:GLN:NE2	2.45	0.47
1:A:500:LYS:HA	1:A:500:LYS:HD3	1.36	0.47
1:A:501:ASN:ND2	4:A:720:HOH:O	2.46	0.47
1:A:145:ALA:HA	1:A:146:ALA:HA	1.58	0.46
1:A:458:ILE:HD11	1:A:461:ILE:HG12	1.97	0.46
1:A:496:VAL:HB	1:A:500:LYS:HB2	1.97	0.46
1:A:288:ALA:HB1	1:A:484:PHE:HE2	1.80	0.45
1:A:460:GLN:HB3	1:A:531:LYS:HB3	1.98	0.45
1:A:82:ILE:HG12	1:A:87:VAL:HG23	1.98	0.45
1:A:302:LYS:HG2	1:A:319:THR:HG23	1.99	0.44
1:A:433:ALA:HB3	1:A:438:ILE:HD11	2.00	0.43
1:A:206:GLN:O	1:A:206:GLN:HG3	2.18	0.43
1:A:309:VAL:HB	1:A:315:ALA:HB2	2.01	0.43
1:A:431:PRO:HA	1:A:506:TYR:O	2.19	0.42
1:A:326:CYS:O	1:A:327:ARG:HB3	2.19	0.42
1:A:148:ASN:ND2	1:A:378:THR:O	2.49	0.42
1:A:496:VAL:HG21	1:A:500:LYS:HG2	2.00	0.42
1:A:254:PRO:HB2	1:A:257:SER:HB3	2.02	0.42
1:A:258:VAL:HB	1:A:262:GLU:HB3	2.00	0.41
1:A:356:ILE:HG12	1:A:361:VAL:HG22	2.01	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:357:LYS:HD2	1:A:362:VAL:HG21	2.03	0.41
1:A:280:MET:HE3	1:A:325:ALA:O	2.21	0.41
1:A:161:GLU:HB2	1:A:373:MET:HG3	2.03	0.40
1:A:227:LYS:HD3	1:A:227:LYS:HA	1.77	0.40
1:A:323:LEU:O	1:A:324:TYR:HD1	2.04	0.40
1:A:485:ALA:O	1:A:486:LYS:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	527/542 (97%)	499 (95%)	22 (4%)	6 (1%)	<b>14</b> <b>13</b>

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	323	LEU
1	A	493	GLY
1	A	499	LYS
1	A	486	LYS
1	A	485	ALA
1	A	327	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	453/466 (97%)	431 (95%)	22 (5%)	25 31

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	VAL
1	A	41	GLU
1	A	120	SER
1	A	121	PHE
1	A	151	LYS
1	A	167	ARG
1	A	175	SER
1	A	206	GLN
1	A	208	SER
1	A	223	THR
1	A	232	TYR
1	A	249	ASN
1	A	289	VAL
1	A	378	THR
1	A	386	SER
1	A	454	GLU
1	A	458	ILE
1	A	469	ASP
1	A	476	SER
1	A	486	LYS
1	A	488	LYS
1	A	500	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	523	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	601	-	53,58,58	3.81	17 (32%)	68,89,89	1.46	7 (10%)
3	NAP	A	602	-	45,52,52	2.31	9 (20%)	56,80,80	1.39	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	7/30/50/50	0/6/6/6
3	NAP	A	602	-	-	2/31/67/67	0/5/5/5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C8A-N7A	-15.02	1.07	1.34
2	A	601	FAD	C4A-N3A	-11.06	1.20	1.35
2	A	601	FAD	C2A-N3A	-9.47	1.16	1.32
2	A	601	FAD	C5A-C4A	8.80	1.64	1.40
3	A	602	NAP	O4D-C1D	7.90	1.52	1.41
3	A	602	NAP	O4B-C1B	6.54	1.50	1.41
2	A	601	FAD	O4B-C1B	6.20	1.49	1.41
2	A	601	FAD	C5A-N7A	-6.02	1.17	1.39
2	A	601	FAD	C2B-C1B	-5.50	1.45	1.53
3	A	602	NAP	C7N-N7N	5.39	1.43	1.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C2-N1	4.80	1.48	1.36
3	A	602	NAP	C2D-C1D	-4.66	1.46	1.53
2	A	601	FAD	C2A-N1A	-4.40	1.25	1.33
2	A	601	FAD	C6A-C5A	3.96	1.58	1.43
2	A	601	FAD	C8M-C8	3.80	1.58	1.51
3	A	602	NAP	C3N-C7N	3.60	1.56	1.50
2	A	601	FAD	C5X-N5	3.44	1.46	1.39
2	A	601	FAD	C6A-N6A	3.05	1.45	1.34
3	A	602	NAP	C2N-N1N	3.00	1.38	1.35
3	A	602	NAP	C3B-C2B	-2.92	1.46	1.52
3	A	602	NAP	C2D-C3D	-2.91	1.45	1.53
2	A	601	FAD	C2-N3	2.57	1.45	1.39
2	A	601	FAD	C3B-C4B	-2.34	1.47	1.53
2	A	601	FAD	C9-C9A	2.20	1.43	1.39
3	A	602	NAP	O4D-C4D	2.16	1.49	1.45
2	A	601	FAD	C2B-C3B	-2.04	1.47	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	C4A-C5A-N7A	-7.37	101.72	109.40
3	A	602	NAP	N3A-C2A-N1A	-5.52	120.04	128.68
3	A	602	NAP	C3D-C2D-C1D	4.67	108.01	100.98
2	A	601	FAD	C4-N3-C2	-3.21	119.71	125.64
3	A	602	NAP	C2D-C3D-C4D	2.85	108.18	102.64
2	A	601	FAD	C4X-C4-N3	2.79	120.28	113.19
2	A	601	FAD	O4-C4-C4X	-2.57	119.78	126.60
2	A	601	FAD	P-O3P-PA	-2.56	124.03	132.83
3	A	602	NAP	C6N-N1N-C2N	-2.41	119.78	121.97
3	A	602	NAP	PN-O3-PA	-2.38	124.68	132.83
2	A	601	FAD	C4X-C10-N10	2.30	119.85	116.48
3	A	602	NAP	C2N-C3N-C4N	2.20	120.75	118.26
2	A	601	FAD	C4X-C10-N1	-2.04	120.01	124.73
3	A	602	NAP	C4A-C5A-N7A	-2.00	107.31	109.40

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	FAD	O4B-C4B-C5B-O5B
2	A	601	FAD	N10-C1'-C2'-O2'
2	A	601	FAD	N10-C1'-C2'-C3'

Continued on next page...

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	601	FAD	C3B-C4B-C5B-O5B
2	A	601	FAD	C2'-C3'-C4'-C5'
3	A	602	NAP	O4B-C4B-C5B-O5B
2	A	601	FAD	O3'-C3'-C4'-C5'
3	A	602	NAP	C2B-O2B-P2B-O2X
2	A	601	FAD	O3'-C3'-C4'-O4'


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	NAP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	529/542 (97%)	-0.15	17 (3%) 47 58	14, 31, 60, 87	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	500	LYS	7.3
1	A	497	PRO	6.6
1	A	494	ALA	5.2
1	A	499	LYS	5.2
1	A	323	LEU	5.0
1	A	498	GLY	3.5
1	A	495	ASN	3.0
1	A	324	TYR	2.9
1	A	493	GLY	2.8
1	A	492	PHE	2.7
1	A	119	ASN	2.7
1	A	490	ARG	2.6
1	A	4	GLN	2.5
1	A	496	VAL	2.4
1	A	151	LYS	2.2
1	A	501	ASN	2.2
1	A	488	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	A	601	53/53	0.96	0.13	17,29,38,41	0
3	NAP	A	602	48/48	0.96	0.12	25,38,59,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.