

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 4, 2024 – 02:06 PM EDT

PDB ID : 8V8K

Title: Crystal Structure of Nanobody NbE

Authors: Koehl, A.; Manglik, A.; Yu, J.; Kumar, A.; Zhang, X.; Martin, C.; Raia, P.;

Steyaert, J.; Ballet, S.; Boland, A.; Stoeber, M.

Deposited on : 2023-12-05

Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467 Xtriage (Phenix) : 1.20.1

EDS : 3.0

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.002 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.38.3

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}(\mathring{A}))$
$R_{free}$	164625	1268 (2.88-2.84)
Clashscore	180529	1351 (2.88-2.84)
Ramachandran outliers	177936	1318 (2.88-2.84)
Sidechain outliers	177891	1319 (2.88-2.84)
RSRZ outliers	164620	1269 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	123	91%		8% •
			31%		
1	В	123	75%	21%	•
1	С	123	22% 85%	9%	7%



## 2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Nanobody NbE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	123	Total	С	N	O	S	0	0	0
1	Λ	120	938	589	157	188	4	0	U	0
1	B	118	Total	С	N	О	S	0	0	0
1	Ъ	110	904	570	150	180	4	U	U	U
1	С	115	Total	С	N	О	S	0	0	0
1		110	859	541	142	172	4	U	U	U



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants	76.56Å 76.56Å 298.60Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.37 - 2.85	Depositor
Resolution (A)	44.37 - 2.85	EDS
% Data completeness	98.7 (44.37-2.85)	Depositor
(in resolution range)	97.0 (44.37-2.85)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.59 (at 2.86Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
D D.	0.286 , 0.331	Depositor
$R, R_{free}$	0.283 , $0.323$	DCC
$R_{free}$ test set	11691 reflections $(10.03\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.7	Xtriage
Anisotropy	0.377	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34, 72.2	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.47, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	2701	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.27	0/958	0.52	0/1295	
1	В	0.26	0/923	0.54	0/1247	
1	С	0.26	0/877	0.52	0/1188	
All	All	0.26	0/2758	0.52	0/3730	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	938	0	901	7	0
1	В	904	0	863	19	0
1	С	859	0	790	10	0
All	All	2701	0	2554	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$	
1:B:88:LYS:HG3	1:B:90:GLU:HG2	1.76	0.67	

Continued on next page...



Continued from previous page...

Continued from pre		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)
1:B:104:GLY:HA3	1:B:109:LYS:HD3	1.79	0.64
1:B:84:MET:HB3	1:B:87:LEU:HD21	1.80	0.63
1:B:68:ARG:HE	1:B:86:SER:HB3	1.64	0.60
1:B:109:LYS:HG2	1:C:115:TRP:CH2	2.35	0.60
1:A:33:ILE:HG21	1:A:36:MET:HG3	1.83	0.59
1:B:109:LYS:HG2	1:C:115:TRP:HH2	1.71	0.55
1:B:106:TYR:O	1:C:115:TRP:NE1	2.39	0.54
1:B:30:ILE:HD11	1:B:80:VAL:HG22	1.89	0.53
1:B:62:ALA:O	1:B:66:LYS:HG3	2.11	0.51
1:C:56:GLY:H	1:C:73:ARG:HH22	1.60	0.49
1:B:40:ARG:HD3	1:B:93:ALA:HB3	1.95	0.48
1:C:20:LEU:HB2	1:C:84:MET:HE2	1.96	0.47
1:B:40:ARG:HG3	1:B:95:TYR:CG	2.49	0.47
1:A:84:MET:HB3	1:A:87:LEU:HD21	1.98	0.46
1:B:33:ILE:HG12	1:B:36:MET:HG3	1.99	0.45
1:B:21:ARG:HD3	1:B:81:TYR:CD1	2.53	0.44
1:C:35:THR:HB	1:C:100:LYS:HB3	1.99	0.44
1:A:14:VAL:HG11	1:A:87:LEU:HD12	2.00	0.44
1:B:68:ARG:NH2	1:B:85:ASN:O	2.51	0.43
1:C:56:GLY:H	1:C:73:ARG:NH2	2.16	0.43
1:B:5:GLN:OE1	1:B:27:SER:HA	2.18	0.42
1:C:38:TRP:O	1:C:50:VAL:HG22	2.19	0.42
1:B:33:ILE:HD12	1:B:33:ILE:HA	1.83	0.42
1:B:109:LYS:HD3	1:B:109:LYS:HA	1.81	0.42
1:B:61:TYR:CZ	1:B:71:ILE:HG22	2.55	0.42
1:A:100:LYS:HG3	1:A:113:ASP:OD1	2.19	0.42
1:C:84:MET:HB3	1:C:87:LEU:HD21	2.02	0.42
1:B:109:LYS:HG2	1:C:115:TRP:CZ3	2.55	0.41
1:A:35:THR:HB	1:A:100:LYS:HB3	2.02	0.41
1:A:54:THR:O	1:A:73:ARG:NH2	2.54	0.41
1:A:66:LYS:HB2	1:A:66:LYS:HE3	1.83	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	121/123 (98%)	119 (98%)	2 (2%)	0	100	100
1	В	114/123 (93%)	110 (96%)	4 (4%)	0	100	100
1	С	111/123 (90%)	105 (95%)	5 (4%)	1 (1%)	14	29
All	All	346/369 (94%)	334 (96%)	11 (3%)	1 (0%)	37	55

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	30	ILE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	100/100 (100%)	99 (99%)	1 (1%)	73	87
1	В	96/100 (96%)	95 (99%)	1 (1%)	73	87
1	С	86/100 (86%)	86 (100%)	0	100	100
All	All	282/300 (94%)	280 (99%)	2 (1%)	81	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	ILE
1	В	50	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

There are no ligands in this entry.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	123/123 (100%)	0.11	2 (1%) 70 67	50, 67, 96, 130	0
1	В	118/123 (95%)	1.43	38 (32%) 1 1	56, 107, 142, 185	0
1	С	115/123 (93%)	1.35	27 (23%) 2 2	54, 137, 169, 217	0
All	All	356/369 (96%)	0.94	67 (18%) 4 4	50, 103, 158, 217	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	122	THR	5.9
1	В	42	ALA	5.4
1	С	6	LEU	5.1
1	С	4	VAL	5.1
1	С	120	PRO	5.1
1	В	14	VAL	4.4
1	В	95	TYR	4.1
1	В	91	ASP	4.0
1	С	23	SER	4.0
1	В	22	LEU	3.6
1	В	43	PRO	3.5
1	В	121	VAL	3.2
1	В	82	LEU	3.1
1	С	97	CYS	3.1
1	В	89	PRO	3.0
1	С	101	TYR	3.0
1	С	25	ALA	3.0
1	В	84	MET	2.9
1	В	71	ILE	2.9
1	С	14	VAL	2.9
1	В	120	PRO	2.9
1	С	121	VAL	2.8
1	С	71	ILE	2.8

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Res	Type	RSRZ
1	В	119	THR	2.8
1	В	12	GLY	2.7
1	В	65	VAL	2.7
1	В	125	SER	2.7
1	В	75	ASN	2.7
1	В	62	ALA	2.7
1	С	36	MET	2.6
1	A	125	SER	2.6
1	В	87	LEU	2.6
1	С	22	LEU	2.5
1	С	80	VAL	2.5
1	С	24	CYS	2.5
1	В	85	ASN	2.5
1	В	57	GLY	2.4
1	С	30	ILE	2.4
1	С	92	THR	2.4
1	В	61	TYR	2.4
1	С	15	GLN	2.3
1	В	50	VAL	2.3
1	В	21	ARG	2.3
1	С	43	PRO	2.3
1	С	114	TYR	2.3
1	В	64	SER	2.3
1	В	51	ALA	2.3
1	С	20	LEU	2.2
1	В	68	ARG	2.2
1	В	38	TRP	2.2
1	В	45	LYS	2.2
1	С	13	LEU	2.2
1	В	70	THR	2.2
1	С	72	SER	2.1
1	В	20	LEU	2.1
1	С	82	LEU	2.1
1	В	60	ASN	2.1
1	В	55	SER	2.1
1	В	107	PHE	2.1
1	С	117	LYS	2.1
1	В	49	LEU	2.1
1	В	106	TYR	2.1
1	A	3	GLN	2.0
1	В	102	TYR	2.0
1	В	90	GLU	2.0

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	С	102	TYR	2.0
1	С	50	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.

### 6.5 Other polymers (i)

There are no such residues in this entry.

