

Full wwPDB EM Validation Report (i)

Nov 23, 2022 – 05:00 PM JST

PDB ID : 7V9L

EMDB ID : EMD-31824

Title : Cryo-EM structure of the SV1-Gs complex.

Authors: Cong, Z.T.; Zhou, F.L.; Zhang, C.; Zou, X.Y.; Zhang, H.B.; Wang, Y.Z.; Zhou,

Q.T.; Cai, X.Q.; Liu, Q.F.; Li, J.; Shao, L.J.; Mao, C.Y.; Wang, X.; Wu, J.H.; Xia, T.; Zhao, L.H.; Jiang, H.L.; Zhang, Y.; Xu, H.E.; Cheng, X.; Yang, D.H.;

Wang, M.W.

Deposited on : 2021-08-26

Resolution : 2.60 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM~structures} \ (\#{ m Entries})$	
Clashscore	158937	4297	
Ramachandran outliers	154571	4023	
Sidechain outliers	154315	3826	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	360	<u>·</u>	60%		6%		34%
2	R	522	5%	43%	6%		51%	
3	Y	71	11%		79%			• 18%
4	N	140	•		79%			11% 10%
5	В	371	•		83%			9% 8%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8047 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	237	Total 1964	C 1243	N 352	O 361	S 8	0	0

• Molecule 2 is a protein called GHRH receptor splice variant 1,GHRH receptor splice variant 1,GHRH receptor splice variant 1,SV1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
2	R	257	Total 2064	C 1377	N 336	O 344	S 7	0	0

• Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	${f Atoms}$				AltConf	Trace	
3	V	58	Total	С	N	О	S	0	0
3	1	90	445	278	78	86	3	U	U

• Molecule 4 is a protein called Nanobody 35.

Mol	Chain	Residues	Atoms				AltConf	Trace	
4	N	126	Total 961	C 599	N 168	O 188	S 6	0	0

• Molecule 5 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	В	341	Total 2613	C 1611	N 470	O 511	S 21	0	0

There are 32 discrepancies between the modelled and reference sequences:




Chain	Residue	Modelled	Actual	Comment	Reference
В	-4	MET	-	initiating methionine	UNP P54311
В	-3	GLY	-	expression tag	UNP P54311
В	-2	SER	-	expression tag	UNP P54311
В	-1	LEU	-	expression tag	UNP P54311
В	0	LEU	-	expression tag	UNP P54311
В	1	GLN	-	expression tag	UNP P54311
В	341	GLY	-	expression tag	UNP P54311
В	342	SER	-	expression tag	UNP P54311
В	343	SER	-	expression tag	UNP P54311
В	344	GLY	-	expression tag	UNP P54311
В	345	GLY	-	expression tag	UNP P54311
В	346	GLY	-	expression tag	UNP P54311
В	347	GLY	-	expression tag	UNP P54311
В	348	SER	-	expression tag	UNP P54311
В	349	GLY	-	expression tag	UNP P54311
В	350	GLY	-	expression tag	UNP P54311
В	351	GLY	-	expression tag	UNP P54311
В	352	GLY	-	expression tag	UNP P54311
В	353	SER	-	expression tag	UNP P54311
В	354	SER	-	expression tag	UNP P54311
В	355	GLY	-	expression tag	UNP P54311
В	356	VAL	-	expression tag	UNP P54311
В	357	SER	-	expression tag	UNP P54311
В	358	GLY	-	expression tag	UNP P54311
В	359	TRP	-	expression tag	UNP P54311
В	360	ARG	-	expression tag	UNP P54311
В	361	LEU	-	expression tag	UNP P54311
В	362	PHE	-	expression tag	UNP P54311
В	363	LYS	-	expression tag	UNP P54311
В	364	LYS	-	expression tag	UNP P54311
В	365	ILE	-	expression tag	UNP P54311
В	366	SER	-	expression tag	UNP P54311




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Guanine nucleotide-binding protein G(s) subunit alpha isoforms short




• Molecule 2: GHRH receptor splice variant 1,GHRH receptor splice variant 1,GHRH receptor splice variant 1,SV1





A SN TTYR A LIA A LIA A LIA A LIA A SN TTRE TT

• Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2




• Molecule 4: Nanobody 35





 \bullet Molecule 5: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	377241	Depositor
Resolution determination method	DIFFRACTION PATTERN/LAYERLINES	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	80	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.785	Depositor
Minimum map value	-0.021	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.022	Depositor
Recommended contour level	0.0644	Depositor
Map size (Å)	234.07999, 234.07999	wwPDB
Map dimensions	224, 224, 224	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.045, 1.045, 1.045	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.29	0/2003	0.47	0/2697	
2	R	0.29	0/2124	0.45	0/2896	
3	Y	0.25	0/451	0.42	0/609	
4	N	0.32	0/981	0.51	0/1329	
5	В	0.30	0/2660	0.53	0/3606	
All	All	0.29	0/8219	0.49	0/11137	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1964	0	1948	11	0
2	R	2064	0	2096	19	0
3	Y	445	0	454	2	0
4	N	961	0	932	11	0
5	В	2613	0	2516	20	0
All	All	8047	0	7946	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
2:R:130:HIS:HB3	2:R:133:PHE:N	<u>distance (Å)</u> 1.77	overlap (Å)
1:A:228:ARG:NH2	5:B:186:ASP:OD1		0.97
	2:R:133:PHE:N	2.12	0.82
2:R:130:HIS:CB		2.47	0.77
5:B:22:ARG:NH1	5:B:258:ASP:O	2.28	0.65
4:N:117:TYR:OH	5:B:228:ASP:OD1	2.19	0.60
5:B:286:LEU:HD22	5:B:327:VAL:HG21	1.87	0.57
2:R:97:ARG:NH2	2:R:318:GLU:OE2	2.38	0.55
5:B:51:LEU:HB2	5:B:336:LEU:HB3	1.89	0.53
5:B:30:LEU:HD23	5:B:262:MET:HB2	1.91	0.52
2:R:119:ASP:OD1	2:R:120:ALA:N	2.43	0.52
5:B:166:CYS:HB2	5:B:180:PHE:HB2	1.93	0.50
1:A:232:ARG:NH2	5:B:204:CYS:HB3	2.27	0.50
5:B:112:VAL:HG23	5:B:124:TYR:HB2	1.94	0.50
4:N:12:VAL:HG11	4:N:86:LEU:HD13	1.94	0.50
1:A:239:ASN:HA	1:A:285:ILE:HD11	1.93	0.49
4:N:64:VAL:HG13	4:N:68:PHE:HB2	1.93	0.49
5:B:163:ASP:O	5:B:165:THR:N	2.45	0.49
2:R:93:LEU:HD13	2:R:319:VAL:HG22	1.95	0.48
4:N:24:ALA:HB1	4:N:27:PHE:CZ	2.48	0.48
4:N:67:ARG:HH22	4:N:90:ASP:CG	2.16	0.48
1:A:206:GLY:HA2	1:A:226:ALA:HB3	1.95	0.48
5:B:130:GLU:HG3	5:B:132:ASN:H	1.78	0.48
1:A:270:LEU:HD21	1:A:344:GLU:HB3	1.94	0.48
5:B:123:ILE:HD11	5:B:139:LEU:HD11	1.96	0.47
4:N:68:PHE:CD1	4:N:83:MET:HA	2.50	0.47
2:R:130:HIS:CG	2:R:133:PHE:N	2.82	0.46
2:R:306:PHE:CZ	2:R:310:ILE:HD11	2.50	0.46
1:A:282:LEU:HA	1:A:285:ILE:HD12	1.96	0.46
5:B:262:MET:SD	5:B:302:ALA:HB2	2.56	0.46
1:A:224:VAL:HG21	1:A:234:TRP:CZ3	2.51	0.46
1:A:45:LEU:HD21	1:A:53:LYS:HB2	1.98	0.45
2:R:164:ASN:ND2	2:R:235:LEU:HB3	2.31	0.45
4:N:88:PRO:O	4:N:91:THR:HG22	2.16	0.45
5:B:139:LEU:HD13	5:B:169:TRP:CG	2.53	0.44
2:R:133:PHE:O	2:R:133:PHE:CG	2.70	0.44
4:N:48:VAL:HG13	4:N:64:VAL:HG21	2.00	0.44
5:B:124:TYR:CE1	5:B:135:VAL:HG22	2.52	0.44
2:R:102:THR:O	2:R:106:THR:HG23	2.18	0.44
5:B:254:ASP:OD1	5:B:256:ARG:N	2.50	0.44
1:A:64:HIS:HB2	1:A:212:PHE:HB2		
		2.00	0.43
4:N:83:MET:HE1	4:N:124:VAL:HG21	2.00	0.43
2:R:139:LYS:HD3	2:R:209:TRP:CZ2	2.54	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		${ m distance}({ m \AA})$	overlap (Å)
5:B:254:ASP:HB2	5:B:261:LEU:HD11	1.99	0.43
2:R:83:ALA:HB1	2:R:313:CYS:SG	2.58	0.43
2:R:304:GLN:O	2:R:308:VAL:HG23	2.19	0.43
2:R:127:ASP:OD1	2:R:128:THR:N	2.51	0.42
1:A:238:PHE:O	1:A:241:VAL:HG22	2.20	0.42
3:Y:48:ASP:HB2	5:B:281:SER:HB3	2.01	0.42
4:N:29:PHE:O	4:N:72:ARG:NH2	2.52	0.42
1:A:215:ASP:OD1	1:A:373:ARG:NH2	2.46	0.42
2:R:267:THR:O	2:R:271:ILE:HG12	2.20	0.41
2:R:294:LEU:HB2	2:R:295:PRO:HD3	2.01	0.41
2:R:273:LEU:HB2	2:R:312:TYR:CE2	2.55	0.41
2:R:94:HIS:HB3	2:R:99:TYR:OH	2.20	0.41
5:B:310:GLY:O	5:B:337:LYS:NZ	2.31	0.41
4:N:68:PHE:CD1	4:N:68:PHE:N	2.89	0.40
2:R:163:LEU:HD23	2:R:239:ILE:HD13	2.03	0.40
3:Y:60:PRO:HD2	5:B:325:MET:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	233/360 (65%)	224 (96%)	9 (4%)	0	100 100
2	R	251/522~(48%)	242 (96%)	9 (4%)	0	100 100
3	Y	56/71 (79%)	56 (100%)	0	0	100 100
4	N	124/140 (89%)	117 (94%)	7 (6%)	0	100 100
5	В	339/371 (91%)	326 (96%)	13 (4%)	0	100 100
All	All	1003/1464 (68%)	965 (96%)	38 (4%)	0	100 100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	P	erce	\mathbf{ntiles}
1	A	214/314~(68%)	212 (99%)	2 (1%)		78	91
2	R	224/444~(50%)	224 (100%)	0		100	100
3	Y	47/58 (81%)	47 (100%)	0		100	100
4	N	104/116~(90%)	104 (100%)	0		100	100
5	В	282/302 (93%)	280 (99%)	2 (1%)		84	94
All	All	871/1234 (71%)	867 (100%)	4 (0%)		89	96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	220	HIS
1	A	280	ARG
5	В	105	TYR
5	В	234	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
5	В	239	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-31824. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map






Z Index: 112



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0644. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $40~\mathrm{nm^3}$; this corresponds to an approximate mass of $36~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



^{*}Reported resolution corresponds to spatial frequency of 0.385 $\rm \AA^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-31824 and PDB model 7V9L. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0644 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0644).



9.4 Atom inclusion (i)



At the recommended contour level, 89% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0644) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8112	0.5050
A	0.8110	0.5130
В	0.8719	0.5560
N	0.8412	0.5320
R	0.7427	0.4310
Y	0.7100	0.4610



