



# wwPDB X-ray Structure Validation Summary Report

Dec 6, 2023 – 02:40 am GMT

PDB ID : 2VBO  
Title : Molecular basis of human XPC gene recognition and cleavage by engineered homing endonuclease heterodimers  
Authors : Redondo, P.; Prieto, J.; Munoz, I.G.; Alibes, A.; Stricher, F.; Serrano, L.; Arnould, S.; Perez, C.; Cabaniols, J.P.; Duchateau, P.; Paques, F.; Blanco, F.J.; Montoya, G.  
Deposited on : 2007-09-14  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: red;"></div> </div> <p style="text-align: center;">83% 12% . .</p>
2	B	153	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 1%; height: 10px; background-color: red;"></div> </div> <p style="text-align: center;">3% 81% 15% . . .</p>
3	C	24	<div style="display: flex; align-items: center;"> <div style="width: 21%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 58%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: orange;"></div> </div> <p style="text-align: center;">21% 58% 21%</p>
4	E	24	<div style="display: flex; align-items: center;"> <div style="width: 58%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 38%; height: 10px; background-color: orange;"></div> </div> <p style="text-align: center;">. 58% 38%</p>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3788 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA ENDONUCLEASE I-CREI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	153	1225	786	210	228	1	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	SER	TYR	conflict	UNP P05725
A	38	ARG	GLN	conflict	UNP P05725
A	42	THR	ALA	conflict	UNP P05725
A	70	SER	ARG	conflict	UNP P05725
A	75	ASN	ASP	conflict	UNP P05725
A	110	GLU	TRP	conflict	UNP P05725
A	111	GLN	ARG	conflict	UNP P05725

- Molecule 2 is a protein called DNA ENDONUCLEASE I-CREI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	152	1232	794	207	230	1	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	28	GLU	LYS	conflict	UNP P05725
B	38	ARG	GLN	conflict	UNP P05725
B	40	LYS	SER	conflict	UNP P05725
B	42	THR	ALA	conflict	UNP P05725
B	44	LYS	GLN	conflict	UNP P05725
B	68	SER	ARG	conflict	UNP P05725
B	70	SER	ARG	conflict	UNP P05725
B	75	ASN	ASP	conflict	UNP P05725
B	110	GLU	TRP	conflict	UNP P05725

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	111	GLN	ARG	conflict	UNP P05725

- Molecule 3 is a DNA chain called 5'-D(\*TP\*CP\*TP\*GP\*CP\*CP\*TP\*TP\*TP\*TP \*TP\*T  
P\*GP\*AP\*AP\*GP\*GP\*AP\*TP\*CP\*CP\*TP\*AP\*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	24	493	236	97	137	23	0	0	0

- Molecule 4 is a DNA chain called 5'-D(\*TP\*TP\*AP\*GP\*GP\*AP\*TP\*CP\*CP\*TP \*TP\*C  
P\*AP\*AP\*AP\*AP\*AP\*AP\*GP\*GP\*CP\*AP\*GP\*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	E	24	485	235	80	147	23	0	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total 3	Ca 3	0	0
5	B	1	Total 1	Ca 1	0	0



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	139	Total 139	O 139	0	0
6	B	112	Total 112	O 112	0	0
6	C	47	Total 47	O 47	0	0
6	E	51	Total 51	O 51	0	0



### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA ENDONUCLEASE I-CREI




- Molecule 2: DNA ENDONUCLEASE I-CREI



- Molecule 3: 5'-D(\*TP\*CP\*TP\*GP\*CP\*CP\*TP\*TP\*TP\*TP \*TP\*TP\*GP\*AP\*AP\*GP\*GP\*AP\*TP\*CP\*CP\*TP\*AP\*A)-3'



- Molecule 4: 5'-D(\*TP\*TP\*AP\*GP\*GP\*AP\*TP\*CP\*CP\*TP \*TP\*CP\*AP\*AP\*AP\*AP\*AP\*AP\*GP\*GP\*CP\*AP\*GP\*A)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.99Å 67.95Å 76.63Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	39.47 – 1.80 39.47 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.47-1.80) 100.0 (39.47-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.150 , 0.225 0.152 , 0.223	Depositor DCC
$R_{free}$ test set	2206 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.3	Xtrriage
Anisotropy	0.127	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3788	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/1247	0.75	2/1684 (0.1%)
2	B	0.56	0/1255	0.87	2/1693 (0.1%)
3	C	1.02	0/555	2.24	34/855 (4.0%)
4	E	0.92	0/541	2.30	38/833 (4.6%)
All	All	0.71	0/3598	1.47	76/5065 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	2
All	All	0	4

There are no bond length outliers.

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	153	ASP	CA-C-O	19.59	161.24	120.10
4	E	3	DT	P-O3'-C3'	12.67	134.90	119.70
4	E	16	DG	O4'-C1'-N9	12.28	116.60	108.00
3	C	13	DA	N1-C2-N3	-11.60	123.50	129.30
4	E	13	DG	O4'-C1'-N9	-11.51	99.95	108.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	149	ARG	Peptide
1	A	150	ALA	Peptide
2	B	150	ALA	Peptide
2	B	152	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1225	0	1255	34	0
2	B	1232	0	1267	26	0
3	C	493	0	271	15	0
4	E	485	0	276	29	0
5	A	3	0	0	0	0
5	B	1	0	0	0	0
6	A	139	0	0	17	0
6	B	112	0	0	9	0
6	C	47	0	0	4	0
6	E	51	0	0	3	0
All	All	3788	0	3069	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:4:DG:H2'	6:E:2006:HOH:O	1.28	1.29
6:A:2022:HOH:O	4:E:2:DC:H5	1.22	1.21
1:A:33:SER:HB3	6:A:2024:HOH:O	0.97	1.15
4:E:5:DC:H6	4:E:5:DC:H5'	1.11	1.13
3:C:23:DG:C1'	6:C:2047:HOH:O	2.03	1.04

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/153 (99%)	143 (95%)	7 (5%)	1 (1%)	22	10
2	B	150/153 (98%)	143 (95%)	5 (3%)	2 (1%)	12	3
All	All	301/306 (98%)	286 (95%)	12 (4%)	3 (1%)	15	5

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	150	ALA
2	B	149	ARG
2	B	150	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/140 (98%)	131 (96%)	6 (4%)	28	14
2	B	139/140 (99%)	135 (97%)	4 (3%)	42	29
All	All	276/280 (99%)	266 (96%)	10 (4%)	35	20

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	151	VAL
2	B	152	LEU
2	B	153	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	123	LEU
1	A	152	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	ASN
2	B	75	ASN
2	B	103	ASN
2	B	92	GLN
1	A	92	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/153 (100%)	-0.27	2 (1%) 77 74	14, 23, 45, 61	0
2	B	152/153 (99%)	-0.15	5 (3%) 46 40	14, 24, 49, 66	0
3	C	24/24 (100%)	-0.50	0 100 100	16, 28, 49, 50	0
4	E	24/24 (100%)	-0.38	0 100 100	15, 31, 49, 51	0
All	All	353/354 (99%)	-0.24	7 (1%) 65 61	14, 24, 49, 66	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	2	ASN	4.0
1	A	119	PRO	2.4
2	B	33	TYR	2.4
1	A	151	VAL	2.3
2	B	31	GLN	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CA	A	1154	1/1	1.00	0.10	15,15,15,15	0
5	CA	A	1155	1/1	1.00	0.09	15,15,15,15	0
5	CA	A	1156	1/1	1.00	0.06	21,21,21,21	0
5	CA	B	1154	1/1	1.00	0.06	24,24,24,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.